

## TABLE OF CONTENTS

	Page
<b>ACKNOWLEDGMENT</b>	iii
<b>ABSTRACT</b> (in English)	v
<b>ABSTRACT</b> (in Thai)	ix
<b>TABLE OF CONTENTS</b>	xiii
<b>LIST OF ILLUSTRATIONS</b>	xviii
<b>LIST OF TABLES</b>	xxvi
 <b>CHAPTER 1 INTRODUCTION</b>	
1.1 Background and problem	1
1.2 Previous investigation	3
1.2.1 Tectonic setting of Tertiary basins in Thailand	3
1.2.2 Stratigraphic Sequence	3
1.2.3 Organic petrography	6
1.3 Objective	10
1.4 Scope, Planning and Methodology	10
1.5 Education advantages	11
 <b>CHAPTER 2 GEOLOGY OF STUDY ARAES</b>	
2.1 Geological setting of Fang basin	12
2.2 Geological setting of Na Hong basin	19
2.3 Geological setting of Li basin	22
2.4 Geological setting of Mae Sot basin	27
2.5 Geological setting of Phitsanulok basin	31
2.6 Geological setting of Suphan Buri basin	38
 <b>CHAPTER 3 SAMPLE COLLECTION AND ANALYTICAL METHODS</b>	
3.1 Sample collection and preparation	43
3.1.1 Sample collection	43
3.1.2 Preparation of samples	46

3.2	Methodology	55
3.2.1	Total organic carbon (TOC) analysis, Total carbon (TC) analysis and Total sulfur (TS) analysis	55
3.2.1.1.	LECO™ CS200 for TC, TOC and TS analysis	55
3.2.1.2	Infrared Radiation, Absorption and Detection	55
3.2.1.3	Sampling preparation for TOC analysis	57
3.2.1.4	TOC, TC and TS analytical procedure	58
3.2.2	Rock-Eval pyrolysis	58
3.2.2.1	Rock-Eval Pyrolysis procedure	61
3.2.3	Gas Chromatography and Gas Chromatography- Mass Spectrometry	61
3.2.3.1	Sample selection	61
3.2.3.2	Extraction and separation	61
3.2.3.3	Gas chromatography	62
3.2.3.4	Gas chromatography-mass spectrometry	63
3.2.4	Petrography	63
3.2.4.1.	Organic petrography	63
3.2.4.2	Vitrinite reflectance measurements	65
3.2.4.3	Sample selection	66
3.2.4.4	Sample preparation	66

## CHAPTER 4 RESULTS AND INTERPRETATION

4.1	Data screening	67
4.1.1	Results of Fang basin samples	70
4.1.2	Results of Na Hong basin	74
4.1.3	Results of Ban Pa Kha sub-basin, Li basin	74
4.1.4	Results of Mae Sot basin	79
4.1.5	Results of P-SK well, Phitsanulok basin	79
4.1.6	Results of Suphanburi Basin	87
4.2	<i>n</i> -Alkane distribution by gas chromatography	104
4.2.1	Fang-MS well, Fang basin	107
4.2.2	Na Hong basin	108

4.2.3	Li basin	108
4.2.4	Mae Sot basin	108
4.2.5	P-SK well, Phitsanulok basin	108
4.2.6	SP1 and SP2 wells, Suphanburi basin	108
4.3	Biomarkers parameters: Gas Chromatography-Mass Spectrometer	109
4.3.1	Fang-MS well, Fang basin	111
4.3.2	Na Hong basin	111
4.3.3	Li basin	111
4.3.4	Mae Sot basin	114
4.3.5	P-SK, Phitsanulok basin	114
4.3.6	SP1 and SP2 wells, Suphanburi basin	114
4.4	Organic Petrographic Results	114
4.4.1	Fang-MS well, Fang basin	115
4.4.2	Na Hong basin	122
4.4.3	Li basin	122
4.4.4	Mae Sot basin	133
4.4.5	P-SK, Phitsanulok basin	139
4.4.6	SP1 and SP2 wells, Suphanburi basin	139


## CHAPTER 5 DISCUSSION

5.1	Depositional environment of organic mater	159
5.2	Source rock quality and petroleum generation	170
5.3	Source rock thermal maturity	180

## CHEPTEP 6 CONCLUSION

6.1	Fang-MS well, Fang basin	190
6.2	Na Hong basin	191
6.3	Li basin	191
6.4	Mae Sot basin	192
6.5	Phitsanulok basin	192
6.6	Suphanburi basin	193

<b>REFERENCES</b>	197
<b>APPENDICES</b>	208
APPENDIX A	209
APPENDIX B	221
APPENDIX C	236
<b>CURRICULUM VITAE</b>	304



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่  
Copyright© by Chiang Mai University  
All rights reserved

## LIST OF FIGURES

<b>Figure</b>	<b>Page</b>
1.1 Cenozoic basins in Thailand. The study areas are in the blocks	2
1.2 Structural map of Northern Thailand showing relationship between conjugate strike-slip faults and the development of N-S trending pull-apart basins.	5
1.3 Generalized stratigraphy of Cenozoic basins in Northern Thailand.	7
1.4 Structural map of Central Thailand showing relationship between conjugate strike-slip faults and the development of N-S trending pull-apart basins.	8
1.5 Generalized stratigraphy of Cenozoic basins in Central Thailand.	9
2.1 Geologic map of Fang basin.	14
2.2 Generalised stratigraphic column for Fang Basin.	17
2.3 Geological cross-section through the Mae Soon structure	18
2.4 Geologic map of Na Hong basin	20
2.5 Stratigraphic succession along quarry wall of coal mine of Na Hong basin.	21
2.6 Geology of Li basin.	23
2.7 Schematic stratigraphy of Ban Pa Kha Formation at Ban Pa Kha Coalfield.	25
2.8 Geological map of Mae Sot basin.	28
2.9 Bouguer anomaly map of Mae Sot basin.	29
2.10 The Teriary stratigraphy colum of Mae Sot basin.	30
2.11 The Phitsanulok Basin is located in central Thailand and location of Sirikit oil field.	31
2.12 Geological map of Phitsanulok basin	33
2.13 Stratigraphic cross-section of the Phitsanulok Basin.	34
2.14 The stratigraphy of the Phitsanulok basin.	36
2.15 The Suphanburi and Kamphaeng Saen basin are located in central Thailand. Location of U Thong and Kamphaeng Saen oil fields and are also shown.	39
2.16 Geological map of Suphanburi basin showing location of U Thong oilfield	40
2.17 The stratigraphy and petroleum system of the Suphanburi basin.	41

3.1	Generalised stratigraphic column for well FANG, showing stratigraphic levels and sample numbers of samples collected.	44
3.2	Generalised stratigraphic column for well PH, showing stratigraphic levels and sample numbers of samples collected.	47
3.3	Generalised stratigraphic column for well SP1, showing stratigraphic levels and sample numbers of samples collected.	50
3.4	Generalised stratigraphic column for well SP2, showing stratigraphic levels and sample numbers of samples collected.	53
3.5	LECO CS-20 gas flow diagram.	56
3.6	Principal of the Rock-Eval pyrolysis device.	59
3.7	Cycle of analysis and example of record.	60
4.1	The plots of TOC content, TC content and TS content against depth show variation of screening data and source rock quality of Fang-MS well.	73
4.2	The plots of $S_1$ , $S_2$ and HI against depth showing variation of screening data, source rock quality and petroleum generation potential of Fang-MS well	75
4.3	The plots of $T_{max}$ and PI against depth showing variation of screening data and maturation of source rock of Fang-MS well	76
4.4	The plots of TOC content, TC content and TS content against depth show variation in screening data and source rock quality of P-SK well	84
4.5	The plots of $S_1$ , $S_2$ and HI against depth show variation of screening data, source rock quality and petroleum generation potential of P-SK well	85
4.6	The plots of $T_{max}$ and PI against depth show variation of screening data and maturation of source rock of P-SK well	86
4.7	The plots of TOC content, TC content and TS content against depth show variation of screening data and source rock quality of SP1 well	93
4.8	The plots of $S_1$ , $S_2$ and HI against depth show variation of screening data, source rock quality and petroleum generation potential of SP1 well	94

4.9	The plots of $T_{\max}$ and PI against depth show variation of screening data and maturation of source rock of SP1 well	95
4.10	The plots of TOC content, TC content and TS content against depth show variation of screening data and source rock quality of SP2 well	96
4.11	The plots of $S_1$ , $S_2$ and HI against depth show variation of screening data, source rock quality and petroleum generation potential of SP2 well	102
4.12	The plots of $T_{\max}$ and PI against depth show variation of screening data and maturation of source rock of SP2 well	103
4.13	Photomicrographs of filamentous lamalginite (FLA), exsudatinite (Ex) surrounding quartz grain (Qtz) and framboidal pyrite (Py) of sample 11859 (588.30 m) from Fang-MS well in cross polarize light (A) and in fluorescence-inducing blue light (B).	117
4.14	Photomicrographs of <i>Botryococcus</i> -type telalginite (Bo) and huminite Fragments (Hum) in groundmass of yellowish brown fluorescing amorphous organic matter (AOM) and liptodetrinite (Lip) of sample 11859 (588.30 m) from Fang-MS well in cross polarize light (A) and in fluorescence-inducing blue light (B).	118
4.15	Photomicrographs of gelinite (Gel), cutinite (Cu), resinite (Re) and exsudatinite (Ex) intruded into cleats of gelinite of sample 11861 (618.70 m) from Fang-MS well in white light (A) and in fluorescence-inducing blue light (B).	119
4.16	Photomicrograph of resinite (Re) and lamalginite (Lam) display yellowish orange color in fluorescence-inducing blue light of sample 11871 (786.40 m) from Fang-MS well.	120
4.17	Photomicrograph of association of disc-shaped <i>Botryococcus</i> -type Telalginite (Bo) and lamalginite (Lam) in groundmass of weakly brownish fluorescing amorphous organic matter and liptodetrinite in fluorescence-inducing blue light of sample 11871 (786.40 m) from Fang-MS well.	120

- 4.18 Photomicrograph of exsudatinite (Ex) surrounding quartz (Qtz) grains and displays greenish yellow color and *Botryococcus?* (Bo?) in fluorescence-inducing blue light of sample 11876 (879.30 m) from Fang-MS well. 121
- 4.19 Photomicrograph of framboidal pyrite (Py) in humic coal (Hum) in white light of sample 11882 (984.5 m) from Fang-MS well. 121
- 4.20 Photomicrographs of association of disc-shaped *Botryococcus*-type telalginite (Bo) and lamalginite (Lam) in groundmass of fluorescing amorphous organic matter and liptodetrinite of sample 14714 from Na Hong basin in cross polarize light (A) and in fluorescence-inducing blue light (B). 124
- 4.21 Photomicrographs of association of disc-shaped *Botryococcus*-type telalginite (Bo) and lamalginite (Lam) in groundmass of fluorescing amorphous organic matter and liptodetrinite; exsudatinite (Ex) filled in pore of fusinite (Fu) layer of sample 14714 from Na Hong basin in white light (A) and in fluorescence-inducing blue light (B). 125
- 4.22 Photomicrographs of resinite (Re), sporinite (Sp), liptodetrinite (Lip) and cutinite (Cu) in densinite (Den) and gelinite (Gel) groundmass in sample 14709 from Na Hong basin in white light (A) and in fluorescence-inducing blue light (B). 126
- 4.23 Photomicrographs of resinite (Re), sporinite (Sp), exsudatinite (Ex) and framboidal pyrite (Py) in densinite (Den) groundmass in sample 14709 from Na Hong basin in white light (A) and in fluorescence-inducing blue light (B) 127
- 4.24 Photomicrographs of lamalginite (Lam), exsudatinite (Ex) and cutinite (Cu) in dentinite groundmass of sample 14719 from Na Hong basin, in white light (A) and white color in fluorescence-inducing blue light (B). 128
- 4.25 Photomicrographs of lamalginite (Lam), sporinite (Sp), exsudatinite (Ex) filled in pore of fusinite (Fu) and framboidal pyrite (Py) in densinite groundmass of sample 14712 from Na Hong basin in white light (A) and in fluorescence-inducing blue light (B). 129



- 4.26 Photomicrographs of lamalginite (Lam) and resinite (Re) in densinite (Den) groundmass of sample 14719 from Na Hong basin in white light (A) and in fluorescence-inducing blue light (B). 130
- 4.27 Photomicrographs of exsudatinite (Ex) in huminite (Hum) and association of disc-shaped *Botryococcus*-type telalginite (Bo), lamalginite and pyrite (Py) in groundmass of fluorescing amorphous organic matter and liptodetrinite of sample 14690 Ban Pa Kha subbasin, Li basin in white light (A) in fluorescence-inducing blue light (B). 131
- 4.28 Photomicrographs of gelinite huminite (Hum) with exsudatinite in cleates and association of disc-shaped *Botryococcus*-type telalginite and lamalginite in groundmass of fluorescing amorphous organic matter and liptodetrinite of sample 14693 from Li basin in white light (A) in fluorescence-inducing blue light (B). 132
- 4.29 Photomicrographs of framboidal pyrite (Py) and association of disc-shaped *Botryococcus*-type telalginite (Bo) and lamalginite in groundmass of fluorescing amorphous organic matter and liptodetrinite of sample 14693 from Li basin in cross polarize light (A) and in fluorescence-inducing blue light (B). 134
- 4.30 Photomicrographs of framboidal pyrite (Py) (white in polarize light and black in blue light) in homogeneous AOM which considered mainly to be derived from alginite of sample 14700 from Mae Sot basin in cross polarize light (A); in fluorescence-inducing blue light (B). 135
- 4.31 Photomicrographs of homogeneous AOM which considered mainly to be derived from alginite of sample 14700 from Mae Sot basin in cross polarize light (A); in fluorescence-inducing blue light (B). 136
- 4.32 Photomicrographs of disc-shaped *Botryococcus*-type telalginite (Bo) in pyrite (Py) rich liptodetrinite groundmass of sample 14702 from Mae Sot basin in cross polarize light (A) in fluorescence-inducing blue light (B). 137

- 4.32 Photomicrographs of disc-shaped *Botryococcus*-type telalginite (B) in liptodetrinite groundmass and exsudatinite (Ex) filled in pore of funginite (Fun) of sample 14702 from Mae Sot basin in white light (A); in fluorescence-inducing blue light (B). 138
- 4.34 Photomicrographs of disc-shaped *Botryococcus*-type telalginite (Bo) in liptodetrinite and fluorescing amorphous organic matter groundmass of sample 14742 from P-SK well in cross polarize light (A); in fluorescence-inducing blue light (B). 141
- 4.35 Photomicrographs of filamentous lamalginite (Lam) in liptodetrinite and fluorescing amorphous organic matter groundmass of sample 14745 from P-SK well in fluorescence-inducing blue light (A and B). Greenish fluorescing lines are exsudationite (Ex) of low number carbon chain. 142
- 4.36 Photomicrograph of cutinite (Cu) and resinite (Re) in the groundmass of *Botryococcus* algae of sample 14755 from P-SK well in fluorescence-inducing blue light. 143
- 4.37 Photomicrograph of resinite (Re) and exsudatinite (Ex) expelled into cleats of huminite of sample 14737 from P-SK well in fluorescence-inducing blue light. 143
- 4.38 Photomicrographs of exsudatinite (Ex) intruded into cleats of huminite of sample 14728 from P-SK well in cross polarize light (A); in fluorescence-inducing blue light (B). 144
- 4.39 Photomicrographs of exsudatinite (Ex) intruded into cleats of huminite, and pyrite (Py) of sample 14748 from P-SK well in white light (A); in fluorescence-inducing blue light (B). 145
- 4.40 Photomicrograph of huminite (Hum), funginite (Fun) and semifusinite (Semi-fu) in groundmass of Liptodetrinite sample 14755 from P-SK well in white light. 146
- 4.41 Photomicrograph of inertinite? (In?) and huminite (Hum) of sample 14737 from P-SK well in white light. 146

- 4.42 Photomicrographs of exsudatinite (Ex) surrounded of quartz grains (Qtz) of sample 14728 from P-SK well in cross polarize light (A); in fluorescence-inducing blue light (B). 148
- 4.43 Photomicrograph of disc-shaped *Botryococcus*-type telalginite (Bo) and pyrite (Py) in liptodetrinite (Lip) and fluorescing amorphous organic matter groundmass of sample 11725 from SP1 well in fluorescence-inducing blue light. 150
- 4.44 Photomicrograph of compacted lamalginite (Lam) and disc-shaped *Botryococcus*-type telalginite (Bo) in groundmass of fluorescing amorphous organic matter and liptodetrinite of sample 11753 from SP1 well in fluorescence-inducing blue light. The brown color of lamalginite indicated the partially expelled of hydrocarbon. 150
- 4.45 Photomicrograph of lamalginite (Lam) and exsudatinite (Ex) surrounded quartz grains (Qtz) of sample 11753 from SP1 well in fluorescence-inducing blue light. 151
- 4.46 Photomicrographs of huminite (Hum) of sample 11726 from SP1 well in white light. 151
- 4.47 Photomicrographs of non-fluorescing mineral matter (Non-fl) and association of disc-shaped *Botryococcus*-type telalginite (Bo) and lamalginite (Lam) in groundmass of fluorescing amorphous organic matter and liptodetrinite of sample 11725 from SP1 well in fluorescence-inducing blue light. 152
- 4.48 Photomicrograph of exsudatinite (Ex), resinite (Re) and non-fluorescing mineral matter (Non-fl) of sample 11743 from SP1 well in fluorescence-inducing blue light. 152
- 4.49 Photomicrographs of lamalginite (Lam) and framboidal pyrite (Py) of sample 11726 from SP1 well in white light (A); in fluorescence-inducing blue light (B). 153
- 4.50 Photomicrograph of association of disc-shaped *Botryococcus*-type telalginite (B) and lamalginite (Lam) in groundmass of fluorescing amorphous organic matter and liptodetrinite of sample 11799 from SP2 well in fluorescence-inducing blue light. 155

4.51 Photomicrograph of lamalginite (Lam) and fluorescing amorphous organic matter (Fl AOM) of sample 11802 from SP2 well in fluorescence-inducing blue light.	155
4.52 Photomicrograph of lamalginite (Lam) in groundmass of liptodetrinite and pyrite (black) of sample 11820 from SP2 well in fluorescence-inducing blue light.	156
4.53 Photomicrograph of resinite (Re) and lamalginite (Lam) of sample 11829 from SP2 well in fluorescence-inducing blue light.	156
4.54 Photomicrographs of disc-shaped <i>Botryococcus</i> -type telalginite (B), lamalginite (Lam), huminite (Hum) and framboidal pyrite (Py) of sample 11825 from SP2 well in cross polarize light (A) and in fluorescence-inducing blue light (B).	157
5.1 TOC/TS versus TOC plot showing relatively high TOC/TS ratios (~5-8) of the samples from Fang-MS well indicating mostly blackish origin	160
5.2 A Pr/nC <sub>17</sub> versus Ph/nC <sub>18</sub> plot of samples studies. The plot is used to indicate redox depositional environments	161
5.3 Triangular diagram showing interpretation of environment from C <sub>27</sub> , C <sub>28</sub> and C <sub>29</sub> regular steranes distribution	163
5.4 TOC/TS versus TOC plot showing relatively high TOC/TS ratios of the samples from Na Hong, Li and Mae Sot basins	165
5.5 TOC/TS versus TOC plot showing relatively high TOC/TS ratios of the samples from P-SK well.	169
5.6 TOC/TS versus TOC plot showing relatively high TOC/TS ratios of the samples from SP1 and SP2 well	171
5.7 (A) S <sub>2</sub> yields plot against the TOC content suggesting a fair to good source rock quality, S <sub>2</sub> and TOC cut-off values based on Peters and Moldowan (1993); (B) Hydrogen Index versus T <sub>max</sub> plot. The kerogen of Fang-MS samples falls within the areas of Type II and III and possess a potential for mixed oil/gas and oil generation	173

- 5.8 (A)  $S_2$  yields plot against the TOC content of Na Hong, Li and Mae Sot samples,  $S_2$  and TOC cut-off values based on Peters and Moldowan (1993).; (B) Hydrogen Index versus  $T_{max}$  plot of Na Hong, Li and Mae Sot samples 174
- 5.9 (A)  $S_2$  yields plot against the TOC content for P-SK samples suggesting a fair to excellent source rock quality,  $S_2$  and TOC cut-off values based on Peters and Moldowan (1993); (B) Hydrocarbon Index versus  $T_{max}$  plot. The kerogens of P-SK samples fall within the areas of Type II and III and possess a potential for mixed oil/gas and oil generation 177
- 5.10 (A)  $S_2$  yields plot against the TOC content of samples from SP1 well suggesting a poor to excellent source rock quality,  $S_2$  and TOC cut-off values based on Peters and Moldowan (1993); (B) Hydrogen Index versus  $T_{max}$  plot of samples from SP1 well. The kerogen falls within the areas of Type II and III and possess a potential for mixed oil/gas and oil generation of SP1 samples 179
- 5.11 (A)  $S_2$  yields plot against the TOC content of samples of SP2 well suggesting a poor to excellent source rock quality,  $S_2$  and TOC cut-off values based on Peters and Moldowan (1993); (B) Hydrogen Index versus  $T_{max}$  plot of samples of SP2 well. The kerogens fall within the areas of Type II and III and possess a potential for mixed oil/gas and oil generation of SP2 samples 181
- 5.12  $C_{31} 22S/(22S+22R)$  homohopane ratio versus  $C_{27} Ts/(Ts+Tm)$  ratio plot suggesting source rock maturity 182
- 5.13  $C_{29} 20S/(20S+20R)$  sterane ratio versus  $C_{29} \beta\beta/(\beta\beta+\alpha\alpha)$  sterane ratio plot suggesting source rock maturity 183
- 5.14 Vitrinite reflectance versus depth plot suggesting source sock maturity of sample from Fang-MS well, Fang basin 184
- 5.15 Vitrinite reflectance versus depth plot suggesting source sock maturity of sample from P-SK well, Phitsanulok basin 187
- 5.16 Vitrinite reflectance versus depth plot suggesting source sock maturity of sample from SP1 and SP2 wells, Suphnburi basin 188

## LIST OF TABLES

<b>Table</b>	<b>Page</b>
3.1 Kerogen classification.	64
4.1 Screening data of samples from Fang-MS well from Fang basin.	71
4.2 The screening data of Na Hong samples.	77
4.3 The screening data of Li samples	78
4.4 The screening data of Mae Sot samples..	80
4.5 Screening data of samples from P-SK well from Phitsanulok basin	82
4.6 Screening data of samples from SP1 well from Suphanburi basin.	88
4.7 Screening data of samples from SP2 well from Suphanburi basin	99
4.8 Gas chromatogram data of n-alkane hydrocarbon from Fang-MS well, Na Hong, Li, Mae Sot, P-SK well, SP1 well and SP2 well.	105
4.9 Summarized data of biomarkers from Fang, Na Hong, Li, Mae Sot, Phitsanulok, and Suphanburi basins.	112
4.10 Organic composition and vitrinite reflectance of the samples of Fang-MS well from Fang basin.	116
4.11 Organic composition and vitrinite reflectance of the samples from Na Hong, Li and Mae Sot basins.	123
4.12 Organic composition and vitrinite reflectance of the samples from P-SK well, Phitsanulok basin.	140
4.13 Organic composition and vitrinite reflectance of the samples from SP1 well, Suphanburi basin.	148
4.14 Organic composition and vitrinite reflectance of the samples from SP2 well, Suphanburi basin.	154