

TABLE OF CONTENTS

	Page
ACKNOWLEDGMENTS	ii
ABSTRACT	VII
LIST OF TABLES	XIII
LIST OF FIGURES	XVII
INTRODUCTION	1
LITERATURE REVIEW	6
Taxonomic Status	6
Soybean Stem Fly Biology	6
Status of Soybean Stem Fly as a Pest	7
Management of Soybean Stem Fly	9
Environmental Control of Soybean Stem Fly	
Using Different Ecological Conditions	10
MATERIALS AND METHODS	14
Sampling Soybean Stem Fly, <u>Melanagromyza</u>	
<u>soj</u> ae (Zehntner) on Soybean	17
Sampling adult populations	17
Sampling larvae and pupae populations	19
Predator Population Densities Assessment	20
Vegetation Diversity in Cropping Systems	21
RESULTS	
Sampling Soybean Stem Fly, <u>Melanagromyza</u>	

	XII
	Page
<u>sojae</u> (Zehntner) on Soybean	24
Sampling adult populations	24
Sampling larvae and pupae populations	27
Predator Population Density Assessment	42
Vegetation Diversity in Various Soybean Cropping Systems	81
The Effect of Cropping Systems on Crop and Weed Yields	87
DISCUSSION AND CONCLUSIONS	94
LITERATURE CITED	114
CURRICULUM VITAE	120

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่  
 Copyright© by Chiang Mai University  
 All rights reserved

## LIST OF TABLES

Table	Page
1. Mean monthly temperature, rainfall and relative humidity during the experimental period.	15
2. Mean densities of soybean stem fly, <u>Melanagromyza sojae</u> (Zhnt.) adults counted on 25 plants in various soybean cropping systems at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).	25
3. Mean densities of soybean stem fly, <u>Melanagromyza sojae</u> (Zhnt.) larvae and pupae in various soybean cropping systems at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).	28
4. Mean percent infestations by <u>Melanagromyza sojae</u> (Zhnt.) in various soybean cropping systems at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).	32
5. Mean rates of parasitization of <u>Melanagromyza sojae</u> (Zhnt.) by	

## Table

## Page

- naturally occurring parasitoid, Eurytoma sp. in various soybean cropping systems at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989). 35
6. Mean soybean stem fly, Melanagromyza sojae (Zhnt.) densities, percent infestations, and rates of parasitization by naturally occurring parasitoid, Eurytoma sp. in various soybean cropping systems at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989). 37
7. Abundance of predators in various soybean cropping systems (7 weeks after planting) at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989). 50
8. Abundance of predators in various soybean cropping systems (8 weeks after planting) at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989). 52

Table	Page
9. Abundance of predators in various soybean cropping systems (9 weeks after planting) at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).	55
10. Abundance of predators in various soybean cropping systems (10 weeks after planting) at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).	58
11. Abundance of predators in various soybean cropping systems (11 weeks after planting) at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).	61
12. Abundance of predators in various soybean cropping systems (12 weeks after planting) at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).	64
13. Mean abundance of predators in various soybean cropping systems at the Multiple Cropping Center Experimental	

## Table

## Page

	Farm, Chiang Mai (1989).	69
14.	Mean natural enemy species diversity measures in various soybean cropping systems at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).	78
15.	Vegetation diversity associated with soybean grown in monoculture and polyculture cropping systems at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).	82
16.	Percentage species composition of weeds in various soybean cropping systems at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).	85
17.	Comparison of weed biomass in various soybean cropping systems at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).	88
18.	Crop and weed yields in soybean monoculture and soybean-corn polyculture systems at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).	90

## LIST OF FIGURES

Figure	Page
<p>1. Layout of randomized complete block design for each eight treatments: Soybean monocultures: weedy throughout the season, weed-free for only two weeks after planting, weed-free for only four weeks after planting, weed-free throughout the season. Soybean/sweet corn polycultures: weed-free throughout the season, weed-free for only two weeks after planting, weed-free for only four weeks after planting, and weedy throughout the season. Each replicated four times at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989). x = soybean, o = corn.</p>	16
<p>2. The locations of four one square meter quadrats selected for sampling vegetation diversity and biomass in soybean monocultures: weedy throughout the season, weed-free for only two</p>	

## Figure

Page

weeks after planting, weed-free for only four weeks after planting, weed-free throughout the season.

Soybean/sweet corn polycultures: weed-free throughout the season, weed-free for only two weeks after planting, weed-free for only four weeks after planting, and weedy throughout the season at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

22

3. The seasonal abundance of adult soybean stem fly, Melanagromyza sojae (Zhnt.) densities in soybean monocultures: (•) weedy throughout the season, (+) weed-free for only two weeks after planting, (×) weed-free for only four weeks after planting, (□) weed-free throughout the season, and soybean/sweet corn polycultures: (X) weed-free throughout the season, (◇) weed-free for only two weeks after planting, (△) weed-free for only four weeks after planting, and (⊗) weedy



## Figure

## Page

throughout the season at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

26

4. Mean densities of soybean stem fly, Melanagromyza sojae (Zhnt.) larvae and pupae in soybean monocultures: (•) weedy throughout the season, (+) weed-free for only two weeks after planting, (×) weed-free for only four weeks after planting, (□) weed-free throughout the season, and soybean/corn polycultures: (X) weed-free throughout the season, (◇) weed-free for only two weeks after planting, (△) weed-free for only four weeks after planting, and (⌘) weedy throughout the season at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

29

5. Mean percent infestation by soybean stem fly, Melanagromyza sojae (Zhnt.) in soybean monocultures: (•) weedy throughout the season, (+)

Figure

Page

weed-free for only two weeks after planting, (X) weed-free for only four weeks after planting, (□) weed-free throughout the season, and soybean/corn polycultures: (X) weed-free throughout the season, (◇) weed-free for only two weeks after planting, (△) weed-free for only four weeks after planting, and (Z) weedy throughout the season at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

34

6. Percentage of soybean stem fly, Melanagromyza sojae (Zhnt.) larvae attacked by the naturally occurring parasitoid, Eurytoma sp. in soybean monocultures: (•) weedy throughout the season, (+) weed-free for only two weeks after planting, (X) weed-free for only four weeks after planting, (□) weed-free throughout the season, and soybean/corn polycultures: (X) weed-free throughout the season, (◇)

## Figure

Page

weed-free for only two weeks after planting, ( $\Delta$ ) weed-free for only four weeks after planting, and ( $\times$ ) weedy throughout the season at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

36

7. Relationship between soybean stem fly, Melanagromyza sojae (Zhnt.) larvae (L) and percent infestation (I) in soybean monocultures (L = .—.—., I = +—+—+) and soybean/corn polycultures (L = \*—\*—\*, I = □—□—□) weedy throughout the season at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

38

8. Relationship between soybean stem fly, Melanagromyza sojae (Zhnt.) larvae (L) and percent infestation (I) in soybean monocultures (L = .—.—., I = +—+—+) and soybean/corn polycultures (L = \*—\*—\*, I = □—□—□) weed-free for only two weeks after planting at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

38

## Figure

9. Relationship between soybean stem fly, Melanagromyza sojae (Zhnt.) larvae (L) and percent infestation (I) in soybean monocultures (L = .—.—., I = +—+—+) and soybean/corn polycultures (L = \*—\*—\*, I = □—□—□) weed-free for only four weeks after planting at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

39

10. Relationship between soybean stem fly, Melanagromyza sojae (Zhnt.) larvae (L) and percent infestation (I) in soybean monocultures (L = .—.—., I = +—+—+) and soybean/corn polycultures (L = \*—\*—\*, I = □—□—□) weed-free throughout the season at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

39

11. Relationship between soybean stem fly, Melanagromyza sojae (Zhnt.) larvae (L) and rate of parasitization (P) in soybean monocultures (L = .—.—., P = +—+—+) and soybean/corn polycultures

## Figure

Page

( L = \*—\*—\*, P = □—□—□ ) weedy throughout the season at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

40

12. Relationship between soybean stem fly, Melanagromyza sojae (Zhnt.) larvae (L) and rate of parasitization (P) in soybean monocultures (L = .—.—., P = +—+—+) and soybean/corn polycultures (L = \*—\*—\*, P = □—□—□) weed-free for only two weeks after planting at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

40

13. Relationship between soybean stem fly, Melanagromyza sojae (Zhnt.) larvae (L) and rate of parasitization (P) in soybean monocultures (L = .—.—., P = +—+—+) and soybean/corn polycultures (L = \*—\*—\*, P = □—□—□) weed-free for only four weeks after planting at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

41

## Figure

## Page

14. Relationship between soybean stem fly, Melanagromyza sojae (Zhnt.) larvae (L) and rate of parasitization (P) in soybean monocultures (L = .—.—., P = +—+—+) and soybean/corn polycultures (L = \*—\*—\*, P = □—□—□) weed-free throughout the season at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

41

15. Relationship between soybean stem fly, Melanagromyza sojae (Zhnt.) larvae (L) and adults (A) in soybean monocultures (L = .—.—., A = +—+—+) and soybean/corn polycultures (L = \*—\*—\*, A = □—□—□) weedy throughout the season at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

43

16. Relationship between soybean stem fly, Melanagromyza sojae (Zhnt.) larvae (L) and adults (A) in soybean monocultures (L = .—.—., A = +—+—+) and soybean/corn polycultures (L =

## Figure

## Page

\*—\*—\*, A = □—□—□ ) weed-free for only two weeks after planting at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

43

17. Relationship between soybean stem fly, Melanagromyza sojae (Zhnt.) larvae (L) and adults (A) in soybean monocultures (L = .—.—., A = +—+—+) and soybean/corn polycultures (L = \*—\*—\*, A = □—□—□) weed-free for only four weeks after planting at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

44

18. Relationship between soybean stem fly, Melanagromyza sojae (Zhnt.) larvae (L) and adults (A) in soybean monocultures (L = .—.—., A = +—+—+) and soybean/corn polycultures (L = \*—\*—\*, A = □—□—□) weed-free throughout the season at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

44

19. Relationship between soybean stem

## Figure

Page

fly, Melanagromyza sojae (Zhnt.) adults (A) and percent infestation (I) in soybean monocultures ( A = .—.—., I = +—+—+ ) and soybean/corn polycultures ( A = \*—\*—\*, I = □—□—□ ) weedy throughout the season at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

45

20. Relationship between soybean stem fly, Melanagromyza sojae (Zhnt.) adults (A) and percent infestation (I) in soybean monocultures ( A = .—.—., I = +—+—+ ) and soybean/corn polycultures ( A = \*—\*—\*, I = □—□—□ ) weed-free for only two weeks after planting at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

45

21. Relationship between soybean stem fly, Melanagromyza sojae (Zhnt.) adults (A) and percent infestation (I) in soybean monocultures ( A = .—.—., I = +—+—+ ) and soybean/corn



## Figure

## Page

polycultures ( A = \*—\*—\*, I = □—□—□ ) weed-free for only four weeks after planting at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

46

22. Relationship between soybean stem fly, Melanagromyza sojae (Zhnt.) adults (A) and percent infestation (I) in soybean monocultures ( A = .—.—., I = +—+—+ ) and soybean/corn polycultures ( A = \*—\*—\*, I = □—□—□ ) weed-free throughout the season at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

46

23. Relationship between soybean stem fly, Melanagromyza sojae (Zhnt.) adults (A) and rate of parasitization (P) in soybean monocultures ( A = .—.—., P = +—+—+ ) and soybean/corn polycultures ( A = \*—\*—\*, P = □—□—□ ) weedy throughout the season at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

47

## Figure

## Page

24. Relationship between soybean stem fly, Melanagromyza sojae (Zhnt.) adults (A) and rate of parasitization (P) in soybean monocultures (A = .—.—., P = +—+—+) and soybean/corn polycultures (A = \*—\*—\*, P = □—□—□) weed-free for only two weeks after planting at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

47

25. Relationship between soybean stem fly, Melanagromyza sojae (Zhnt.) adults (A) and rate of parasitization (P) in soybean monocultures (A = .—.—., P = +—+—+) and soybean/corn polycultures (A = \*—\*—\*, P = □—□—□) weed-free for only four weeks after planting at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

48

26. Relationship between soybean stem fly, Melanagromyza sojae (Zhnt.) adults (A) and rate of parasitization (P) in soybean monocultures (A = .—.—., P =

## Figure

Page

+—+—+ ) and soybean/corn polycultures (A = \*—\*—\*, P = □—□—□) weed-free throughout the season at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

48

27. Mean spider densities in soybean monocultures: (•) weedy throughout the season, (+) weed-free for only two weeks after planting, (×) weed-free for only four weeks after planting, (□) weedy throughout the season, and soybean/corn polycultures: (×) weed-free throughout the season, (◇) weed-free for only two weeks after planting, (△) weed-free for only four weeks after planting, and (X̄) weedy throughout the season at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

68

28. Mean coccinellid beetle densities in soybean monocultures: (•) weedy throughout the season, (+) weed-free for only two weeks after planting,

Figure

Page

(\*) weed-free for only four weeks after planting, (□) weed-free throughout the season, and soybean/corn polycultures: (X) weed-free throughout the season, (◇) weed-free for only two weeks after planting, (△) weed-free for only four weeks after planting, and (⊗) weedy throughout the season at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

70

29. Mean pentatomid bug, Podisus sp. densities in soybean monocultures: (\*) weedy throughout the season, (+) weed-free for only two weeks after planting, (\*) weed-free for only four weeks after planting, (□) weed-free throughout the season, and soybean/corn polycultures: (X) weed-free throughout the season, (◇) weed-free for only two weeks after after planting, (△) weed-free for only four weeks after planting, and (⊗) weedy throughout the season at the Multiple Cropping

## Figure

Page

Center Experimental Farm, Chiang Mai  
(1989).

72

30. Mean big-eyed bug, Geocoris sp. densities in soybean monocultures: (•) weedy throughout the season, (+) weed-free for only two weeks, (×) weed-free for only four weeks after planting, (□) weed-free throughout the season, and soybean/corn polycultures: (X) weed-free throughout the season, (◇) weed-free for only two weeks after planting, (△) weed-free for only four weeks after planting, and (⊗) weedy throughout the season at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).

74

31. Mean pentatomid bug, Stiretrus sp. densities in monocultures: (•) weedy throughout the the season, (+) weed-free for only two weeks after planting, (×) weed-free for only four weeks after planting, (□) weedy throughout the season, and soybean/corn

## Figure

Page

polycultures: (X) weed-free throughout the season, (◇) weed-free for only two weeks after planting, (△) weed-free for only four weeks after planting, and (⊗) weedy throughout the season at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989). 75

32. Mean syrphid densities in soybean monocultures: (\*) weedy throughout the season, (+) weed-free for only two weeks after planting, (×) weed-free for only four weeks after planting, (□) weed-free throughout the season and soybean/corn polycultures: (X) weed-free throughout the season, (◇) weed-free for only two weeks after planting, (△) weed-free for only four weeks after planting, and (⊗) weedy throughout the season at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989). 76

33. Total biomass in soybean monocultures: weedy throughout the

season, weed-free for only two weeks after planting, weed-free for only four weeks after planting, weedy throughout the season and soybean/corn polycultures: weed-free throughout the season, weed-free for only two weeks after planting, weed-free for only four weeks after planting, and weedy throughout the season at the Multiple Cropping Center Experimental Farm, Chiang Mai (1989).