

## สารบัญ

|                                      | หน้า |
|--------------------------------------|------|
| กิตติกรรมประกาศ                      | ค    |
| บทคัดย่อภาษาไทย                      | ง    |
| บทคัดย่อภาษาอังกฤษ                   | ฉ    |
| สารบัญตาราง                          | ญ    |
| สารบัญภาพ                            | ฎ    |
| สารบัญตารางภาคผนวก                   | ฐ    |
| อักษรย่อ                             | ท    |
| บทที่ 1 บทนำ                         | 1    |
| วัตถุประสงค์                         | 2    |
| บทที่ 2 การตรวจเอกสาร                | 3    |
| ลิตีต                                | 3    |
| คอเลสเตรอล                           | 7    |
| แนวทางการลดคอเลสเตรอลในเนื้อและไขไก่ | 11   |
| เยื่อใย                              | 12   |
| สมุนไพรร                             | 14   |
| โปรไบโอติก                           | 15   |
| แร่ธาตุ                              | 16   |
| กรดไขมันชนิดไม่อิ่มตัวเชิงซ้อน       | 23   |
| โคติน และโคโตซาน                     | 30   |
| บทที่ 3 อุปกรณ์ และวิธีการทดลอง      | 32   |
| อุปกรณ์ที่ใช้ในการทดลอง              | 32   |
| วิธีการทดลอง                         | 34   |
| การวิเคราะห์ข้อมูลทางสถิติ           | 39   |
| สถานที่ทำการทดลอง                    | 39   |

สารบัญ (ต่อ)

|  | หน้า |
|--|------|
| บทที่ 4 ผลการทดลอง                               | 40   |
| ส่วนประกอบของกรดไขมันในน้ำมันจากพืชที่ใช้ทดลอง   | 40   |
| การศึกษาในไก่ไข่ (การทดลองที่ 1)                 | 41   |
| สมรรถภาพการผลิตและคุณภาพไข่                      | 41   |
| คอเลสเตอรอลในเลือดและไข่                         | 41   |
| กรดไขมันในไข่                                    | 43   |
| ทองแดงในอวัยวะภายในและมูล                        | 45   |
| การศึกษาในนกกกระทาไข่ (การทดลองที่ 2)            | 47   |
| สมรรถภาพการผลิตและคุณภาพไข่                      | 47   |
| คอเลสเตอรอลในไข่                                 | 48   |
| กรดไขมันในไข่                                    | 49   |
| ทองแดงในมูล                                      | 50   |
| ต้นทุนการผลิตไข่ไก่ และไข่นกกกระทา               | 51   |
| บทที่ 5 วิจัยรณและสรุปลผลการทดลอง                | 53   |
| วิจัยรณผลการทดลอง                                | 53   |
| สรุปลผลการทดลอง                                  | 59   |
| เอกสารอ้างอิง                                    | 61   |
| ภาคผนวก  | 69   |
| ภาคผนวก ก. การวิเคราะห์ตัวอย่างในห้องปฏิบัติการ  | 69   |
| ภาคผนวก ข. ข้อมูลด้านสมรรถภาพการผลิต             | 74   |
| ภาคผนวก ค. ข้อมูลการวิเคราะห์ความแปรปรวน (ANOVA) | 98   |

สารบัญตาราง

| ตารางที่  | หน้า |
|---|------|
| 1. Cholesterol content per 100 g. of fresh egg matter in individual weeks of the laying period.   | 10   |
| 2. The relative biological value of different Cu sources in poultry.  | 18   |
| 3. Fatty acid composition of some plant oils compared with tallow.  | 29   |
| 4. Feed formulation and chemical composition of experimental laying hen diets.  | 37   |
| 5. Feed formulation and chemical composition of experimental laying Japanese quails diets.  | 39   |
| 6. Fatty acid profiles (% of total fatty acid) in palm, soybean and linseed oils which is investigated in this study.   | 40   |
| 7. Production performance of laying hens fed diets containing different sources of plant oil and various levels of copper and chromium during 168 days.   | 42   |
| 8. Serum and yolk cholesterol content of laying hens fed diets containing different sources of plant oil and various levels of copper or chromium during 168 days and after changing to the control diets for 28 day. | 43   |
| 9. Fatty acid profiles in egg yolk of laying hens fed diets containing different sources of plant oil and various levels of copper and chromium during 168 days.  | 44   |
| 10. Copper content in visceral organs of laying hens fed diets containing different sources of plant oil and various levels of copper and chromium during 168 days.   | 46   |
| 11. Copper in excreta of laying hens containing different sources of plant oil and various levels of copper and chromium during 168 days and after changing to the control diets for 28 days.                         | 47   |

## สารบัญตาราง (ต่อ)

| ตารางที่  | หน้า |
|---|------|
| 12. Production performance of Japanese quails fed diets containing different sources of plant oil and various levels of copper and chromium during 84 days.                                   | 48   |
| 13. Yolk cholesterol content of Japanese quails fed diets containing different sources of plant oil and various levels of copper and chromium during 84 days.                                 | 48   |
| 14. Fatty acid profiles (% of total fatty acid) in egg yolk of Japanese quails fed diets containing different sources of plant oil and various levels of copper or chromium during 84 days.   | 49   |
| 15. Copper content in excreta of Japanese quails fed diets containing different sources of plant oil and various levels of copper and chromium during 84 days.                                | 50   |
| 16. Cost of production of laying hens and Japanese quails fed diets containing different sources of plant oil and various levels of copper and chromium during 168 and 84 days, respectively. | 51   |

## สารบัญภาพ

| ภาพที่  | หน้า |
|---|------|
| 1. Structural formula of cholesterol                        | 7    |
| 2. Cholesterol content per egg and per 100 g of yolk        | 11   |
| 3. Metabolic pathways of essential fatty acids in mammals   | 25   |
| 4. Flax seed and oil  | 28   |
| 5. Structure of chitin                                      | 31   |
| 6. Automated chemistry analyzer (MERCK; Mega)               | 33   |
| 7. Gas chromatography (Shimadzu; GC-14B)                    | 33   |
| 8. Atomic absorption spectrophotometer (Perkin Elmer; 3100) | 33   |
| 9. The form of chromium and copper use in the experiment    | 36   |
| 10. Oils from soybean, palm and linseed                     | 36   |

## สารบัญตารางภาคผนวก

| ตารางภาคผนวก ข. ที่ (Table Appendix B.)   | หน้า |
|---|------|
| B-1 Egg production of laying hens fed diets containing different sources of plant oil and various levels of copper and chromium during 168 days.                                      | 74   |
| B-2 Feed intake of laying hens fed diets containing different sources of plant oil and various levels of copper and chromium during 168 days.   | 74   |
| B-3 Feed used (kg)/doz. egg of laying hens fed diets containing different sources of plant oil and various levels of copper and chromium during 168 days.                             | 75   |
| B-4 Feed used (kg)/egg of laying hens fed diets containing different sources of plant oil and various levels of copper and chromium during 168 days.                                  | 75   |
| B-5 Body weight gain (g) of laying hens fed diets containing different sources of plant oil and various levels of copper and chromium during 168 days.                                | 76   |
| B-6 Egg weight (g) of laying hens fed diets containing different sources of plant oil and various levels of copper and chromium during 168 days.                                      | 76   |
| B-7 Egg specific gravity of laying hens fed diets containing different sources of plant oil and various levels of copper and chromium which added in to control diet during 168 days. | 77   |
| B-8 Haugh unit of laying hens fed diets containing different sources of plant oil and various levels of copper and chromium during 168 days.  | 77   |
| B-9 Yolk color (score) of laying hens fed diets containing different sources of plant oil and various levels of copper and chromium during 168 days.                                  | 78   |

## สารบัญตารางภาคผนวก (ต่อ)

| <b>Table Appendix B.</b>  | <b>หน้า</b> |
|---|-------------|
| B-10 Egg shell thickness (mm) of laying hens fed diets containing different sources of plant oil and various levels of copper and chromium during 168 days.   | 78          |
| B-11 Egg grading (%) of laying hens fed diets containing different sources of plant oil and various levels of copper and chromium during 168 days.  | 79          |
| B-12 Serum cholesterol (mg/dl) of laying hens fed diets containing different sources of plant oil and various levels of copper and chromium during 168 days.  | 80          |
| B-13 Cholesterol in yolk and egg of laying hens fed diets containing different sources of plant oil and various levels of copper and chromium during 168 days.  | 81          |
| B-14 Fatty acid profiles (% of total fatty acid) in egg yolk of laying hens fed diets containing different sources of plant oil and various levels of copper and chromium during 168 days.                      | 82          |
| B-15 Fatty acid profiles (g/100g yolk) in egg yolk of laying hens fed diets containing different sources of plant oil and various levels of copper and chromium during 168 days.                                | 84          |
| B-16 Fatty acid profiles (g/egg) in egg yolk of laying hens fed diets containing different sources of plant oil and various levels of copper and chromium during 168 days.                                      | 86          |
| B-17 Copper content in excreta of laying hens fed diets containing different sources of plant oil and various levels of copper and chromium during 168 days and after changing to the control diets for 28 day. | 88          |
| B-18 Copper content (mg/kg) in visceral organs of laying hens fed diets containing different sources of plant oil and various levels of copper and chromium during 168 days.                                    | 89          |
| B-19 Egg production (%) of Japanese quails fed diets containing different sources of plant oil and various levels of copper and chromium during 84 days.  | 89          |



สารบัญตารางภาคผนวก (ต่อ)

| <b>Table Appendix B.</b>  | <b>หน้า</b> |
|---|-------------|
| B-20 Feed intake (g/day) of Japanese quails fed diets containing different sources of plant oil and various levels of copper and chromium during 84 days.   | 90          |
| B-21 Feed used (kg)/100 eggs of Japanese quails fed diets containing different sources of plant oil and various levels of copper and chromium during 84 days.                                       | 90          |
| B-22 Feed used (kg)/kg. egg of Japanese quails fed diets containing different sources of plant oil and various levels of copper and chromium during 84 days.  | 90          |
| B-23 Egg weight (g) of Japanese quails fed diets containing different sources of plant oil and various levels of copper and chromium during 84 days.  | 91          |
| B-24 Yolk color (score) of Japanese quails fed diets containing different sources of plant oil and various levels of copper and chromium during 84 days.  | 91          |
| B-25 Egg shell thickness (mm) of Japanese quails fed diets containing different sources of plant oil and various levels of copper and chromium during 84 days.                                      | 91          |
| B-26 Cholesterol in yolk and egg of Japanese quails fed diets containing different sources of plant oil and various levels of copper and chromium during 84 days.                                   | 92          |
| B-27 Copper content in excreta of Japanese quails fed diets containing different sources of plant oil and various levels of copper and chromium during 84 days.                                     | 92          |
| B-28 Retention time, area peak and fatty acid profiles in egg yolk of Japanese quails fed diets containing different sources of plant oil and various levels of copper and chromium during 84 days. | 93          |
| B-29 Cholesterol content in laying hen and Japanese quails egg yolk from various report sources during 1980-2003.   | 94          |
| B-30 Fatty acid profiles (% of total fatty acid) of laying hens egg yolk from various reports.  | 95          |



## สารบัญตารางภาคผนวก (ต่อ)

| <b>Table Appendix B.</b>   | <b>หน้า</b> |
|--|-------------|
| B-31 Fatty acid profiles (% of total fatty acid) of Japanese quails egg yolk from various reports. | 96          |
| B-32 Fatty acids profiles of some plant oils compared with tallow (g/100g fat).                    | 97          |
| <br>   |             |
| <b>Table Appendix C.</b>   |             |
| C-1 ANOVA: Egg production of laying hens for 168 days.   | 98          |
| C-2 ANOVA: Feed intake of laying hens for 168 days.  | 98          |
| C-3 ANOVA: Feed/doz. egg of laying hens for 168 days.  | 98          |
| C-4 ANOVA: Feed/kg egg of laying hens for 168 days.  | 98          |
| C-5 ANOVA: Body weight gain of laying hens for 168 days.   | 98          |
| C-6 ANOVA: Egg weight of laying hens for 168 days.   | 99          |
| C-7 ANOVA: Haugh unit of laying hens for 168 days.   | 99          |
| C-8 ANOVA: Egg specific gravity of laying hens for 168 days.                                       | 99          |
| C-9 ANOVA: Egg shell thickness of laying hens for 168 days.  | 99          |
| C-10 ANOVA: Yolk color of laying hens for 168 days.  | 99          |
| C-11 ANOVA: Egg grading No. 5 (< 50 g.) of laying hens for 168 days.                               | 100         |
| C-12 ANOVA: Egg grading No. 4 (51-55 g.) of laying hens for 168 days.                              | 100         |
| C-13 ANOVA: Egg grading No. 3 (56-60 g.) of laying hens for 168 days.                              | 100         |
| C-14 ANOVA: Egg grading No. 2 (61-65 g.) of laying hens for 168 days.                              | 100         |
| C-15 ANOVA: Egg grading No. 1 (66-70 g.) of laying hens for 168 days.                              | 100         |
| C-16 ANOVA: Egg grading No. 0 (>70 g.) of laying hens for 168 days.                                | 101         |
| C-17 ANOVA: Serum cholesterol of laying hens for 168 days.   | 101         |
| C-18 ANOVA: Serum cholesterol of laying hens (after changing to the control diet)                  | 101         |
| C-19 ANOVA: Yolk cholesterol of laying hens for 168 days. (mg/g yolk)                              | 101         |
| C-20 ANOVA: Yolk cholesterol of laying hens (after changing to the control diet, mg/g yolk)        | 101         |
| C-21 ANOVA: Yolk cholesterol of laying hens for 168 days. (mg/egg)                                 | 102         |

## สารบัญตารางภาคผนวก (ต่อ)

| Table Appendix C.  | หน้า |
|--|------|
| C-22 ANOVA: Yolk cholesterol of laying hens (after changing to the control diet, mg/egg) | 102  |
| C-23 ANOVA: Fatty acid profile of laying hens (C16:0, % of total fatty acid)             | 102  |
| C-24 ANOVA: Fatty acid profile of laying hens (C16:1, % of total fatty acid)             | 102  |
| C-25 ANOVA: Fatty acid profile of laying hens (C18:0, % of total fatty acid)             | 102  |
| C-26 ANOVA: Fatty acid profile of laying hens (C18:1, % of total fatty acid)             | 103  |
| C-27 ANOVA: Fatty acid profile of laying hens (C18:2, % of total fatty acid)             | 103  |
| C-28 ANOVA: Fatty acid profile of laying hens (C18:3, % of total fatty acid)             | 103  |
| C-29 ANOVA: Fatty acid profile of laying hens (C20:5, % of total fatty acid)             | 103  |
| C-30 ANOVA: Fatty acid profile of laying hens (C22:6, % of total fatty acid)             | 104  |
| C-31 ANOVA: Fatty acid profile of laying hens (SFA, % of total fatty acid)               | 104  |
| C-32 ANOVA: Fatty acid profile of laying hens (UFA, % of total fatty acid)               | 104  |
| C-33 ANOVA: Fatty acid profile of laying hens (MUFA, % of total fatty acid)              | 104  |
| C-34 ANOVA: Fatty acid profile of laying hens (PUFA, % of total fatty acid)              | 105  |
| C-35 ANOVA: Fatty acid profile of laying hens (UFA/SFA, % of total fatty acid)           | 105  |
| C-36 ANOVA: Fatty acid profile of laying hens (n-3, % of total fatty acid)               | 105  |
| C-37 ANOVA: Fatty acid profile of laying hens (n-6, % of total fatty acid)               | 105  |
| C-38 ANOVA: Fatty acid profile of laying hens (n-6/n-3, % of total fatty acid)           | 106  |
| C-39 ANOVA: Fatty acid profile of laying hens (C16:0, g/100g yolk)                       | 106  |
| C-40 ANOVA: Fatty acid profile of laying hens (C16:1, g/100g yolk)                       | 106  |
| C-41 ANOVA: Fatty acid profile of laying hens (C18:0, g/100g yolk)                       | 106  |
| C-42 ANOVA: Fatty acid profile of laying hens (C18:1, g/100g yolk)                       | 106  |
| C-43 ANOVA: Fatty acid profile of laying hens (C18:2, g/100g yolk)                       | 107  |
| C-44 ANOVA: Fatty acid profile of laying hens (C18:3, g/100g yolk)                       | 107  |
| C-45 ANOVA: Fatty acid profile of laying hens (C20:5, g/100g yolk)                       | 107  |
| C-46 ANOVA: Fatty acid profile of laying hens (C22:6, g/100g yolk)                       | 107  |
| C-47 ANOVA: Fatty acid profile of laying hens (SFA, g/100g yolk)                         | 107  |
| C-48 ANOVA: Fatty acid profile of laying hens (UFA, g/100g yolk)                         | 108  |
| C-49 ANOVA: Fatty acid profile of laying hens (MUFA, g/100g yolk)                        | 108  |

สารบัญตารางภาคผนวก (ต่อ)

| Table Appendix C.  | หน้า |
|--|------|
| C-50 ANOVA: Fatty acid profile of laying hens (PUFA, g/100g yolk)                              | 108  |
| C-51 ANOVA: Fatty acid profile of laying hens (UFA/SFA, g/100g yolk)                           | 108  |
| C-52 ANOVA: Fatty acid profile of laying hens (n-3, g/100g yolk)                               | 108  |
| C-53 ANOVA: Fatty acid profile of laying hens (n-6, g/100g yolk)                               | 109  |
| C-54 ANOVA: Fatty acid profile of laying hens (n-6/n-3, g/100g yolk)                           | 109  |
| C-55 ANOVA: Fatty acid profile of laying hens (C16:0, g/egg)                                   | 109  |
| C-56 ANOVA: Fatty acid profile of laying hens (C16:1, g/egg)                                   | 109  |
| C-57 ANOVA: Fatty acid profile of laying hens (C18:0, g/egg)                                   | 109  |
| C-58 ANOVA: Fatty acid profile of laying hens (C18:1, g/egg)                                   | 110  |
| C-59 ANOVA: Fatty acid profile of laying hens (C18:2, g/egg)                                   | 110  |
| C-60 ANOVA: Fatty acid profile of laying hens (C18:3, g/egg)                                   | 110  |
| C-61 ANOVA: Fatty acid profile of laying hens (C20:5, g/egg)                                   | 110  |
| C-62 ANOVA: Fatty acid profile of laying hens (C22:6, g/egg)                                   | 110  |
| C-63 ANOVA: Fatty acid profile of laying hens (SFA, g/egg)                                     | 111  |
| C-64 ANOVA: Fatty acid profile of laying hens (UFA, g/egg)                                     | 111  |
| C-65 ANOVA: Fatty acid profile of laying hens (MUFA, g/egg)                                    | 111  |
| C-66 ANOVA: Fatty acid profile of laying hens (PUFA, g/egg)                                    | 111  |
| C-67 ANOVA: Fatty acid profile of laying hens (UFA/SFA, g/egg)                                 | 111  |
| C-68 ANOVA: Fatty acid profile of laying hens (n-3, g/egg)                                     | 112  |
| C-69 ANOVA: Fatty acid profile of laying hens (n-6, g/egg)                                     | 112  |
| C-70 ANOVA: Fatty acid profile of laying hens (n-6/n-3, g/egg)                                 | 112  |
| C-71 ANOVA: Copper in heart of laying hens for 168 days.                                       | 112  |
| C-72 ANOVA: Copper in liver of laying hens for 168 days.                                       | 112  |
| C-73 ANOVA: Copper in gizzard of laying hens for 168 days.                                     | 113  |
| C-74 ANOVA: Copper in gastrointestinal tract of laying hens for 168 days.                      | 113  |
| C-75 ANOVA: Copper in excreta of laying hens for 168 days (fresh).                             | 113  |
| C-76 ANOVA: Copper in excreta of laying hens for 168 days (AD).                                | 113  |
| C-77 ANOVA: Copper retention of laying hens for 168 days.                                      | 114  |
| C-78 ANOVA: Copper in excreta of laying hens (after changing to the control diets, fresh wt.). | 114  |

## สารบัญตารางภาคผนวก (ต่อ)

| <b>Table Appendix C.</b>   | <b>หน้า</b> |
|--|-------------|
| C-79 ANOVA: Copper in excreta of laying hens (after changing to the control diets, AD) | 114         |
| C-80 ANOVA: Copper retention of laying hens (after changing to the control diets).     | 114         |
| C-81 ANOVA: Egg production of Japanese quails for 84 days.                             | 114         |
| C-82 ANOVA: Feed intake of Japanese quails for 84 days.                                | 114         |
| C-83 ANOVA: Feed/100 egg of Japanese quails for 84 days.                               | 115         |
| C-84 ANOVA: Feed/kg egg of Japanese quails for 84 days.                                | 115         |
| C-85 ANOVA: Body weight gain of Japanese quails for 84 days.                           | 115         |
| C-86 ANOVA: Egg weight of Japanese quails for 84 days.                                 | 115         |
| C-87 ANOVA: Egg shell thickness of Japanese quails for 84 days.                        | 115         |
| C-88 ANOVA: Yolk color of Japanese quails for 84 days.                                 | 115         |
| C-89 ANOVA: Yolk cholesterol of Japanese quails for 84 days (mg/g yolk).               | 116         |
| C-90 ANOVA: Yolk cholesterol of Japanese quails for 84 days (mg/egg).                  | 116         |
| C-91 ANOVA: Copper in excreta of Japanese quails for 84 days (fresh wt.).              | 116         |
| C-92 ANOVA: Copper in excreta of Japanese quails for 84 days (AD).                     | 116         |
| C-93 ANOVA: Copper retention of Japanese quails for 84 days.                           | 116         |

## อักษรย่อ

|      |   |                      |        |   |                           |
|------|---|----------------------|--------|---|---------------------------|
| ก.   | = | กรัม                 | ppb    | = | part per billion          |
| กก.  | = | กิโลกรัม             | ppm    | = | part per million          |
| ซม.  | = | เซนติเมตร            | FID    | = | flame ionization detector |
| ม.   | = | เมตร                 | GC     | = | gas chromatography        |
| มม.  | = | มิลลิเมตร            | m      | = | meter                     |
| มล.  | = | มิลลิลิตร            | nm     | = | nanometer                 |
| มก.  | = | มิลลิกรัม            | mm     | = | millimeter                |
| ดล.  | = | เดซิลิตร             | μm     | = | micrometer                |
| °ซ   | = | องศาเซลเซียส         | ANOVA  | = | analysis of variance      |
| g    | = | gram                 | CV     | = | coefficient of variation  |
| mg   | = | milligram            | df     | = | degree of freedom         |
| kg   | = | kilogram             | MS     | = | mean square               |
| kcal | = | kilocalorie          | SS     | = | sum of square             |
| ml   | = | milliliter           | SOV    | = | source of variation       |
| dl   | = | deciliter            | S.E.M. | = | standard error of mean    |
| μl   | = | microliter           | Ave.   | = | average                   |
| Exp. | = | experiment           |        |   |                           |
| vs.  | = | versus               |        |   |                           |
| AD   | = | air dry              |        |   |                           |
| DM   | = | dry matter           |        |   |                           |
| CP   | = | crude protein        |        |   |                           |
| EE   | = | ether extract        |        |   |                           |
| ME   | = | metabolizable energy |        |   |                           |
| CF   | = | crude fiber          |        |   |                           |
| Ca   | = | calcium              |        |   |                           |
| aP   | = | available phosphorus |        |   |                           |