

เอกสารอ้างอิง

จรรยา มนีโขติ. 2547. ข้าวชีพชีช กัญแจงที่กำลังคุกคามชาวนา. หนังสือพิมพ์แนวหน้า ปีที่ 25 ฉบับที่ 8579.

สงกรานต์ จิตรากร. 2532. ทรัพยากรเชื้อพันธุ์ข้าวป่าในประเทศไทย. วารสารวิชาการเกษตร เล่มที่ 1: 69-83.

สงกรานต์ จิตรากร, ฉวีวรรณ ภูมิญาโณ, ผกวรรณ ภู่สุวรรณ, กัมปนาท มุขดี. 2538. การบันทึก ลักษณะและวิเคราะห์ลักษณะข้าวป่าในประเทศไทย. วารสารวิชาการเกษตรเล่มที่ 3: 197-218.

Anuntalabhochai, A., R. Chudet, J. Chiangda and P. Apavatjrut. 2000. Genetic diversity within Lychee (*Litchi chinensis* Sonn.) based on RAPD analysis. International Symposium on Tropical and Subtropical Fruit. 26th Nov – 1st Dec 2000. Cairns, Australia.

Barbier P., 1987. Ecological genetic study on wild rice populations from Thailand. M.S. Thesis submitted to Nagoya Univ., Nagoya, Japan, 122 p.

Barbier P.,and A. Ishihama, 1991. Relatedness of annual and perennial strains of *Oryza rufipogon* investigated with nuclear DNA sequences. In Rice Genetics II. Proc.2nd Int. Rice Gen. Symp. pp. 731-734. IRRI. Philippines.

Chang T.T., Y. Pan, Q. Chu, R. Peiris, G. and C. Loresto, 1991. Cytogenetic, electrophoretic, and root studies of javanica rice. In Rice Genetics II. Proc.2nd Int. Rice Gen. Symp. pp. 21-31. IRRI. Philippines.

Chitrakon S., 1995. Characterization, evaluation and utilization of wild rice germplasm in Thailand. 143 p.

Doyle, J.J. and J.L. Doyle. 1987. A rapid DNA isolation Procedure for small quantities of fresh leaf tissue. *Focus* 12: pp. 13-15.

Franskel H., H.D. Brown, J. Burdon, 1991. The Conservation of Plant Biodiversity. Cambridge: Cambridge University Press. 299 p.

Grealy D. R., T.H. Tai, C.H. Sneller, 2002. Identification of red rice, rice, and hybrid populations using microsatellite markers. In Weed Science. Vol.50: pp. 333-339.

- Goldstein D.T. and C.Schlotterer 1998. *Microsatellites Evolution and Applications*. Oxford University press. 352 p.
- IRRI-IBPGR. 1980. Descriptors for rice *Oryza sativa* L. IRRI, P. O. Box 933, Manila, Philippines. 21 p.
- International Rice Research Institute., 1991. Program report for 1989. Philippines.
- Kataya M.T., 1997. Relationships between chromosome number and genomic constitutions in genus *Oryza*. In Science of The Rice Plant. Vol.3 Genetics. (Eds.) T. Matsuo, Y. Futsuhara, F. Kikuchi, and H. Yamaguchi. pp. 39-48. Food and Agriculture Policy Research Center, Tokyo, Japan.
- Kumar, S., K. Tamura, I.B. Jakobsen, and M. Nei. 2001. *MEGA2: Molecular Evolutionary Genetics Analysis* software, Bioinformatics Vol. 17, 12: 1244-1245.
- Lu B. R., 2004. Gene flow from cultivated rice: ecological consequences. ISB News Report. May 2004. pp. 4-6.
- Morishima H.,Y. Shinanoto, Y. Sano and Y.I. Sato. 1984. Observations on wild and cultivated rices in Thailand for ecological genetic study in Southeast Asia. Report of Study-tour in 1983. Report Nat. Inst. Genet. Japan. 82 p.
- Morishima H.,Y. Shinanoto, Y. Sano, Y.I. Sato, S. Chitrakon, P. Barbier, T. Sato, and H. Yamagishi., 1996. Monitoring wild rice populations in permanent study site in Thailand. In Rice Genetics III. Proc. 3rd Int. Rice Gen. Symp. (Ed.) G. S. Khush. 377 p. IRRI. Philippines.
- Morishima H., 1991. Association between Pow-1 variation and seed productivity potential in wild rice. In Rice Genetic 2. Proc.2nd Int. Rice Gen. Symp. pp. 1-10. IRRI. Philippines.
- Nei, M. 1972. Genetic distance between populations. Am Nat 106: 283 – 292.
- Newton, C.R. and A. Graham. 1994. PCR. Introduction to Biotechniques. ZENECA Pharmaceuticals, Mereside, Alderley Park, Macclesfield. Bios scientific publishers in association with the Biochemical Society, 161p.
- Oka H.I. and H. Morishima, 1967. Variations in the breeding systems of a wild rice, *Oryza perennis*. Evolution 21. pp. 249-258.
- Oka H.I., 1988. Origin of Cultivated Rice. Japan Scientific Societies Press. National Institute of Genetics, Japan. 254 p.

- Power L.E. and R. McSorley, 2000. Ecological Principles of Agriculture. Delmar. Thomson Learning. 433 p.
- Rohlf, F. J. 1998. NTSYSpc. Numerical Taxonomy and Multivariate Analysis System. Version 2.0. Department of Ecology and Evolution, state University of New York, Stony Brook, NY.
- Rong J., H. Xia, Y. Zhu, Y. Wang and B. Lu. 2004. Asymmetric gene flow between traditional and hybrid rice varieties (*Oryza sativa*) indicated by nuclear simple sequence repeats and implications for germplasm conservation. In *New Phytologist*. Vol. 163: pp. 439-445.
- Saiki, R.F., D.H. Gelfand and S. Stoffel. 1988. Primer-detected enzymatic amplification of DNA with a thermostable DNA polymerase. *Science* 239: 487 – 491.
- Vaughan D A., 1994. The Wild Relatives of Rice. IRRI-IBPGR, Philippines: A genetic Resources Handbook. IRRI. Philippines. 137 p.
- Willium, J.G.K., A.R. Kubelik, and K.J. Livak. 1990. DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucl. Acids Res* 18: 6531 – 6535.
- Xie Z.W. and Y. Zhou, 1998. Phylogenetic relationship of genus *Oryza* as revealed by RAPD analysis. In IRRN Vol.23. pp. 5-8.
- Yeh, F.C., R.C. Yang, T. Boyle, Z.H. Ye, and J.X. Mao. 1977. POPGENE, the user-friendly shareware for population genetic analysis. Molecular Biology and Biotechnology Center. University of Alberta, Canada.