


TABLE OF CONTENTS

	Page
Acknowledgements	iii
Abstract in Thai	v
Abstract in English	x
Table of contents	xv
List of Tables	xviii
List of Figures	xxx
Chapter 1 Introduction	1
Chapter 2 Literature Reviews	4
1. Area : attributes and usage	4
2. The late production of Kaew mango strategies in the Upper North	11
2.1 Delayed flowering	11
2.2 Extension of panicle growth	12
2.3 Delaying fruit maturation	13
3. Opportunity of late production in fruit tree	15
Chapter 3 Material and Methods	17
1. Promising areas for producing late season Kaew mango	17
2. Farmers' interview	19
3. Experiments in the delayed harvestings	20
3.1 Delaying flowering	20
3.1.1 Pruning	20
3.1.2 Panicle thinning	21
3.2 Extension of panicle growth	21
3.2.1 Paclobutrazol concentrations and time of application	21
3.2.2 Paclobutrazol concentrations and application stage	22
3.3 Delaying fruit maturation	23

3.3.1 Fruit bagging	23
3.3.2 Gibberellin application	23
3.3.2.1 Gibberellin concentrations and number of applications	23
3.3.2.2 Gibberellin concentrations and fruit age	23
3.3.2.3 Gibberellin application and fruit age	24
4. Combination of technology in extending the harvest time of Kaew mango	24
5. Assessment of the farmers response in adopting delayed harvesting techniques	24
Chapter 4 Results and Discussions	26
1. Promising areas for producing late season Kaew mango	26
2. Understanding of the target farmers and area	32
3. Experiments in the delayed harvestings	54
3.1 Delaying flowering	54
3.1.1 Pruning	54
3.1.2 Panicle thinning	70
3.2. Extension of panicle growth	76
3.2.1 PBZ concentrations and time of applications	76
3.2.2 PBZ concentrations and application stage	92
3.3 Delaying the preharvest fruit maturity	115
3.3.1 Fruit bagging	115
3.3.2 GA concentrations and number of applications	125
3.3.3 GA concentrations and fruit age	140
3.3.4 GA application and fruit age	151
4. Combination of technology for delayed harvesting of Kaew mango	172
5. Assessment of the farmers' opinions and views on the practicality of the new technology	194
Chapter 5 Conclusions and Recommendations	216

References	219
Appendices	233
Curriculum vitae	246



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่

Copyright© by Chiang Mai University

All rights reserved

LIST OF TABLES

Table		Page
1	Planting area of fruit trees in the 12 districts of Chiang Mai	31
2	Gender of mango growers in Chiang Dao district	32
3	Age of mango growers in Chiang Dao district	33
4	Education level of mango growers in Chiang Dao district	34
5	Family size of mango growers in Chiang Dao district	34
6	Size of family labor of mango growers in Chiang Dao district	35
7	Farming experience of 64 mango growers in Chiang Dao district	36
8	Source of local knowledge gained by mango growers in Chiang Dao district	37
9	Kind of knowledge gained by mango growers in Chiang Dao district	37
10	Clone of Kaew mango cultivated by growers in Chiang Dao district	38
11	Reason for choice of Kaew mango planting given by growers in Chiang Dao district	39
12	Reason why Kaew mango is no longer attractive by growers in Chiang Dao district	40
13	Farm size of mango growers in Chiang Dao district	41
14	Source of water for mango orchard in Chiang Dao district	41
15	Size of mango orchard in Chiang Dao district	42
16	Tree spacing of mango orchard in Chiang Dao district	43
17	Farmers' interest in late season production in Chiang Dao district	43
18	Objective of mango pruning given by mango growers in Chiang Dao district	44
19	Amount of tree pruning practiced by mango growers in Chiang Dao district	45
20	Pruning time by mango growers in Chiang Dao district	46
21	Farmers' practicing in the delayed harvesting of Kaew mango in Chiang Dao district	47

Table	Page
22 Criteria for a choice of plant bioregulator by mango growers in Chiang Dao district	48
23 Common harvesting time of Kaew mango in Chiang Dao district	49
24 Required harvesting time of Kaew mango in Chiang Dao district	50
25 Number of harvesting time of Kaew mango in Chiang Dao district	51
26 Fruit grading practiced by mango growers in Chiang Dao district	51
27 Fruit grading of Kaew mango in Chiang Dao district	51
28 Type of mango trader in Chiang Dao district	52
29 Price of Kaew mango in June and July in Chiang Dao district	52
30 Source of mango growers' income in Chiang Dao district	53
31 Flush appearance and flush production of Kaew mango trees after pruning in different months	55
32 Quantity and size of Kaew mango flush after pruning in different months	56
33 Flower development of Kaew mango trees after pruning in different months	58
34 Panicle size, floral sex and floral sex ratio of Kaew mango trees after pruning in different months	59
35 Days to each developmental stage of Kaew mango trees after pruning in different months	61
36 Fruiting panicle of Kaew mango after pruning in different months	62
37 Number of fruit per panicle of Kaew mango after pruning in different months	63
38 Yield of Kaew mango at harvesting after pruning in different months	64
39 Fruit size, fruit weight and flesh percentage of Kaew mango after pruning in different months	65
40 Size and weight of Kaew mango seed after pruning in different months	66
41 Peel color of Kaew mango after pruning in different months	68
42 Fruit stalk toughness, fruit firmness and flesh color of Kaew mango after pruning in different months	69
43 Fruit quality of Kaew mango after pruning in different months	70

Table	Page
44 Return bloom appearance of axillary panicles after thinning at different panicle lengths	71
45 Panicle appearance of axillary panicle after thinning at different panicle lengths	72
46 Axillary panicle length after thinning at different panicle lengths	73
47 Axillary panicle diameter after thinning at different panicle lengths	73
48 Floral sex percentage and ratio of axillary panicle after thinning at different panicle lengths	74
49 Days from panicle appearance to full bloom stage of control and axillary panicle after thinning at different panicle lengths	75
50 Panicle development of Kaew mango after spraying with different PBZ concentrations at panicle length of 1 and 5 cm	77
51 Male and perfect flowers of Kaew mango after spraying with different PBZ concentrations at panicle length of 1 and 5 cm	78
52 Days to each developmental stage of Kaew mango after spraying with different PBZ concentrations at panicle length of 1 and 5 cm	79
53 Number of fruit per panicle of Kaew mango after spraying with different PBZ concentrations at panicle length of 1 and 5 cm	81
54 Yield of Kaew mango fruit after spraying with different PBZ concentrations at panicle length of 1 and 5 cm	83
55 Fruit weight, fruit size and flesh content of Kaew mango at harvest after spraying with different PBZ concentrations at panicle length of 1 and 5 cm	84
56 Weight and size of Kaew mango seeds at harvest after spraying with different PBZ concentrations at panicle length of 1 and 5 cm	86
57 Internal qualities of Kaew mango fruits at harvest after spraying with different PBZ concentrations at panicle length of 1 and 5 cm	88
58 Peel color of Kaew mango fruit at harvest after spraying with different PBZ concentrations at panicle length of 1 and 5 cm	90

Table	Page	
59	Flesh color of Kaew mango fruit at harvest after spraying with different PBZ concentrations at panicle length of 1 and 5 cm	91
60	Moisture percentage, chlorophyll content, total nonstructural carbohydrate (TNC) and reducing sugar (RS) of Kaew mango leaves treated with three PBZ concentrations at panicle appearance stage	92
61	Flowering percentage and panicle size of Kaew mango treated with three PBZ concentrations at panicle appearance stage	93
62	Span of time from panicle appearance to full bloom, floral sex percentage and floral sex ratio of Kaew mango treated with three PBZ concentrations at panicle appearance stage	94
63	Moisture, chlorophyll, total nonstructural carbohydrate (TNC) and reducing sugar (RS) contents of Kaew mango leaves treated with three PBZ concentrations at panicle appearance stage	95
64	Span of time from full bloom to peanut stage, fruited panicle and number of fruit per panicle of Kaew mango at peanut stage treated with three PBZ concentrations at panicle appearance stage	96
65	Weight and size of Kaew mango fruit at peanut stage treated with three PBZ concentrations at panicle appearance stage	97
66	Moisture and chlorophyll contents of Kaew mango leaves and fruits at peanut stage treated with three PBZ concentrations at panicle appearance stage	98
67	Total nonstructural carbohydrate (TNC) and reducing sugar (RS) contents in leaves and fruits of Kaew mango at peanut stage after treated with three PBZ concentrations at panicle appearance stage	99
68	Span of time from peanut to bird's egg stage, fruited panicle and number of fruit per panicle of Kaew mango at bird's egg stage after treated with three PBZ concentrations at panicle appearance	99
69	Weight and size of Kaew mango fruit at bird's egg stage after treated with three PBZ concentrations at panicle appearance stage	100

Table	Page	
70	Moisture and chlorophyll contents of Kaew mango leaves and fruits at bird's egg stage after treated with three PBZ concentrations at panicle appearance stage	101
71	Total nonstructural carbohydrate (TNC) and reducing sugar (TNC) contents of Kaew mango at bird's egg stage after treated with three PBZ concentrations at panicle appearance stage	102
72	Span of time from bird's egg stage to hen's egg stage, fruited panicle and number of fruit per panicle of Kaew mango at hen's egg stage after treated with three PBZ concentrations at panicle appearance stage	103
73	Weight and size of Kaew mango fruit at hen's egg stage after treated with three PBZ concentrations at panicle appearance stage	104
74	Seed weight and flesh content of Kaew mango at hen's egg stage after treated with three PBZ concentrations at panicle appearance stage	104
75	Moisture and chlorophyll contents of Kaew mango leaves and fruits at hen's egg stage after treated with three PBZ concentrations at panicle appearance stage	105
76	Total nonstructural carbohydrate (TNC) and reducing sugar (RS) contents of Kaew mango leaf and fruit at hen's egg stage after treated with three PBZ concentrations at panicle appearance stage	106
77	Span of time from full bloom to harvest stage, fruited panicle and number of fruit per panicle of Kaew mango after treated with three PBZ concentrations at panicle appearance stage	107
78	Yield of Kaew mango fruits treated with three PBZ concentrations at panicle appearance stage	108
79	Size and weight of fully maturity Kaew mango fruits after treated with three PBZ concentrations at panicle appearance stage	108
80	Size and weight of Kaew mango seeds and flesh content treated with three PBZ concentrations at panicle appearance stage	109

Table	Page
81 Peel color of Kaew mango fruits treated with three PBZ concentrations at panicle appearance stage measured	110
82 Fruit firmness and fruit stalk toughness of Kaew mango treated with three PBZ concentrations at panicle appearance stage	111
83 Flesh color of Kaew mango fruits treated with three PBZ concentrations at panicle appearance stage	112
84 Total soluble solids (TSS) and titratable acidity (TA) of Kaew mango fruits treated with three PBZ concentrations at panicle appearance stage	112
85 Moisture and chlorophyll content of Kaew mango fruits treated with three PBZ concentrations at panicle appearance stage	113
86 Total nonstructural carbohydrate (TNC) and reducing sugar (RS) contents of Kaew mango fruits and leaves treated with three PBZ concentrations at panicle appearance stage	114
87 Fruit width of Kaew mango after bagging at 30, 45 and 60 DAF	115
88 Fruit length of Kaew mango after bagging at 30, 45 and 60 DAF	116
89 Fruit thickness of Kaew mango after bagging at 30, 45 and 60 DAF	117
90 Fruit drop of Kaew mango after bagging at 30, 45 and 60 DAF	117
91 Fruit retention of Kaew mango after bagging at 30, 45 and 60 DAF	118
92 Fruit weight and flesh content of fully mature Kaew mango at harvesting after bagging at 30, 45 and 60 DAF	119
93 Size and weight of Kaew mango seeds at harvesting after bagging at 30, 45 and 60 DAF	120
94 Internal fruit qualities of fully mature Kaew mango at harvesting after bagging at 30, 45 and 60 DAF	121
95 Peel color of fully mature Kaew mango fruits at harvesting after bagging at 30, 45 and 60 DAF	123
96 Flesh color of fully mature Kaew mango fruits at harvesting after bagging at 30, 45 and 60 DAF	124

Table	Page
97 Days to harvesting of Kaew mango after bagging at 30, 45 and 60 DAF	125
98 Fruit width of Kaew mango after GA applications at 80 DAF	126
99 Fruit length of Kaew mango after GA applications at 80 DAF	127
100 Fruit thickness of Kaew mango after GA applications at 80 DAF	128
101 Days to harvesting, size and weight of Kaew mango fruits treated with different GA concentrations and number of applications at 80 DAF	130
102 Size and weight of Kaew mango seeds and flesh percentage of fruits treated with different GA concentrations and number of applications at 80 DAF	131
103 Fruit drop percentage of Kaew mango treated with different GA concentrations and number of applications at 80 DAF	133
104 Number of retained fruit per panicle of Kaew mango treated with different GA concentrations and number of applications at 80 DAF	134
105 Fruit stalk toughness, fruit firmness, contents of total soluble solids (TSS) and titratable acidity (TA) of Kaew mango treated with different GA concentrations and number of applications at 80 DAF	136
106 Peel color of Kaew mango fruits treated with different GA concentrations and number of applications at 80 DAF	138
107 Flesh color of Kaew mango fruits treated with different GA concentrations and number of applications at 80 DAF	139
108 Fruit width of Kaew mango treated with different GA concentrations and fruit ages	140
109 Fruit length of Kaew mango treated with different GA concentrations and fruit ages	141
110 Fruit thickness of Kaew mango treated with different GA concentrations and fruit ages	142
111 Fruit weight, flesh content and fruit size at harvesting of Kaew mango fruits treated with different GA concentrations and fruit ages	143

Table	Page
112 Weight and size at harvesting of Kaew mango seeds treated with different GA concentrations and fruit ages	145
113 Yield and days to harvesting of Kaew mango treated with different GA concentrations and fruit ages	146
114 Internal fruit qualities of Kaew mango treated with different GA concentrations and fruit ages	148
115 Peel color of Kaew mango fruits treated with different GA concentrations and fruit ages	150
116 Flesh color of Kaew mango fruits treated with different GA concentrations and fruit ages	151
117 Fruit retention percentage of Kaew mango treated with GA 50 ppm and different fruit ages	152
118 Total nonstructural carbohydrate (TNC) contents in Kaew mango leaves and fruits treated with GA 50 ppm and different fruit ages	154
119 Reducing sugar (RS) contents in Kaew mango leaves and fruits treated with GA 50 ppm and different fruit ages	156
120 Chlorophyll contents in Kaew mango leaves and fruits treated with GA 50 ppm and different fruit ages	158
121 Fruit weight of Kaew mango treated with GA 50 ppm and different fruit ages	159
122 Fruit width of Kaew mango treated with GA 50 ppm and different fruit ages	160
123 Fruit length of Kaew mango treated with GA 50 ppm and different fruit ages	160
124 Fruit thickness of Kaew mango treated with GA 50 ppm and different fruit ages	161
125 Seed weight of Kaew mango treated with GA 50 ppm and different fruit ages	161
126 Seed width of Kaew mango treated with GA 50 ppm and different fruit ages	162
127 Seed length of Kaew mango treated with GA 50 ppm and different fruit ages	162
128 Seed thickness of Kaew mango treated with GA 50 ppm and different fruit ages	163

Table	Page	
129	Flesh content of Kaew mango fruits treated with GA 50 ppm and different fruit ages	163
130	Leaf moisture content of Kaew mango treated with GA 50 ppm and different fruit ages	164
131	Fruit moisture content of Kaew mango treated with GA 50 ppm and different fruit ages	165
132	Peel color of Kaew mango fruits at shoulder side treated with GA 50 ppm and different fruit ages	166
133	Peel color of Kaew mango fruits at middle side treated with GA 50 ppm and different fruit ages	167
134	Peel color of Kaew mango fruits at apex side treated with GA 50 ppm and different fruit ages	168
135	Days to harvesting and yield of Kaew mango treated with GA 50 ppm and different fruit ages	169
136	Fruit stalk toughness, fruit firmness, total soluble solids (TSS), titratable acidity (TA) and flesh color of Kaew mango treated with GA 50 ppm	171
137	Panicle diameter of Kaew mango after spraying PBZ 1000 ppm on panicle appearance	173
138	Panicle length of Kaew mango after spraying PBZ 1000 ppm on panicle appearance	173
139	Blooming percentage of Kaew mango after spraying PBZ 1000 ppm on panicle appearance	174
140	Floral percentage and floral sex ratio of Kaew mango after spraying PBZ 1000 ppm on panicle appearance	175
141	Days to each developmental stage of Kaew mango after application of two bioregulators	177
142	Fruited panicle percentage of Kaew mango after application of two bioregulators	179

Table	Page
143 Number of fruit per panicle of Kaew mango after application of two bioregulators	180
144 Fruit size at harvesting after application of two bioregulators	181
145 Seed size and flesh content of Kaew mango at harvesting after application of two bioregulators	182
146 Moisture content of Kaew mango fruits after application of two bioregulators	182
147 Chlorophyll content of Kaew mango after application of two bioregulators	183
148 Total nonstructural carbohydrate (TNC) contents of Kaew mango after application of two bioregulators	184
149 Reducing sugar (RS) contents of Kaew mango after application of two bioregulators	185
150 Peel color at shoulder side of Kaew mango fruits at interval seven days from 85 DAF until harvesting	188
151 Peel color at middle side of Kaew mango fruits at interval seven days from 85 DAF until harvesting	189
152 Peel color at apex side of Kaew mango fruits at interval seven days from 85 DAF until harvesting	190
153 Flesh color of Kaew mango fruits at interval seven days from 85 DAF until harvesting	191
154 Fruit firmness, fruit stalk toughness, total soluble solids (TSS) and titratable acidity (TA) of Kaew mango at harvesting after application of two bioregulators	192
155 Yield of Kaew mango after application of two bioregulators	193
156 Gender of farmers who interested in producing late season Kaew mango	195
157 Age of the farmers who interested in producing late season Kaew mango	195
158 Education level of farmers who interested in producing late season Kaew mango	196

Table	Page
159 Orchard size of the farmers who interested in producing late season Kaew mango	197
160 Age of Kaew mango tree mentioned by farmers who interested in producing late season Kaew mango	198
161 Experience years of the farmers who interested in producing late season Kaew mango	199
162 Occupation category of the farmers who interested in producing late season Kaew mango	199
163 Assets of the farmers who interested in producing late season Kaew mango	200
164 Farm gate price of Kaew mango between year 2002-2004 at Mae Ore Nai village, Chiang Mai	201
165 Reasons for low price of Kaew mango given by the farmers who interested in producing late season Kaew mango	202
166 Opinions of the mango growers on the improving of Kaew mango price	203
167 Farmers' recognition of the area producing late season Kaew mango	203
168 Farmers' recognition of better earning by delayed harvesting of Kaew mango	204
169 Farmers' experience of practicing delayed harvesting of Kaew mango	204
170 Farmers' acceptance of using plant bioregulator	205
171 Possibility of producing late season Kaew mango	206
172 Farmers' requirement for producing late season Kaew mango	206
173 Farmers' opinions about the practical method to produce late season Kaew mango	207
174 Farmers' expectation of the delayed harvesting technology	207
175 Limitation of the delayed harvesting technique	208
176 Usefulness of the delayed harvesting technology	209
177 Appropriateness of the delayed harvesting technology	209
178 Farmers' opinion about input cost of producing a kilogram of in season Kaew mango	210

Table	Page
179 Farmers' acceptance of the additional cost for producing a kilogram of late season Kaew mango	211
180 Target harvesting period of mango growers	212
181 Required minimum price/kg for late season Kaew mango	212
182 Farmers' confidence in the delayed harvesting experiment	213
183 Practicability of the delayed harvesting technology	214
184 Transfer of the delayed harvesting technology to the neighbor	214

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
 Copyright© by Chiang Mai University
 All rights reserved

LIST OF FIGURES

Figure	Page
1 Position of Mae Ore Nai village, Chiang Dao district, Chiang Mai Province	6
2 The average temperature and rainfall at Chiang Dao district in 2002	7
3 The average temperature and rainfall at Chiang Dao district in 2003	8
4 The average temperature and rainfall at Chiang Dao district in 2004	9
5 Conceptual framework for producing the late season of Kaew mango in the rainfed uplands of Upper North	18
6 Diagram to identify promising area for producing late season of Kaew mango	19
7 The potential areas at district level for producing late season Kaew mango	27
8 The geological characteristics of the potential areas (dark shading)	29
9 Formation of axillary panicle after thinning	70