

Table of Contents

	Page
Acknowledgements	iii
Abstract (English)	v
Abstract (Thai)	x
Table of Contents	xiii
List of Tables	xvi
List of Illustrations	xxi
Abbreviations and Symbols	xxii
Chapter 1 Introduction	1
1.1 Background	1
1.2 Objectives	3
Chapter 2 Literature Review	4
2.1 Hybrid Rice	5
2.2 Heterosis	5
2.3 Genetic Basis of Heterosis	6
2.4 Floral Biology of Hybrid	6
2.5 Sequence and Duration of Blooming	7
2.6 Stamen Characteristics	8
2.7 Stigma Characteristics	10
2.8 Extent of Heterosis	11
2.9 Cytoplasmic Genetic Male Sterility	12
2.10 Genetics of Fertility Restoration	13
2.11 Male Sterility System	14

2.11.1	Cytoplasmic- genetic Male Sterility	15
2.11.2	Enviromental-sensitive genetic male sterility	16
	Photoperiod-sensitive male sterility	17
	Thermosensitive-sensitive male sterility	17
2.11.3	Chemically-induced male sterility	19
2.12	Hybrid rice production	21
2.12.1	Three-line hybrid	21
	CMS line (A- line)	21
	Maintainer (B- line)	22
	Restorer line (R-line)	23
2.12.2	Two-line hybrid	25
2.13.3	One-line method	26
2.13	Breeding CMS Lines with a High Outcrossing Rate	27
Chapter 3 Appropriate Restorer Lines for Hybrid Rice Production		28
3.1	Introduction	28
3.2	Materials and Methods	29
3.2.1	Pollen Fertility and Agronomic Character Study	30
3.2.2	Test of restorer lines (R-line)	31
3.3	Results	33
3.3.1	Pollen Fertility and Agronomic Character Study	33
3.3.2	Relationship between Pollen Fertility and Spikelet	36
	Fertility of R-lines	
3.3.3	Fertility Restoration of R-line	36
3.3.4	Evaluation of Variety/line Performance	39

3.4 Discussion	46
Chapter 4 Combining ability of inbred lines and heterosis of F1 hybrids	48
4.1 Introduction	48
4.2 Materials and Methods	49
4.3 Results	54
4.3.1 Estimation of Combining ability	54
4.3.2 Estimation of Heterosis	78
4.4 Discussion	94
Chapter 5 Yield Trial and Agronomic Character Evaluation for F1 Hybrid Lines.	97
5.1 Introduction	97
5.2 Materials and Methods	98
5.2.1 Yield Trial of Hybrid Cross	98
5.2.2 Blast Disease and Brown planthopper Resistance Test	101
5.3 Results	104
5.3.1 Yield Trial of Hybrid Cross	104
5.3.2 Blast Disease Resistance Evaluation	107
5.3.3 Brown Planthopper Resistance Estimation	108
5.3.4 Physical and Chemical Evaluation of Grains	109
5.4 Discussion	121
Chapter 6 General Discussion	124
References	129
Curriculum vitae	141

List of Tables

Table	Description of Table	Page
2.1	Differences between the CMS line and its corresponding maintainer line	22
3.1	Agronomic characters of A-line and B-line rice studied at Pathumthani Rice Research Center in rainy season 2002	35
3.2	Pollen fertility and spikelet fertility of F1 hybrids of rice varieties studied at Pathumthani Rice Research Center, dry season 2002	41
3.3	Percentage of filled grain of F1 hybrids derived from crossing between A- line and R- line studied at Pathumthani Rice Research Center, dry season 2002	42
3.4	Grain yield (gm/ plant) of F1 hybrids studied at Pathumthani Rice Research Center, dry season 2002	43
3.5	Number of panicles per hill of F1 hybrids studied at Pathumthani Rice Research Center, dry season 2002	44
3.6	Number of grains per panicle of F1 hybrids studied at Pathumthani Rice Research Center, dry season 2002	45
4.1	Analysis of variance of grain yield and yield components of F1 hybrid grown at Pathumthani Rice Research Center in dry season 2003	57
4.2	Estimate of GCA effects of 4 CMS-testers and 16 R-lines were for grain yield and yield components evaluated at Pathumthani Rice Research Center in dry season 2003	58
4.3	Estimate of SCA effect of crosses combination for grain yield (kg/rai) among F1 hybrid of rice grown at Pathumthani Rice Research Center in dry season 2003	60

Table	Description of Table	Page
4.4	Estimate of SCA effect of crosses combination for plant height among F1 hybrid of rice grown at Pathumthani Rice Research Center in dry season 2003	61
4.5	Estimate of Estimate of SCA effect of crosses combination for number of panicle/ hill among F1 hybrid of rice grown at Pathumthani Rice Research Center in dry season 2003	62
4.6	Estimate of SCA effect of crosses combination for number of grain / panicle among F1 hybrid of rice grown at Pathumthani Rice Research Center in dry season 2003	63
4.7	Estimate of SCA effect of crosses combination for %filled grain/panicle among F1 hybrid of rice grown at Pathumthani Rice Research Center in dry season 2003	64
4.8	Estimate of SCA effect of crosses combination for 1,000 - grain weight among F1 hybrid of rice grown at Pathumthani Rice Research Center in dry season 2003	65
4.9	Analysis of variance of grain yield and yield components of F1 hybrid grown at Pathumthani Rice Research Center in rainy season 2003	69
4.10	Estimate of GCA effects of 4 CMS-testers and 16 R-lines were evaluate for grain yield and yield component evaluate at Pathumthani Rice Research Center in rainy season 2003	70

Table	Description of Table	Page
4.11	Estimate of SCA effect of cross combination for grain yield (kg/rai) among F1 hybrids of rice grown at Pathumthani Rice Research Center in rainy season 2003	72
4.12	Estimate of SCA effect of cross combination for plant height among F1 hybrids of rice grown at Pathumthani Rice Research Center in rainy season 2003	73
4.13	Estimate of of SCA effect of cross combination for number of panicle/ hill among F1 hybrids of rice grown at Pathumthani Rice Research Center in rainy season 2003	74
4.14	Estimate of SCA effect of cross combination for number of grain / panicle among F1 hybrids of rice grown at Pathumthani Rice Research Center in rainy season 2003	75
4.15	Estimate of SCA effect of cross combination for 1,000 - grain weight among F1 hybrids of rice grown at Pathumthani Rice Research Center in rainy season 2003	76
4.16	Estimate of SCA effect of cross combination for %filled grain/panicle among F1 hybrids of rice grown at Pathumthani Rice Research Center in rainy season 2003	77

4.17	Heterosis (H) and heterobeltiosis (Hb) for grain yield of F1 hybrids grown at Pathumthani Rice Research Center in dry season 2003	80
------	---	----

Table	Description of Table	Page
4.18	Heterosis (H) and heterobeltiosis (Hb) for plant height of F1 hybrids grown at Pathumthani Rice Research Center in dry season 2003	81
4.19	Heterosis (H) and heterobeltiosis (Hb) for panicle per hill of F1 hybrids grown at Pathumthani Rice Research Center in dry season 2003	82
4.20	Heterosis (H) and heterobeltiosis (Hb) for number of grain per panicle of F1 hybrids grown at Pathumthani Rice Research Center in dry season 2003	83
4.21	Heterosis (H) and heterobeltiosis (Hb) for percent filled grain of F1 hybrids grown at Pathumthani Rice Research Center in dry season 2003	84
4.22	Heterosis (H) and heterobeltiosis (Hb) for 1000-grain weight of F1 hybrids grown at Pathumthani Rice Research Center in dry season 2003	85
4.23	Heterosis (H) and heterobeltiosis (Hb) for grain yield of F1 hybrids grown at Pathumthani Rice Research Centre in rainy season 2003	88

4.24	Heterosis (H) and heterobeltiosis (Hb) for plant height of F1 hybrids grown at Pathumthani Rice Research Centre in rainy season 2003	89
------	--	----

Table	Description of Table	Page
4.25	Heterosis (H) and heterobeltiosis (Hb) for panicle per hill of F1 hybrids grown at Pathumthani Rice Research Centre in rainy season 2003	90
4.26	Heterosis (H) and heterobeltiosis (Hb) for no. of grain per panicle of F1 hybrids grown at Pathumthani Rice Research Centre in rainy season 2003	91
4.27	Heterosis (H) and heterobeltiosis (Hb) for percent filled grain of F1 hybrids grown at Pathumthani Rice Research Centre in rainy season 2003	92
4.28	Heterosis (H) and heterobeltiosis (Hb) for 1000-grain weight of F1 hybrids grown at Pathumthani Rice Research Centre in rainy season 2003	93
5.1	Grain yield among F1 hybrids crosses of yield trial conducted at Pathumthani Rice Research Center, dry season 2004	112
5.2	Yield components among F1 hybrids crosses of yield trial conducted at Pathumthani Rice Research Center, dry season 2004	113

5.3	Grain yield among F1 hybrids crosses of yield trial conducted at Pathumthani Rice Research Center, rainy season 2004	114
5.4	Yield components among F1 hybrids crosses of yield trial conducted at Pathumthani Rice Research Center, rainy season 2004	116
5.5	Reaction of F1 hybrids for blast and brown planthopper resistance	118
5.6	Physical and chemical characteristic of grain of F1 hybrids	119

List of Illustrations

Figure	Description of Figure	Page
2.1	The spikelet of rice	10
2.2	Presence of certain dominant restores gene (s) in the nucleus makes a line capable of restoring fertility in the hybrid derived from it and a CMS line	24
2.3	Schematic description of the use of EGMS line for developing two-line hybrids	25
2.4	Megasporogenesis and megagametogenesis in <i>Arabidopsis</i>	26
3.1	Pollen of F1 hybrid stained with I-IK solution	37
3.2	Relationship between pollen fertility and spikelet fertility in F1 hybrid crosses	38

Abbreviations

CMS	cytoplasmic genetic male sterile
GCA	general combining ability
SCA	specific combining ability
EGMS	environmental genic male sterility
TGMS	thermo sensitive genic male sterile
PGMS	photoperiod sensitive genic male sterile
A-line	male sterile parent involving cytoplasm
M-line	maintainer line
R-line	restorer line
SPR1	Supanburi 1
CNT1	Chainat 1
KHSPR	Khao Hawm Supanburi
KLK1	Khao Hawm Klongluang 1
PTT1	Khaow Hawm Pathumthani 1
R	resistance
S	susceptible
MR	moderately resistance
MS	moderately susceptible
VR	very resistance
VS	very susceptible
BC	backcross
CK	check
GA ₃	gibberellic acid