

TABLE OF CONTENTS

Acknowledgements	iii
Abstract (English)	v
Abstract (Thai)	x
List of Tables	xvii
List of Figures	xxii
Introduction	1
Chapter 1 Literature review	4
1.1 Cultivated rice	4
1.1.1 Asian cultivated rice (<i>O. sativa</i> L.)	4
1.1.2 African rice (<i>O. glaberrima</i> Stedu.)	5
1.2 Wild rice	5
1.2.1 Common wild rice (<i>O. rufipogon</i> Griff.)	5
1.2.2 <i>O. barthii</i> A. Chev. (formerly known as <i>O. brevilugata</i>)	6
1.3 Cultivated rice and wild rice in Thailand	7
1.4. Weedy rice	8
1.5. Gene flow	9
1.5.1 Crop-to-wild gene flow	9
1.5.2 Crop-to-weed gene flow	10
1.5.3 Crop-to-crop gene flow	10
1.6 Gene flow assessment	11

1.7 Interspecific hybridization	12
1.7.1 Reproductive barriers	13
1.7.2 F ₁ fitness	13
Chapter 2 Characterization of cultivated and common wild rice populations	15
2.1 Introduction	15
2.2 Materials and methods	17
2.3 Results	21
2.4 Discussion	39
Chapter 3 Interspecific hybridization between cultivated rice (<i>O. sativa</i>) and common wild rice (<i>O. rufipogon</i>)	42
3.1 Introduction	42
3.2 Materials and methods	44
3.2.1 Experiment 1: Crossability between cultivated rice x wild rice	44
3.2.2 Experiment 2: Evaluation of F ₁ hybrids of cultivated rice x wild rice	44
3.2.3 Experiment 3: Evaluation of F ₂ generations between cultivated rice and wild rice.	45
3.3 Results	46
3.3.1 Crossability between cultivated rice x wild rice	46
3.3.2 Evaluation of F ₁ and F ₂ generations of cultivated rice x wild rice	48

3.4 Discussion	84
Chapter 4 Evidence of gene flow between cultivated rice (<i>O. sativa</i> L.) and wild rice (<i>O. rufipogon</i> Griff.)	87
4.1 Introduction	87
4.2 Materials and methods	88
4.3. Results	91
4.4. Discussion	108
Chapter 5 General Discussion	111
References	115
Appendix	124
Curriculum vitae	141

LIST OF TABLES

Table	Page
2.1 Description of cultivated rice varieties and common wild rice populations used in the experiment	19
2.2 Primer sequences, repeat motif, expected PCR product size, annealing temperature and chromosomal locations of seven microsatellite primers	20
2.3 Morphological characteristics and Shannon-Weaver Index (H') of cultivated rice varieties and common wild rice populations	22
2.4 Physiological characteristics of eight cultivated rice varieties and four common wild rice populations	28
2.5 Allele size ranges in base pairs of seven microsatellite loci in eight cultivated varieties and three wild rice populations	31
2.6 Number of specific and shared alleles (% in parentheses) in seven microsatellite loci based on eight cultivated rice varieties and three wild rice populations	31
2.7 Gene diversity (H_e) of eight cultivated and three wild rice populations based on seven microsatellite loci	38
2.8 Partition of genetic diversity of cultivated and wild rice within and between eight cultivated rice and three wild rice populations based on seven microsatellite loci	38
3.1 Number of spikelets fertilized, number and percent seed set (%) of	47

	crosses between eight cultivated rice (as female) and three common wild rice (as male) parents	
3.2	Number of spikelets fertilized, number of seed set and percent seed set (%) of reciprocal crosses between wild rice x cultivated rice and back crosses between F ₁ hybrids to cultivated and wild rice parents	47
3.3	Seed germination (%) and seedling survival (%) of F ₁ hybrids and their parents	49
3.4	Seed germination (%) and seedling survival (%) of F ₂ generation and their parents	50
3.5	Pigment presentation on plant parts of F ₁ and segregation of F ₂ between six cultivated rice (CR) and wild rice from Kanchanburi (KC) compared with their parents	52
3.6	Pigment presentation on plant part of F ₁ and segregation of F ₂ between six cultivated rice (CR) and wild rice from Nakorn Nayok (NY) compared with their parents	53
3.7	Morphological characteristics of F ₁ hybrids and F ₂ populations between cultivated rice and wild rice (KC and NY) compared with their parents	55
3.8	Days to flowering of F ₁ hybrids between cultivated rice (CR) and wild rice (KC and NY) compared with their parents	57
3.9	Mean, range and standard deviation of days to flowering of F ₂ populations between cultivated rice (CR) x wild rice (KC and NY)	57
3.10	Culm length (cm) of F ₁ hybrids between cultivated rice (CR) and wild	59

	rice (KC and NY) compared with their parents	
3.11	Mean, range and standard deviation of culm length (cm) of F ₂ populations between cultivated rice (CR) x wild rice (KC and NY)	59
3.12	Panicle length (cm) of F ₁ hybrids between eight cultivated rice (CR) varieties and two wild rice (KC and NY) compared with their parents	62
3.13	Mean, range and standard deviation of panicle length (cm) of F ₂ populations between cultivated rice (CR) x wild rice (KC and NY)	62
3.14	Number of panicles plant ⁻¹ of F ₁ hybrids between cultivated rice (CR) and wild rice (KC and NY) compared with their parents	64
3.15	Mean, range and standard deviation of number of panicle ⁻¹ of F ₂ populations between cultivated rice (CR) x wild rice (KC and NY)	64
3.16	Number of spikelets panicle ⁻¹ of F ₁ hybrids between cultivated rice (CR) and wild rice (KC and NY) compared with their parents	68
3.17	Mean, range and standard deviation of number of spikelets panicle ⁻¹ of F ₂ populations between cultivated rice (CR) x wild rice (KC and NY)	68
3.18	Number of seeds panicle ⁻¹ of F ₁ hybrids between cultivated rice (CR) and wild rice (KC and NY) compared with their parents	70
3.19	Mean, range and standard deviation of number of seed panicle ⁻¹ of F ₂ populations between cultivated rice (CR) x wild rice (KC and NY)	70
3.20	Seed fertility (%) of F ₁ hybrids between eight cultivated rice (CR) and two wild rice (KC and NY) compared with their parents	73
3.21	Mean, range and standard deviation of seed fertility (%) of F ₂	73

	populations between cultivated rice (CR) x wild rice (KC and NY)	
3.22	Seed shattering (%) of F ₁ hybrids between cultivated rice (CR) and wild rice (KC and NY) compared with their parents	75
3.23	Mean, range and standard deviation of seed shattering (%) of F ₂ populations between cultivated rice (CR) x wild rice (KC and NY)	75
3.24	Seed width (mm) of F ₁ hybrids between cultivated rice (CR) and wild rice (KC and NY) compared with their parents	79
3.25	Mean, range and standard deviation of seed length (mm) of F ₂ populations between cultivated rice (CR) x wild rice (KC and NY)	79
3.26	Seed length (mm) of F ₁ hybrids between cultivated rice (CR) and wild rice (KC and NY) compared with their parents	80
3.27	Mean, range and standard deviation of seed length (mm) of F ₂ populations between cultivated rice (CR) x wild rice (KC and NY)	80
4.1	Frequency of six morphological characteristics in cultivated, weedy and wild rice	93
4.2	Allele frequencies of crop rice, weedy rice and wild rice evaluated with three microsatellite markers, RM1, RM164 and RM211	99
4.3	Frequency (%) of alleles specific to cultivated rice (SPR1 and CNT1) and wild rice (KC1 and KC2) in crop rice, weedy rice and wild rice based on three microsatellite markers	100
4.4	Genotype frequencies of crop rice, weedy rice and wild rice evaluated with three microsatellite loci	101
4.5	Genotypic frequency (%) of cultivated rice (SPR1 and CNT1), wild	103

rice (KC1 and KC2) and hybrid types in crop rice, weedy rice and wild rice based on three microsatellite markers

- 4.6 Gene diversity (H_e), Fixation index (F_{IS}) and outcrossing rate (t) cultivated, weedy and wild rice based on three microsatellite loci (number in parentheses indicate standard deviation) 107
- 4.7 Genetic differentiation among populations (F_{ST}) matrix values determined across three microsatellite loci in crop rice, weedy rice and wild rice 107

LIST OF FIGURES

Figure		Page
2.1	Morphological characteristics of cultivated rice.	23
2.2	Pigment presentation on different plant parts of purple rice (Kum Doi Sa Ket).	23
2.3	Morphological characteristics of wild rice.	24
2.4	Microsatellite amplification products from eight cultivated and three wild rice genomic DNA with primer RM1.	32
2.5	Microsatellite amplification products from eight cultivated and three wild rice genomic DNA with primer RM164.	32
2.6	Microsatellite amplification products from eight cultivated and three wild rice genomic DNA with primer RM167.	33
2.7	Microsatellite amplification products from eight cultivated and three wild rice genomic DNA with primer RM211.	33
2.8	Microsatellite amplification products from eight cultivated and three wild rice genomic DNA with primer RM588.	34
2.9	Microsatellite amplification products from eight cultivated and three wild rice genomic DNA with primer RM225.	34
2.10	Microsatellite amplification products from eight cultivated and three wild rice genomic DNA with primer RM341.	35
3.1	Distribution of days to flowering of F ₂ populations between	58

	cultivated rice x wild rice.	
3.2	Distribution of culm length (cm) of F ₂ populations between cultivated rice x wild rice.	60
3.3	Distribution of panicle length (cm) of F ₂ populations between cultivated rice x wild rice.	63
3.4	Distribution of number of panicles plant ⁻¹ of F ₂ populations between cultivated rice x wild rice.	65
3.5	Distribution of number of spikelets panicle ⁻¹ of F ₂ populations between cultivated rice x wild rice.	69
3.6	Distribution of number of seeds panicle ⁻¹ of F ₂ populations between cultivated rice x wild rice.	71
3.7	Distribution of seed fertility (%) of F ₂ populations between cultivated rice x wild rice.	74
3.8	Distribution of seed shattering (%) of F ₂ populations between cultivated rice x wild rice.	76
3.9	Seed shape of F ₁ hybrids between cultivated rice x KC wild rice compared with their parents.	81
3.10	Seed shape of F ₁ hybrids between cultivated rice x NY wild rice compared with their parents.	82
3.11	Seed shape of F ₂ populations between cultivated rice x KC wild rice (left) and cultivated rice x wild rice.	83
4.1	Location of rice samples were collected from Kanchanaburi,	90

	Thailand.	
4.2	Weedy rice invaded into rice field. Rice plants were taller than cultivated rice plus panicle with awn were recognized as weedy rice.	90
4.3	Morphological characteristics of weedy rice.	94
4.4	Distribution of days to flowering of purified cultivated rice from RRI, farmers' field, weedy rice and wild rice.	95
4.5	Distribution of culm length (cm) of purified cultivated rice from RRI, farmers' field, weedy rice and wild rice.	96
4.6	Distribution of seed shattering (%) of purified cultivated rice from RRI, farmers' field, weedy rice and wild rice.	97
4.7	Microsatellite amplification products generated from cultivated rice (SPR1) farmers' field genomic DNA with primer RM1 (a), RM164 (b) and RM211 (c).	104