## VIII

## TABLE OF CONTENTS

		Page
ACK	NOWLEDGMENT	Ш
ABS'	TRACT IN THAI	IV
ABSTRACT IN ENGLISH		O O VI
TAB	LE OF CONTENTS	VIII
LIST	OF TABLES	X
LIST OF FIGURES		XIV
CHA	PTER 1 INTRODUCTION	503
1.1	Background information	1
1.2	Objectives	4
1.3	Education/application advantages	4
СНА	PTER 2 LITERATURE REVIEW	5
2.1	Soybean	5
2.2	Soybean yogurt	10
2.3	Starter micro-organisms used in fermented dairy products	12
2.4	Probiotic	13
2.5	Factors that affect the growth of probiotics	20
2.6	Prebiotic	21
2.7	Utilization of stabilizers in fermented dairy products	26
	phyright <sup>©</sup> by Chiang Mai	
СНА	PTER 3 MATERIALS AND METHODS	33
3.1	Materials Control Cont	$\bigcirc$ $\bigcirc$ 33 $\bigvee$ $\bigcirc$
3.2	Chemical Reagents	33
3.3	Equipments	34
3 /	Mathods	25

CHA	PTER 4 RESULTS AND DISSCUSSION	51
4.1	Growth curves of probiotic bacteria in soymilk yogurt with	51
	yogurt starter bacteria	
4.2	Basic formula of soybean yogurt	56
4.3	The effect of soymilk yogurt ingredients on the viability of	59
	probiotic bacteria in soymilk yogurt during storage at 4 <sup>0</sup> C	
	for 21 days.	
4.4	The effect of nutrient compounds on the viability of probiotic	72
	bacteria in soymilk yogurt during storage at 4 °C for 21 day	
CHA	PTER 5 CONCLUSION AND RECOMMENDATION	96
5.1	Conclusion	96
5.2	Recommendation	98
REFE	ERENCES	99
APPE	ENDIX A	105
APPE	ENDIX B	112
CUR	RICULUM VITAE	136

ลิขสิทธิมหาวิทยาลัยเชียงใหม Copyright<sup>©</sup> by Chiang Mai University All rights reserved

Table		Page
2.1	The nutitional value of 100 g soybean	6
2.2	Comparative chemical analysis (g 100g <sup>-1</sup> ) of soymilk and cow's milk	7
2.3	Typical composition of crude and refined soybean oil	9
2.4	Physiological and biochemical characteristics	15
	of Bifidobacterium spp. and L. acidophilus	
2.5	Commercial products containing Bifidobacterium spp.	17
	and Lactobacillus acidophilus	
2.6	The functional properties of fibersol-2 in its applications	23
2.7	The amounts of volume of hydrocolloids used by the U.S.A industries	27
4.1	Chemical properties of soymilk yogurt fermented with S. thermophilus,	52
	L. bulgaricus and B. bifidum or L. acidophilus at 43°C for 42 h	
4.2	Microbiological properties of soymilk yogurt fermented with	53
	S. thermophilus and L. bulgaricus and B. bifidum	
	or L. acidophilus at 43°C for 42 h	
4.3	Chemical and microbiological properties of soymilk yogurt during	56
	storage at 4 <sup>0</sup> C for 21 days	
4.4	Physical and chemical properties of soymilk yogurt during storage	57
	at 4 <sup>o</sup> C for 21 days.	
4.5	The effect of experimental factors on the properties of soymilk yogurt	61
	added with B. bifidum based on a Plackett and Burman design	
4.6	Major and minor experimental factors that affected the properties of	63
	soymilk yogurt added with B. Bifidum based on	
	a Plackett and Burman design	
4.7	The effect of experimental factors on the properties of soymilk yogurt	64
	added with L. acidophilus based on a Plackett and Burman design	

Table		Page
4.8	Major and minor experimental factors that affected the properties of soymilk yogurt added with <i>L. acidophilus</i> based on a Plackett and Burman design	67
4.9	Suitable levels of major soymilk yogurt ingredients for soymilk yogurt added with B. bifidum using a 2 <sup>3</sup> Factorial experiment in center point	69
4.10	Optimal levels of soymilk yogurt ingredients for soymilk yogurt added with B. bifidum	70
4.11	Suitable levels of major soymilk yogurt ingredients for soymilk yogurt added with L. acidophilus using a 2 <sup>2</sup> Factorial experiment in center point	71
4.12	Optimal levels of soymilk yogurt ingredients for soymilk yogurt added with L. acidophilus	71
4.13	Chemical and microbiological properties of soymilk yogurt added with B. bifidum and in the presence of 5 % (w/w) fibersol-2 during storage at 4°C for 21 days	73
4.14	Physical and chemical properties of soymilk yogurt added with B. bifidum and in the presence of 5 % (w/v) fibersol-2 during storage at 4°C for 21 days	74
4.15	Chemical and microbiological properties of soymilk yogurt added with L. acidophilus and in the presence of 5% (w/v) Fructo-oligosaccharide during storage at 4°C for 21 days	76
4.16	Phisical and chemical properties of soymilk yogurt added with L. acidophilus and in the presence of 5% (w/v) Fructo-oligosaccharide during storage at 4°C for 21 days	77

Table		Page
4.17	Chemical and Microbiological properties of soymilk yogurt products	80
	from B. bifidum added Fructo-oligosaccharide during a storage at 4 <sup>0</sup> C for 21 day	
4.18	Physical and chemical properties of soymilk yogurt added with B. bifidu	m 81
	and in the presence of 5% (w/v) Fructo-oligosaccharide during storage at 4°C for 21 days	
4.19	Chemical and microbiological properties of soymilk yogurt added with	83
4.17	L. acidophilus and in the presence of 5% (w/v)	503
	fructo-oligosaccharide during storage at 4°C for 21 days	
4.20	Physical and chemical properties of soymilk yogurt added with	84
	L. acidophilus and in the presence of 5% (w/v) fructo-oligosaccharide	
	during storage at 4°C for 21 days	
4.21	Chemical and Microbiological properties of soymilk yogurt	87
	added with B. bifidum and in the presence of 5% (w/w)	
	tomato extract during storage at 4 <sup>0</sup> C for 21 days	
4.22	Physical and chemical properties of soymilk yogurt added with	89
	B. bifidum and in the presence of 5% (w/v) tomato extract	
	during storage at 4°C for 21 days	
4.23	Reductions in the viable numbers of S. thermophilus, L. bulgaricus	90
	and B. bifidum in soymilk yogurt added with different nutrient-	
	compounds during storage at 4°C for 21 days	
4.24	Chemical and microbiological properties of soymilk yogurt	92
	added with L. acidophilus and in the presence of 5% (w/v)	
	tomato extract during storage at 4°C for 21 days	

	Page
Physical and chemical properties of soymilk yogurt added with  L. acidophilus and in the presence of 5% (w/v) tomato extract	94
Reductions in the viable numbers of S. thermophilus, L. bulgaricus and L. acidophilus in soymilk yogurt added with different nutrient-compounds during storage at 4°C for 21 days	95
	Physical and chemical properties of soymilk yogurt added with L. acidophilus and in the presence of 5% (w/v) tomato extract during storage at 4°C for 21 days  Reductions in the viable numbers of S. thermophilus, L. bulgaricus and L. acidophilus in soymilk yogurt added with different nutrient-

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright<sup>©</sup> by Chiang Mai University All rights reserved

## XIV

# LIST OF FIGURES

Figure	างมยนติ	Page
4.1	The growth curve of S. thermophilus, L. bulgaricus and L. acidophilus	54
	(log CFU/ml) in soymilk yogurt during fermentation at 43°C for 42 h.	
4.2	The growth curve of S. thermophilus, L. bulgaricus and B. bifidum	55
	(log CFU/ml) in soymilk yogurt during fermentation at 43°C for 42 h.	
4.3	The number of S. thermophilus, L. bulgaricus	74
	and B. bifidum (log CFU/ml) in soymilk yogurt added with	
	fibersol-2 during storage at 4 <sup>0</sup> C for 21 days	
4.4	The number of S. thermophilus, L. bulgaricus and	77
	L. acidophilus (log CFU/ml) in soymilk yogurt added with	
	fibersol-2 during storage at 4°C for 21 days	
4.5	The number of S. thermophilus, L. bulgaricus and	81
	B. bifidum (log CFU/ml) soymilk yogurt added with	
	fructo-oligosaccharide during storage at 4°C for 21 days	
4.6	The number of S. thermophilus, L. bulgaricus and	85
	L. acidophilus (log CFU/ml) in soymilk yogurt added with	
•	fructo-oligosaccharide during storage at 4°C for 21 days	
4.7	The number of S. thermophilus, L. bulgaricus and	. 88
	B. bifidum (log CFU/ml) in soymilk yogurt added with	The
	tomato extract during storage at 4°C for 21 days	
4.8	The number of S. thermophilus, L. bulgaricus and L. acidophilus	91
	(log CFU/ml) in soymilk yogurt added with tomato extract during	
	storage at 4°C for 21 days	