

เอกสารอ้างอิง

- “การส่งออกข้าว.” (2547). [ระบบออนไลน์]. แหล่งที่มา [http://www.thairice.org/demo/html/aboutrice/about-rice 8-2.html](http://www.thairice.org/demo/html/aboutrice/about-rice%208-2.html). [7 พฤศจิกายน 2549].
- “จุดกำเนิด และประวัติข้าวไทย.” (2547). [ระบบออนไลน์]. แหล่งที่มา [http://www.Thairice.org/html/aboutrice/about-rice 1.html](http://www.Thairice.org/html/aboutrice/about-rice%201.html). [7 พฤศจิกายน 2549].
- ดวงจันทร์ เสงส์สวัสดิ์. (2538). ข้าวและน้ำมันรำข้าวช่วยลดความเสี่ยงต่อโรคหัวใจ โดยลดการสังเคราะห์คอเลสเตอรอลในร่างกาย. วารสารอาหาร, 2,140.
- นัยนา บุญทวีวัฒน์ และเรวดี จงส์วัฒน์. (2545). น้ำมันรำข้าวทางเลือกเพื่อสุขภาพของคนไทย. กรุงเทพฯ: สำนักพิมพ์โอเดียนสโตร์.
- นิธิยา รัตนাপนนท์. (2545). เคมีอาหาร. กรุงเทพฯ: สำนักพิมพ์โอเดียนสโตร์.
- นิธิยา รัตนাপนนท์. (2548). วิทยาศาสตร์การอาหารของไขมันและน้ำมัน. กรุงเทพฯ: สำนักพิมพ์โอเดียนสโตร์.
- เนตรนภิส วัฒนสุขชาติ. (2538). ประโยชน์ของข้าวต่อระบบอวัยวะต่างๆในร่างกาย. กรุงเทพฯ: สำนักพิมพ์โอเดียนสโตร์.
- วราพร พงษ์ธรรมาพานิช “ การวิเคราะห์เอกลักษณ์และปริมาณโทโคเฟอรอลและโอรีซานอลในกระบวนการผลิตน้ำมันรำข้าว” วิทยานิพนธ์ปริญญาวิทยาศาสตรมหาบัณฑิต มหาวิทยาลัยเทคโนโลยีพระจอมเกล้าธนบุรี 2543.
- ศักดิ์ บวร. (2536). วิตามินอี. กรุงเทพฯ: สำนักพิมพ์สมิต.
- อรอนงค์ นัยวิกุล. (2547). ข้าว : วิทยาศาสตร์และเทคโนโลยี. กรุงเทพฯ: สำนักพิมพ์มหาวิทยาลัยเกษตรศาสตร์.
- Abidi, S.L. (2003). Tocopherol-derived minor constituents in selected plant seed oils. *J. Am. Oil Chem. Soc.*, 80, 327-333.
- Ahmed, M.K., Daun, J.K., and Przybylski, R. (2005). FT-IR methodology for quantitation of total tocopherol, tocotrienols and plastochromanol-8 in vegetable oils. *J. Food Compos. Anal.*, 18, 359-364.
- Alibaba.com Corporation. [online]. Available : http://www.alibaba.com/catalog/11027475/Rice_Bran_Distilled_Fatty_Acid.html [2006, May 22].

- Almeida, M. E. M., Rusig, O., and Guzman, E. C. (2001). Tocopherols extraction from soybean oil deodorizer distillate by saponification: the effect of acetone/water solution composition. IFT Annual Meeting-New Orleans, Louisiana.
- Amaral, J. S., Alves, M. R., Seabra, R. M. and Oliveira, B. P. P. (2005). Vitamin E composition of walnuts (*Juglans regia* L.): A 3-year comparative study of different cultivars. *J. Agric. Food Chem*, 53, 5467-5472.
- AOCS. (1997). *Official methods and recommended Practices of the American Oil Chemists' Society* (5th ed.). Washington, DC: American Oil Chemists' Society Press. (Method Ce 8-89).
- Aoun, E., Rima, J., Chidiac, G., and Hanna, K. (2005). High-performance liquid chromatographic and spectrofluorometric determination of α -tocopherol in a natural plant: *Ferula hermonis* (Zaloooh root). *J. Food Compos. Anal*, 18, 607-615.
- Ausman, L.M., Rong, N., and Nicolosi, R.J. (2005). Hypocholesterolemic effect of physically refined rice bran oil: studies of cholesterol metabolism and early atherosclerosis in hypercholesterolemic hamsters. *J. Nutr. Biochem*, 16, 521-529.
- Azlina, M. F. N., Nafeeza, M. I. and Khalid, B. A. K. (2005). Effect of tocotrienol on lipid peroxidase in experimental gastritis induced by restraint stress. *J. Nutr*, 4, 69-72.
- Bera, D., Lahiri, D., and Nag, A. (2006). Studies on a natural antioxidant for stabilization of edible oil and comparison with synthetic antioxidants. *Food Engineering*, 74, 542-545.
- Bereau, D., Benjelloun-Mlayah, B., Banoub, J., and Bravo, R. (2003). FA and unsaponifiable composition of five amazonizn palm kernel oils. *J. Am. Oil Chem. Soc*, 80, 49-53.
- Bernardini, E. (1985). *Oilseeds, oils and fats, Raw Materials*. New York: Publishing House.
- Bernardini, E. (1985). *Oilseeds, oils and fats, Oils and fat processing*. New York: Publishing House.

- Bonvehi, J.S., Coll, F.V., and Rius, I.A. (2000). Liquid chromatographic determination of tocopherols and tocotrienols in vegetable oils, formulated prepalations, and biscuits. *J. AOAC International*, 83, 627-634.
- Borher, J.R.Z., Gonçalves, L.A.G., and Felício, P.E. (2002). α - and γ -Tocopherol levels in Nelore steer blood plasma after a single oral treatment of soybean oil deodorizer distillate (SODD). *Meat Sci*, 61, 301-306.
- Bourgeois, C. (1992) . Determination of vitamin E : tocopherols and tocotrienols. London: Elsevier science publishers, Ltd.
- Carlucci, G., Mazzeo, P., Governatore, S.D., Giacomo, G.D., and Re G.D. (2001). Liquid chromatographic method for the analysis of tocopherols in malt sprouts with supercritical fluid extraction. *J. Chromatogr. A*, 935, 87-91.
- Carpenter, A.P. (1979). Determination of tocopherol in vegetable oils. *J. Am. Oil Chem. Soc*, 56, 668-671.
- Chang, L.-C., Chang, H.-T., and Sun, S.W. (2006). Cyclodextrin-modified microemulsion electrokinetic chromatography for separation of α -, γ -, δ -tocopherol and α -tocopherol acetate. *J. Chromatogr. A*, 1110, 227-234.
- Chase, G.W., Akoh, C.C., and Eitenmiller, R.R. (1994). Analysis of tocopherol in vegetable oils by high-performance liquid chromatography: comparison of fluorescence and evaporative light-scattering detection. *J. Am. Oil Chem. Soc*, 71, 877-880.
- Chávez-Servín, J.L., Castellote, A.I., and López-Sabater, M.C. (2006). Simultaneous analysis of vitamin A and E in infant milk-based formulae by normal-phase high-performance liquid chromatography-diode array detection using a short narrow-bore column. *J. Chromatogr. A*, 1122, 138-143.
- Che, Man, Y.B., Ammawath, W., and Mirghani, M.E.S. (2005). Determining α -tocopherol in refined bleached and deodorized palm olein by Fourier transform infrared spectroscopy. *Food Chem*, 90, 323-327.
- Chen, M.H., and Bergman, C.J. (2005). A rapid procedure for analyzing rice bran tocopherol, tocotrienol and γ -oryzanol contents. *J. Food Compos. Anal*, 18, 139-151.

- Cherukuri, V., Cheruvanky, R., Lynch, I., and MacPeak, D.L. (1999). Process for obtaining micronutrient enriched rice bran oil. Patent No.: 5,989,344.
- Choo, Y.M., Ma, A.N., Yahaya, H., Yamauchi, Y., Bounoshita, M., and Saito, M. (1996). Separation of crude palm oil components by semipreparative supercritical fluid chromatography. *J. Am. Oil Chem. Soc.*, 73, 523-525.
- Chu, B.S., Baharin, B.S., Man-Che, B.S., and Quek, S.Y. (2004). Separation of vitamin E from palm fatty acid distillate using silica : I Equilibrium of batch adsorption. *J. Food Engineering*, 62, 97-103.
- Chu, B.S., Baharin, B.S., Man-Che, B.S., and Quek, S.Y. (2004). Separation of vitamin E from palm fatty acid distillate using silica : III Batch desorption study. *J. Food Engineering*, 64, 1-7.
- Chu, B.S., Baharin, B.S., and Quek, S.Y. (2002). Factors affecting pre-concentration of tocopherols and tocotrienols from palm fatty acid distillate by lipase-catalysed hydrolysis. *Food Chem*, 79, 55-59.
- Chu, B.S., Quek, S.Y., and Baharin, B.S. (2003). Optimization of enzymatic hydrolysis for concentration of vitamin E in palm fatty acid distillate. *Food Chem*, 80, 295-302.
- Chun, J., Lee, J., Ye, L., Exler, J., and Eitenmiller, R.R. (2006). Tocopherol and tocotrienol contents of raw and processed fruits and vegetables in the United States diet. *J. Food Compos. Anal*, 19, 196-204.
- Chun, J., Ye, L., Lee, J., and Eitenmiller, R. R. (2003). Effect of peanut butter manufacture on vitamin E. *J. Food Sci*, 68, 2211-2214.
- Cunha, S.C., Amaral, J.S., Fernandes, J.O., and Oliveira, M.B.P.P. (2006). Quantification of tocopherols and tocotrienols in portuguese olive oils using HPLC with three different detection systems. *J. Agric. Food Chem*, 54, 3351-3356.
- Delgado-Zamarreno, M.M., Bustamante-Rangel, M., Sanchez-Perez, A., and Carabias-Martinez R. (2004). Pressurized liquid extraction prior to liquid chromatography with electrochemical detection for the analysis of vitamin E isomers in seeds and nuts. *J. Chromatogr. A*, 1056, 249-252.

- Dionisi, F., Prodoliet, J., and Tagliaferri, E. (1995). Assessment of olive oil adulteration by reversed-phase high-performance liquid chromatography/ amperometric detection of tocopherols and tocotrienols. *J. Am. Oil Chem. Soc*, 72, 1505-1511.
- Eitenmiller, R. and Lee, J. (2004). Vitamin E: Food chemistry, composition, and analysis. New York: Marcel Dekker, Inc.
- Evans, J.C., Kodali, D.R., and Addis, P.B. (2002). Optimal tocopherol concentrations to inhibit soybean oil oxidation. *J. Am. Oil Chem. Soc*, 79, 47-51.
- Feeter, D.K. (1974). Determination of tocopherols, sterols, and steryl esters in vegetable oil distillates and residues. *J. Am. Oil Chem. Soc*, 51, 184-187.
- Fernandez, R.X.E., Shier, N.W., and Watkins, B.A. (2000). Effect of alkalisaponification, enzymatic hydrolysis and storage time on the total carotenoid concentration of costa Rican crude palm oil. *J. Food Compos. Anal*, 13, 179-187.
- Gast, K., Jungfer, M., Saure, C., and Brunner, G. (2005). Purification of tocopherols from edible oil. *J. Supercrit. Fluid*, 34, 17-25.
- German, J.B. (2002). Antioxidant. In: Branen, A.L., Davidson, P.M., Salminen, S., and Thorngate, J.H. III (Ed.), *Food additives*. New York: Marcel Dekker, Inc.
- Gianello, R., Libinaki, R., Azzi, A., Gavin, P.D., Negis, Y., Zingg, J.-M., Holt, P., Keah, H.-H., Griffey, A., Smallridge, A., West, S.M., and Ogru, E. (2005). α -Tocopheryl phosphate: a novel, natural form of vitamin E. *Free Radical Biol. & Med*, 39, 970-976.
- Gómez-Coronado, D.J.M., Ibañez, E., Rupérez, F.J., and Barbas, C.(2004). Tocopherol measurement in edible products of vegetable origin. *J. Chromatogr A*, 1054, 227-233.
- Graciano-Verdugo, A.Z., Peralta, E., González-Ríos, H., Soto-Valdez, H. (2006). Determination of α -tocopherol in low-density polyethylene (LDPE) films by diffuse reflectance Fourier transform infrared (DRIFT-IR) spectroscopy compared to high performance liquid chromatography (HPLC). *Analytica Chimica. Acta*, 557, 367-372.

- Gülçın, I., Oktay, M., Kireççi, E., and Küfrevioğlu, O.I. (2003). Screening of antioxidant and antimicrobial activities of anise (*Pimpinella anisum* L.) seed extracts. *Food Chem*, 83, 371-382.
- Hadolin, M., Skerget, M., Knez, Z., and Bauman, D. (2001). High pressure extraction of vitamin E-rich oil from *Silybum marianum*. *Food Chem*, 74, 355-364.
- Hargrove, K.L., and Jr. (1994) . Processing and utilization of rice bran in the United States. In: Marshall, W. E. and Wadsworth, J. I. Rice science and technology. New York: Marcel Dekker, Inc.
- Hu, W., Wells, J.H., Shin, T.-S., and Godber, J.S. (1995) . Recovery of vitamin E and oryzanols from isopropanol-crude rice bran oil miscella using direct-saponification and liquid-liquid extraction. IFT Annual Meeting 1995. Dep. of Biol. & Agric. Eng., Louisiana State Univ. Agric. Cent., Baton Rouge, LA 70803-4505, USA, 201 .
- Hu, W., Wells, J.H., Shin, T.-S., and Godber, J.S. (1996). Comparison of isopropanol and hexane for extraction of vitamin E and oryzanols from stabilized rice bran. *J. Am. Oil Chem. Soc*, 73, 1653-1656.
- Huo, Nelis, H.J., Lavens, P., Sorgeloos, P., and Leenheer, A.P. de. (1996). Determination of vitamin E in aquatic organisms by high-performance liquid chromatography with fluorescence detection. *Analytical-Biochem*, 242, 123-128.
- Ibáñez, E., Benavides, A.M.H., Señoráns, F.J., and Reglero, G. (2002). Concentration of sterols and tocopherols from olive oil with supercritical carbon dioxide. *J. Am. Oil Chem. Soc*, 79, 1255-689.
- Ibáñez, E., Palacios, J., Señoráns, F.J., Santa-María, G., Tabera, J., and Reglero, G. (2000). Isolation and separation of tocopherols from olive by-products with supercritical fluids. *J. Am. Oil Chem. Soc*, 77, 187-190.
- Indira, T.N., Hemavathy, J., Khatoon, S., Krishna, A.G.G., and Bhattacharya, S. (2000). Water degumming of rice bran oil : a response surface approach. *J. Food Engineering*, 43, 83-90.

- Iqbal, S., Bhangar, M.I., and Anwar, F. (2005). Antioxidant properties and components of some commercially available varieties of rice bran in Pakistan. *Food Chem*, 93, 265-272.
- Iqbal, S., Bhangar, M.I., and Anwar, F. (2007). Antioxidant properties and components of bran extracts from selected wheat varieties commercially available in Pakistan. *LWT*, 40, 361-367.
- Jacobs, L. (2005). Process for the production of tocotrienols. Patent No.: US 6,838,104 B2.
- Joint FAO/WHO Expert Committee on Food Additives (JECFA). (2005). General standard for food additives. *Codex Stan*, 192-1995.
- Julio, S., Susana, N., and Alfonso, V. (2000). Thermal stability of some commercial synthetic antioxidants. *J. Food Sci*, 77, 933-936.
- Kamal-Eldin, A., Gorgen, S., Pettersson, J., and Lampi, A.-M. (2000). Normal-phase high-performance liquid chromatography of tocopherols and tocotrienols: comparison of different chromatographic columns. *J. Chromatogr. A*, 881, 217-227.
- Kanaya, Y., Doi, T., Sasaki, H., Fujita, A., Matsuno, S., Okamoto, K., Nakano, Y., Tsujiwaki, S., Furuta, H., Nishi, M., Tsuno, T., Taniguchi, H., and Nanjo, K. (2004). Rice bran extract prevents the elevation of plasma peroxy lipid in KKAY diabetic mice. *Diabetes Research & Clinical Practice*, 66, 157-160.
- Khor, H.T., Chieng, D.Y., and Ong, K.K. (1995). Tocotrienols inhibit liver HMG CoA reductase activity in the guinea pig. *Nutrition Research*, 15, 537-544.
- Kim, J.S. (2005). Radical scavenging capacity and antioxidant activity of the E vitamin fraction in rice bran. *J. Food Sci*, 70, 208-213.
- Kim, J.S., Godber, J.S., King, J.M., and Prinyawiwatkul, W. (2001). Inhibition of cholesterol autoxidation by the nonsaponifiable fraction in rice bran in an aqueous model system. *J. Am. Oil Chem. Soc*, 78, 685-689.
- Kramer, J.K.G., Blais, L., Fouchard, R.C., Melnyk, R.A., and Kallury, K.M.R. (1997). A rapid method for the determination of vitamin E forms in tissues and diet by high-performance liquid chromatography using a normal-phase diol

- column. *Cent. for Food & Animal Res., Agric. & Agri-Food*, 32, 323-330.
- Lavedrine, F., Ravel, A., Poupard, A., and Alary, J. (1997). Effect of geographic origin, variety and storage on tocopherol concentrations in walnuts by HPLC. *Food-chem*, 58, 135-140.
- Matthaus, B., Vosmann, K., Pham, L.Q., and Aitzetmueller, K. (2003). FA and tocopherol composition of Vietnamese oilseeds. *J. Am. Oil Chemists Soc.*, 80, 1013-1020.
- Meijboom, P.W., and Jongenotter, G.A. (1979). A quantitative determination of tocotrienols and tocopherols in palm oil by TLC-GLC. *J. Am. Oil Chemists Soc.*, 56, 33-35.
- Mendes, M.F., Pessoa, F.L.P., and Uller, A.M.C. (2005). Optimization of the process of concentration of vitamin E from DDSO using supercritical CO₂. *Brazilian J. Chem Engineering*, 22, 83-91.
- Mendes, M.F., Pessoa, F.L.P., and Uller, A.M.C. (2002). An economic evaluation based on an experimental study of the vitamin E concentration present in deodorizer distillate of soybean oil using supercritical CO₂. *J. Supercrit. Fluid*, 23, 257-265.
- Minhajuddin, M., Beg, Z.H., and Iqbal, J. (2005). Hypolipidemic and antioxidant properties of tocotrienol rich fraction isolated from rice bran oil in experimentally induced hyperlipidemic rats. *Food and chemical Toxicology*, 43, 747-753.
- Mohamed, H.M.A., and Awatif, I.I. (1998). The use of sesame oil unsaponifiable matter as a natural antioxidant. *Food Chem*, 62, 269-276.
- Nagai, T., Inoue, R., Inoue, H., and Suzuki, N. (2003). Preparation and antioxidant properties of water extract of propolis. *Food Chem*, 80, 29-33.
- Nicolosi, R.J., Rogers, E.J., Ausman, L.M., and Orthoefer, F.T. (1994). Rice bran oil and its health benefits. In: Marshall, W. E. and Wadsworth, J. I. *Rice science and technology*. New York: Marcel Dekker, Inc.
- Nyström, L., Tanja, A., Lampi, A.-M., Moreau, R.A., and Piironen, V. (2007). A comparison of the antioxidant properties of steryl ferulates with tocopherol at high temperatures. *Food Chem*, 101, 947-954.

- Park, S. R., Kim, Y. H., Park, H. J., and Lee, Y. S. (2004). Stability of tocopherols and tocotrienols extracted from unsaponifiable fraction of rice bran under various temperature and oxygen condition [online]. Available : http://www.cropscience.org.au/icsc2004/poster/5/1/1/654_park.html. [2005, June 14].
- Pazos, M., Sanchez, L., and Medina, I. (2005). α -Tocopherol oxidation in fish muscle during chilling and frozen storage. *J. Agri and Food Chem*, 53, 4000-4005.
- Pokorny, J., Yanishlieva, N., and Gordon, M. (2001). Antioxidants in food practical applications. New York: CRC press.
- Ponte Vedra Soap Shoppe, Inc. [online]. Available : <http://www.pvsoap.com/carrierbasespecialtyoils>. [2005, Jan 22].
- Puspitasari-Nienaber, N.L., Ferruzzi, M.G., and Schwartz, S.J. (2002). Simultaneous detection of tocopherols, carotenoids, and chlorophylls in vegetable oils by direct injection C30 RP-HPLC with coulometric electrochemical array detection. *J. Am. Oil Chem. Soc*, 80, 49-53.
- Qureshi, A.A., Bradlow, B.A., Salser, W.A., and Brace, L.D. (1997). Novel tocotrienols of rice bran modulate cardiovascular disease risk parameters of hypercholesterolemic humans. *J. Nutr. Biochem*, 8, 290-298.
- Rangkadilok, N., Sitthimonchai, S., Worasuttayangkurn, L., Mahidol, C., Ruchirawat, M., and Satayavivad, J. (2007). Evaluation of free radical scavenging and antityrosinase activities of standardized longan fruit extract. *Food & Chem. Toxicology*, 45, 328-336.
- Rattanathanalerk, M., Chiewchan, N., and Srichumpoung. (2005). Effect of thermal processing on the quality loss of pineapple juice. *J. Food Engineering*, 66, 259-265.
- Rogers, E.J., Rice, S.M., Nicolosi, R.J., Carpenter, D.R., McClelland, C.A., and Romanczyk, L.J. (1993). Identification and quantitation of γ -oryzanol components and simultaneous assessment of tocopherols in rice bran oil. *J. Am. Oil Chemists Soc*, 70, 301-307.
- Rossell, B.BSC., ARCH., and Dphil. (1999). Oil and Fats : Vegetable oils and fats.

New York: Leatherhead Publishing.

- Rossi, M., Alamprese, C., and Ratti, S. (2007). Tocopherols and tocotrienols as free radical-scavengers in refined vegetable oils and their stability during deep-fat frying. *Food Chem*, 102, 812-817.
- Ryynanen, M., Lampi, A.M., Salo-Vaananen, P., Ollilainen, V., and Piironen, V. (2004). A small-scale sample preparation method with HPLC analysis for determination of tocopherols and tocotrienols in cereals. *J. Food Compos. Anal*, 17, 749-765.
- Santos, J.C.O., Santos, I.M.G.D., Souza, A.G.D., Prasad, S., and Santos, A.V.D. (2002). Thermal stability and kinetic study on thermal decomposition of commercial edible oils by thermogravimetry. *J. Food Sci*, 67, 1393-1398.
- Sarmiento, C.M.P., Ferreira, S.R.S., and Hense, H. (2006). Supercritical fluid extraction (SFE) of rice bran oil to obtain fractions enriched with tocopherols and tocotrienols. *J. Chem. Engineering*, 23, 243-249.
- Scalia, S., Renda, A., Ruberto, G., Bonina, F., and Menegatti, E. (1995). Assay of vitamin A palmitate and vitamin E acetate in cosmetic creams and lotions by supercritical fluid extraction and HPLC. *J. Pharm. & Biomed. Anal*, 13, 273-277.
- Senevirathne, M., Kim, S.-H., Siriwardhana, N., Ha, J.-H., Lee, K.-W., and Jeon, Y.-J. (2006). Antioxidant potential of *Ecklonia cava* on reactive oxygen species scavenging, metal chelating, reducing power and lipid peroxidation inhibition. *Food Sci Tech Int*, 12, 27-38.
- Seppanen, C.M., Rahmani, M., and Saari Csallany, A. (2003). Simultaneous determination of chlorophylls, pheophytins, β -carotene, tocopherols, and tocotrienols in olive and soybean oils by high performance liquid chromatography. *J. Food Sci*, 68, 1644-1647.
- Setiadi, D.H., Chass, G.A., Torday, I.I., Varro, A., and Papp, J.G. (2003). Vitamin E models : Shortened sidechain models of α , β , γ and δ tocopherol and tocotrienol-a density functional study. *J. Molecular Structure*, 637, 11-26.
- Sheppard, A.J., Pennington, J.A.T., and Weihrauch, J.L. (1993). Analysis and distribution of vitamin E in vegetable oils and foods. In: Packer, L. and Fuchs,

J. Vitamin E in Health and Disease. New York: Marcel Dekker, Inc.

- Sivakumar, G., Bacchetta, L., Gatti, R., and Zappa, G. (2005). HPLC screening of natural vitamin E from Mediterranean plant biofactories- a basic tool for pilot-scale bioreactors production of α -tocopherol. *J. Plant Physiol*, 162, 1280-1283.
- Slover, H.T., Thompson JR, R.H., and Merola, G.V. (1983). Determination of tocopherols and sterols by capillary gas chromatography. *J. Am. Oil Chem. Soc*, 60, 1524-1528.
- Szulczewska-Remi, A., Nogala-Kalucka, M., Kwiatkowski, J., Lampart-Szczapa, E., and Rudzinska, M. (2004). Preparation of tocochromanol concentrate from red palm oil, Industrial Chemistry Research Institute, Warsaw, Poland.
- Tasioula-Margari, M. and Okogeri, O. (2001). Simultaneous determination of phenolic compounds and tocopherols in virgin olive oil using HPLC and UV detection. *Food Chem*, 74, 377-383.
- Tavčar-Kalcher, G. and Vengušt, A. (2007). Stability of vitamins in premixes. *Animal Feed Sci & Tech*, 132, 148-154.
- Taylor, J.B., Richar, T.M., Wilhelm, C.L., Chrysam, M.M., Otterburn, M., and Leveille, G.A. (1997). Rice-bran oil antioxidant. Nabiso Inc.
- Therhault, A., Chao, J.T., and Gapor, A. (2002). Tocotrienol is the most effective vitamin E for reducing endothelial expression of adhesion molecules and adhesion to monocytes. *Atherosclerosis*, 160, 21-30.
- Therhault, A., Chao, J.T., Wang, Q., Gapor, A., and Adeli, K. (1999). Tocotrienol : A review of its therapeutic potential. *Clinical Biochemistry*, 23, 309-319.
- USAID: United States Agency International Development from the American people. (2007). Fortification basics : stability.
- U.S. Department of Health and Human Services, Food and Drug Administration, Center for Drug Evaluation and Research (CDER) and Center for Biologics Evaluation and Research (CBER). (2003). Guidance for Industry Q1A(R2) Stability Testing of New Drug Substances and Products. ICH.

- Verhé, R., Verleyen, T., Hoed, V.V., and Greyt, W.D. (2006). Influence of refining of vegetable oils on minor components. *J. Oil Palm Research*, 168-179.
- Verleyen, T., Verhe, R., Garcia, L., Dewettinck, K., and Huyghebaert, Greyt, W.D. (2001). Gas chromatographic characterization of vegetable oil deodorization distillate. *J. Chromatogr A*, 921, 277-285.
- Weber, C., Podda, M., Rallis, M., Thiele, J.J., Traber, M.G. and Packer, L. (1997). Efficacy of topically applied tocopherols and tocotrienols in protection of murine from oxidative damage induced by uv-irradiation. *Free Radical Biology and Medicine*, 22, 761-769.
- Wilson, T.A., Nicolosi, R.J., Woolfrey, B., and Kritchevsky, D. (2007). Rice bran oil and oryzanol reduce plasma lipid and lipoprotein cholesterol concentrations and aortic cholesterol ester accumulation to a greater extent than ferulic acid in hypercholesterolemic hamsters. *J. Nutr. Biochem*, 18, 105-112.
- Ye, L., Landen, W.O., and Eitenmiller, R.R. (2001). Simplified extraction procedure and HPLC determination for total vitamin E and β -carotene of reduced-fat mayonnaise. *J. Food Sci*, 66, 78-82.
- Yen, G.-C., Chang, Y.-C., and Su, S.-W. (2003). Antioxidant activity and active compounds of rice koji fermented with *Aspergillus candidus*. *Food Chem*, 83, 49-54.
- Yen, S.C.W., Pin-Der-Duh. (1996). Extraction and identification of antioxidant components from the leaves of mulberry (*Morus alba* L.). *J. Agri Food Chem*, 44, 1687-1690.
- Yoshida, Y., Niki, E. and Noguchi, N. (2003). Comparative study on the action of tocopherols and tocotrienols as antioxidant : chemical and physical effects. *Chemistry and Physics of Lipid*, 123, 63-75.
- Zadernowski, R., Naczki, M., and Amarowicz, R. (2003). Tocopherols in sea buckthorn (*Hippophaë rhamnoides* L.) berry oil. *J. Am. Oil Chem. Soc*, 80, 55-58.
- Zhou, Z., Robards, K., Helliwell, S., and Blanchard, C. (2002). Composition and functional properties of rice. *J. Food Sci and Tech*, 37, 849-868.