

REFERENCES

1. Mocarski ES J. Cytomegalovirus and their replication. In: Fields B, Knipe D, Howley P, editors. Fields Virology Vol 2. 3rd ed. Philadelphia; 1996. p. 2447-2523.
2. Boeckin M, Bowder R. Cytomegalovirus infection in marrow transplantation. In: Buckner CD, editor. Technical and Biological Components of Marrow Transplantation. Boston; 1995. p. 97-136.
3. Mendez J, Sia I, Paya C. Human cytomegalovirus. In: Lennette E, Smith T, editors. Laboratory Diagnosis of Viral Infection. New York; 1999. p. 361-372.
4. Pass RF. Cytomegalovirus. In: Knipe DM, Howley PM, Griffin DE, editors. Fields Virology Vol 2. 4th ed. Philadelphia; 2001. p. 2629-2705.
5. Gozlan J, Laporte JP, Lesage S, Labopin M, Najman A, Gorin NC, et al. Monitoring of cytomegalovirus infection and disease in bone marrow recipients by reverse transcription-PCR and comparison with PCR and blood and urine cultures. J Clin Microbiol 1996; 34(9): 2085-8.
6. Britt WJ, Mach M. Human cytomegalovirus glycoproteins. Intervirology 1996; 39 (5-6): 401-12.
7. Spaete RR, Gehrz RC, Landini MP. Human cytomegalovirus structural proteins. J Gen Virol 1994; 75(Pt 12): 3287-308.
8. Greenaway PJ, Oram JD, Downing RG, Patel K. Human cytomegalovirus DNA: BamHI, EcoRI and PstI restriction endonuclease cleavage maps. Gene 1982; 18(3): 355-60.
9. Hayward GS, Ambinder R, Ciuffo D, Hayward SD, LaFemina RL. Structural organization of human herpesvirus DNA molecules. J Invest Dermatol 1984; 83(1 Suppl): 29s-41s.
10. Oram JD, Downing RG, Akrigg A, Dollery AA, Duggleby CJ, Wilkinson GW, et al. Use of recombinant plasmids to investigate the structure of the human cytomegalovirus genome. J Gen Virol 1982; 59(Pt 1): 111-29.
11. Spector DH, Hock L, Tamashiro JC. Cleavage maps for human cytomegalovirus DNA strain AD169 for restriction endonucleases *EcoRI*, *BglII*, and *HindIII*. J Virol 1982; 42(2): 558-82.

12. Weststrate MW, Geelen JL, van der Noordaa J. Human cytomegalovirus DNA: physical maps for restriction endonucleases *Bgl*III, *Hind*III and *Xba*I. *J Gen Virol* 1980; 49(1): 1-21.
13. Colberg-Poley AM. Functional roles of immediate early proteins encoded by the human cytomegalovirus UL36-38, UL115-119, TRS1/IRS1 and US3 loci. *Intervirology* 1996; 39(5-6): 350-60.
14. Spector DH. Activation and regulation of human cytomegalovirus early genes. *Intervirology* 1996; 39(5-6): 361-77.
15. Stenberg RM. The human cytomegalovirus major immediate-early gene. *Intervirology* 1996; 39(5-6): 343-9.
16. Stinski MF, Thomsen DR, Stenberg RM, Goldstein LC. Organization and expression of the immediate early genes of human cytomegalovirus. *J Virol* 1983; 46(1): 1-14.
17. Ahn K, Angulo A, Ghazal P, Peterson PA, Yang Y, Fruh K. Human cytomegalovirus inhibits antigen presentation by a sequential multistep process. *Proc Natl Acad Sci U S A* 1996; 93(20): 10990-5.
18. Chee MS, Bankier AT, Beck S, Bohni R, Brown CM, Cerny R, et al. Analysis of the protein-coding content of the sequence of human cytomegalovirus strain AD169. *Curr Top Microbiol Immunol* 1990; 154: 125-69.
19. Iskenderian AC, Huang L, Reilly A, Stenberg RM, Anders DG. Four of eleven loci required for transient complementation of human cytomegalovirus DNA replication cooperate to activate expression of replication genes. *J Virol* 1996; 70(1): 383-92.
20. Stasiak PC, Mocarski ES. Transactivation of the cytomegalovirus ICP36 gene promoter requires the alpha gene product TRS1 in addition to IE1 and IE2. *J Virol* 1992; 66(2): 1050-8.
21. Goldmacher VS, Bartle LM, Skaletskaya A, Dionne CA, Kedersha NL, Vater CA, et al. A cytomegalovirus-encoded mitochondria-localized inhibitor of apoptosis structurally unrelated to Bcl-2. *Proc Natl Acad Sci U S A* 1999; 96(22): 12536-41.
22. Jones TR, Wiertz EJ, Sun L, Fish KN, Nelson JA, Ploegh HL. Human cytomegalovirus US3 impairs transport and maturation of major histocompatibility complex class I heavy chains. *Proc Natl Acad Sci U S A* 1996; 93(21): 11327-33.

23. Lundquist CA, Meier JL, Stinski MF. A strong negative transcriptional regulatory region between the human cytomegalovirus UL127 gene and the major immediate-early enhancer. *J Virol* 1999; 73(11): 9039-52.
24. Borysiewicz LK, Rodgers B, Morris S, Graham S, Sissons JG. Lysis of human cytomegalovirus infected fibroblasts by natural killer cells: demonstration of an interferon-independent component requiring expression of early viral proteins and characterization of effector cells. *J Immunol* 1985; 134(4): 2695-701.
25. Riddell SR, Greenberg PD. T cell therapy of human CMV and EBV infection in immunocompromised hosts. *Rev Med Virol* 1997; 7(3): 181-192.
26. Li CR, Greenberg PD, Gilbert MJ, Goodrich JM, Riddell SR. Recovery of HLA-restricted cytomegalovirus (CMV)-specific T-cell responses after allogeneic bone marrow transplant: correlation with CMV disease and effect of ganciclovir prophylaxis. *Blood* 1994; 83(7): 1971-9.
27. Bogner E, Radsak K, Stinski MF. The gene product of human cytomegalovirus open reading frame UL56 binds the pac motif and has specific nuclease activity. *J Virol* 1998; 72(3): 2259-64.
28. Krosky PM, Underwood MR, Turk SR, Feng KW, Jain RK, Ptak RG, et al. Resistance of human cytomegalovirus to benzimidazole ribonucleosides maps to two open reading frames: UL89 and UL56. *J Virol* 1998; 72(6): 4721-8.
29. Landini MP, Mach M. Searching for antibodies specific for human cytomegalovirus: is it diagnostically useful? When and how. *Scand J Infect Dis Suppl* 1995; 99: 18-23.
30. Paya CV, Smith TF, Ludwig J, Hermans PE. Rapid shell vial culture and tissue histology compared with serology for the rapid diagnosis of cytomegalovirus infection in liver transplantation. *Mayo Clin Proc* 1989; 64(6): 670-5.
31. Betts RF, Schmidt SG. Cytolytic IgM antibody to cytomegalovirus in primary cytomegalovirus infection in humans. *J Infect Dis* 1981; 143(6): 821-6.
32. Zanghellini F, Boppana SB, Emery VC, Griffiths PD, Pass RF. Asymptomatic primary cytomegalovirus infection: virologic and immunologic features. *J Infect Dis* 1999; 180(3): 702-7.

33. Beninga J, Kropff B, Mach M. Comparative analysis of fourteen individual human cytomegalovirus proteins for helper T cell response. *J Gen Virol* 1995; 76(Pt 1): 153-60.
34. Davignon JL, Clement D, Alriquet J, Michelson S, Davrinche C. Analysis of the proliferative T cell response to human cytomegalovirus major immediate-early protein (IE1): phenotype, frequency and variability. *Scand J Immunol* 1995; 41(3): 247-55.
35. He H, Rinaldo CR, Jr., Morel PA. T cell proliferative responses to five human cytomegalovirus proteins in healthy seropositive individuals: implications for vaccine development. *J Gen Virol* 1995;76(Pt 7): 1603-10.
36. Quinnan GV, Jr., Kirmani N, Rook AH, Manischewitz JF, Jackson L, Moreschi G, et al. Cytotoxic t cells in cytomegalovirus infection: HLA-restricted T- lymphocyte and non-T-lymphocyte cytotoxic responses correlate with recovery from cytomegalovirus infection in bone-marrow-transplant recipients. *N Engl J Med* 1982; 307(1): 7-13.
37. Berencsi K, Gonczol E, Endresz V, Kough J, Takeda S, Gyulay Z, et al. The N-terminal 303 amino acids of the human cytomegalovirus envelope glycoprotein B (UL55) and the exon 4 region of the major immediate early protein 1 (UL123) induce a cytotoxic T-cell response. *Vaccine* 1996; 14(5): 369-74.
38. Boppana SB, Britt WJ. Recognition of human cytomegalovirus gene products by HCMV-specific cytotoxic T cells. *Virology* 1996; 222(1): 293-6.
39. Borysiewicz LK, Hickling JK, Graham S, Sinclair J, Cranage MP, Smith GL, et al. Human cytomegalovirus-specific cytotoxic T cells. Relative frequency of stage-specific CTL recognizing the 72-kD immediate early protein and glycoprotein B expressed by recombinant vaccinia viruses. *J Exp Med* 1988; 168(3): 919-31.
40. Kern F, Surel IP, Faulhaber N, Frommel C, Schneider-Mergener J, Schonemann C, et al. Target structures of the CD8(+)-T-cell response to human cytomegalovirus: the 72-kilodalton major immediate-early protein revisited. *J Virol* 1999; 73(10): 8179-84.
41. Reusser P, Riddell SR, Meyers JD, Greenberg PD. Cytotoxic T-lymphocyte response to cytomegalovirus after human allogeneic bone marrow transplantation: pattern of recovery and correlation with cytomegalovirus infection and disease. *Blood* 1991; 78(5): 1373-80.

42. Walter EA, Greenberg PD, Gilbert MJ, Finch RJ, Watanabe KS, Thomas ED, et al. Reconstitution of cellular immunity against cytomegalovirus in recipients of allogeneic bone marrow by transfer of T-cell clones from the donor. *N Engl J Med* 1995; 333(16): 1038-44.
43. Jones TR, Hanson LK, Sun L, Slater JS, Stenberg RM, Campbell AE. Multiple independent loci within the human cytomegalovirus unique short region down-regulate expression of major histocompatibility complex class I heavy chains. *J Virol* 1995; 69(8): 4830-41.
44. Jones TR, Sun L. Human cytomegalovirus US2 destabilizes major histocompatibility complex class I heavy chains. *J Virol* 1997; 71(4): 2970-9.
45. Wiertz EJ, Jones TR, Sun L, Bogyo M, Geuze HJ, Ploegh HL. The human cytomegalovirus US11 gene product dislocates MHC class I heavy chains from the endoplasmic reticulum to the cytosol. *Cell* 1996; 84(5): 769-79.
46. Ahn K, Gruhler A, Galocha B, Jones TR, Wiertz EJ, Ploegh HL, et al. The ER-luminal domain of the HCMV glycoprotein US6 inhibits peptide translocation by TAP. *Immunity* 1997; 6(5): 613-21.
47. Hengel H, Koopmann JO, Flohr T, Muranyi W, Goulmy E, Hammerling GJ, et al. A viral ER-resident glycoprotein inactivates the MHC-encoded peptide transporter. *Immunity* 1997; 6(5): 623-32.
48. Lehner PJ, Karttunen JT, Wilkinson GW, Cresswell P. The human cytomegalovirus US6 glycoprotein inhibits transporter associated with antigen processing-dependent peptide translocation. *Proc Natl Acad Sci U S A* 1997; 94(13): 6904-9.
49. Karre K, Welsh RM. Viral decoy vetoes killer cell. *Nature* 1997; 386(6624): 446-7.
50. Reyburn HT, Mandelboim O, Vales-Gomez M, Davis DM, Pazmany L, Strominger JL. The class I MHC homologue of human cytomegalovirus inhibits attack by natural killer cells. *Nature* 1997; 386(6624): 514-7.
51. Bodaghi B, Jones TR, Zipeto D, Vita C, Sun L, Laurent L, et al. Chemokine sequestration by viral chemoreceptors as a novel viral escape strategy: withdrawal of chemokines from the environment of cytomegalovirus-infected cells. *J Exp Med* 1998; 188(5): 855-66.
52. Benedict CA, Butrovich KD, Lurain NS, Corbeil J, Rooney I, Schneider P, et al. Cutting edge: a novel viral TNF receptor superfamily member in virulent strains of human cytomegalovirus. *J Immunol* 1999; 162(12): 6967-70.

53. Zhu H, Shen Y, Shenk T. Human cytomegalovirus IE1 and IE2 proteins block apoptosis. *J Virol* 1995; 69(12): 7960-70.
54. Fish KN, Britt W, Nelson JA. A novel mechanism for persistence of human cytomegalovirus in macrophages. *J Virol* 1996; 70(3): 1855-62.
55. Spear GT, Lurain NS, Parker CJ, Ghassemi M, Payne GH, Saifuddin M. Host cell-derived complement control proteins CD55 and CD59 are incorporated into the virions of two unrelated enveloped viruses. Human T cell leukemia/lymphoma virus type I (HTLV-I) and human cytomegalovirus (HCMV). *J Immunol* 1995; 155(9): 4376-81.
56. Spiller OB, Morgan BP, Tufaro F, Devine DV. Altered expression of host-encoded complement regulators on human cytomegalovirus-infected cells. *Eur J Immunol* 1996; 26(7): 1532-8.
57. Ibanez CE, Schrier R, Ghazal P, Wiley C, Nelson JA. Human cytomegalovirus productively infects primary differentiated macrophages. *J Virol* 1991; 65(12): 6581-8.
58. Taylor-Wiedeman J, Sissons JG, Borysiewicz LK, Sinclair JH. Monocytes are a major site of persistence of human cytomegalovirus in peripheral blood mononuclear cells. *J Gen Virol* 1991; 72(Pt 9): 2059-64.
59. Sinzger C, Plachter B, Grefte A, The TH, Jahn G. Tissue macrophages are infected by human cytomegalovirus in vivo. *J Infect Dis* 1996; 173(1): 240-5.
60. Bonkowsky HL, Lee RV, Klatskin G. Acute granulomatous hepatitis. Occurrence in cytomegalovirus mononucleosis. *Jama* 1975; 233(12): 1284-8.
61. Stratta RJ, Shaefer MS, Markin RS, Wood RP, Kennedy EM, Langnas AN, et al. Clinical patterns of cytomegalovirus disease after liver transplantation. *Arch Surg* 1989; 124(12): 1443-9; discussion 1449-50.
62. Wingard JR, Mellits ED, Sostrin MB, Chen DY, Burns WH, Santos GW, et al. Interstitial pneumonitis after allogeneic bone marrow transplantation. Nine-year experience at a single institution. *Medicine (Baltimore)* 1988; 67(3): 175-86.
63. Boppana SB, Pass RF, Britt WJ, Stagno S, Alford CA. Symptomatic congenital cytomegalovirus infection: neonatal morbidity and mortality. *Pediatr Infect Dis J* 1992; 11(2): 93-9.

64. Morgello S, Cho ES, Nielsen S, Devinsky O, Petit CK. Cytomegalovirus encephalitis in patients with acquired immunodeficiency syndrome: an autopsy study of 30 cases and a review of the literature. *Hum Pathol* 1987; 18(3): 289-97.
65. Stagno S, Reynolds DW, Pass RF, Alford CA. Breast milk and the risk of cytomegalovirus infection. *N Engl J Med* 1980; 302(19): 1073-6.
66. Alford CA, Stagno S, Pass RF. Natural history of perinatal cytomegaloviral infection. *Ciba Found Symp* 1979; 77: 125-47.
67. Nesmith JD, Pass RF. Cytomegalovirus Infection in Adolescents. *Adolesc Med* 1995; 6(1): 79-90.
68. Murph JR, Baron JC, Brown CK, Ebelhack CL, Bale JF, Jr. The occupational risk of cytomegalovirus infection among day-care providers. *Jama* 1991; 265(5): 603-8.
69. Pass RF, Hutto C, Lyon MD, Cloud G. Increased rate of cytomegalovirus infection among day care center workers. *Pediatr Infect Dis J* 1990; 9(7): 465-70.
70. Pass RF, Boppana S. Cytomegalovirus. In: Jeffries DJ, Hudson CN, editors. *Viral infections in obstetrics and Gynaecology*. New York; 1999. p. 35-56.
71. Stagno S, Pass RF, Dworsky ME, Alford CA, Jr. Maternal cytomegalovirus infection and perinatal transmission. *Clin Obstet Gynecol* 1982; 25(3): 563-76.
72. Dworsky M, Yow M, Stagno S, Pass RF, Alford C. Cytomegalovirus infection of breast milk and transmission in infancy. *Pediatrics* 1983; 72(3): 295-9.
73. Asanuma H, Numazaki K, Nagata N, Hotsubo T, Horino K, Chiba S. Role of milk whey in the transmission of human cytomegalovirus infection by breast milk. *Microbiol Immunol* 1996; 40(3): 201-4.
74. Vochem M, Hamprecht K, Jahn G, Speer CP. Transmission of cytomegalovirus to preterm infants through breast milk. *Pediatr Infect Dis J* 1998; 17(1): 53-8.
75. Smith DM. Rhabdomyosarcoma of the head and neck. *Can J Otolaryngol* 1974; 3(4): 618-24.
76. Gleaves CA, Smith TF, Shuster EA, Pearson GR. Rapid detection of cytomegalovirus in MRC-5 cells inoculated with urine specimens by using low-speed centrifugation and monoclonal antibody to an early antigen. *J Clin Microbiol* 1984; 19(6): 917-9.

77. Griffiths PD, Panjwani DD, Stirk PR, Ball MG, Ganczakowski M, Blacklock HA, et al. Rapid diagnosis of cytomegalovirus infection in immunocompromised patients by detection of early antigen fluorescent foci. *Lancet* 1984; 2(8414): 1242-5.
78. Paya CV, Wold AD, Smith TF. Detection of cytomegalovirus infections in specimens other than urine by the shell vial assay and conventional tube cell cultures. *J Clin Microbiol* 1987; 25(5): 755-7.
79. Rabella N, Drew WL. Comparison of conventional and shell vial cultures for detecting cytomegalovirus infection. *J Clin Microbiol* 1990; 28(4): 806-7.
80. Grefte JM, van der Gun BT, Schmolke S, van der Giessen M, van Son WJ, Plachter B, et al. The lower matrix protein pp65 is the principal viral antigen present in peripheral blood leukocytes during an active cytomegalovirus infection. *J Gen Virol* 1992; 73(Pt 11): 2923-32.
81. Erice A, Holm MA, Gill PC, Henry S, Dirksen CL, Dunn DL, et al. Cytomegalovirus (CMV) antigenemia assay is more sensitive than shell vial cultures for rapid detection of CMV in polymorphonuclear blood leukocytes. *J Clin Microbiol* 1992; 30(11): 2822-5.
82. Iberer F, Tscheliessnigg K, Halwachs G, Rehak P, Wasler A, Petutschnigg B, et al. Definitions of cytomegalovirus disease after heart transplantation: antigenemia as a marker for antiviral therapy. *Transpl Int* 1996; 9(3): 236-42.
83. Patel R, Smith TF, Espy M, Wiesner RH, Krom RA, Portela D, et al. Detection of cytomegalovirus DNA in sera of liver transplant recipients. *J Clin Microbiol* 1994; 32(6): 1431-4.
84. Delgado R, Lumbreras C, Alba C, Pedraza MA, Otero JR, Gomez R, et al. Low predictive value of polymerase chain reaction for diagnosis of cytomegalovirus disease in liver transplant recipients. *J Clin Microbiol* 1992; 30(7): 1876-8.
85. Manez R, Kusne S, Rinaldo C, Aguado JM, St George K, Grossi P, et al. Time to detection of cytomegalovirus (CMV) DNA in blood leukocytes is a predictor for the development of CMV disease in CMV-seronegative recipients of allografts from CMV-seropositive donors following liver transplantation. *J Infect Dis* 1996; 173(5): 1072-6.
86. Drouet E, Boibieux A, Michelson S, Ecochard R, Biron F, Peyramond D, et al. Polymerase chain reaction detection of cytomegalovirus DNA in peripheral blood leukocytes as a predictor of cytomegalovirus disease in HIV-infected patients. *Aids* 1993; 7(5): 665-8.

87. Flood J, Drew WL, Miner R, Jekic-McMullen D, Shen LP, Kolberg J, et al. Diagnosis of cytomegalovirus (CMV) polyradiculopathy and documentation of *in vivo* anti-CMV activity in cerebrospinal fluid by using branched DNA signal amplification and antigen assays. *J Infect Dis* 1997; 176(2): 348-52.
88. Griscelli F, Barrois M, Chauvin S, Lastere S, Bellet D, Bourhis JH. Quantification of human cytomegalovirus DNA in bone marrow transplant recipients by real-time PCR. *J Clin Microbiol* 2001; 39(12): 4362-9.
89. Guiver M, Fox AJ, Mutton K, Mogulkoc N, Egan J. Evaluation of CMV viral load using TaqMan CMV quantitative PCR and comparison with CMV antigenemia in heart and lung transplant recipients. *Transplantation* 2001; 71(11): 1609-15.
90. Boeckh M, Boivin G. Quantitation of cytomegalovirus: methodologic aspects and clinical applications. *Clin Microbiol Rev* 1998; 11(3): 533-54.
91. GRAHAM RT. Polymerase chain reaction: basic principles and automation. In: McPHERSON MJ, QUIRKE P, TAYLOR GR, editors. *PCR 1*. Oxford: Oxford University Press; 1996. p. 1-13.
92. Danise A, Cinque P, Vergani S, Candino M, Racca S, De Bona A, et al. Use of polymerase chain reaction assays of aqueous humor in the differential diagnosis of retinitis in patients infected with human immunodeficiency virus. *Clin Infect Dis* 1997; 24(6): 1100-6.
93. Knox CM, Chandler D, Short GA, Margolis TP. Polymerase chain reaction-based assays of vitreous samples for the diagnosis of viral retinitis. Use in diagnostic dilemmas. *Ophthalmology* 1998; 105(1): 37-44; discussion 44-5.
94. Lee-Wing MW, Hodge WG, Diaz-Mitoma F. The prevalence of herpes family virus DNA in the conjunctiva of patients positive and negative for human immunodeficiency virus using the polymerase chain reaction. *Ophthalmology* 1999; 106(2): 350-4.
95. Evans PC, Soin A, Wreghitt TG, Alexander GJ. Qualitative and semiquantitative polymerase chain reaction testing for cytomegalovirus DNA in serum allows prediction of CMV related disease in liver transplant recipients. *J Clin Pathol* 1998; 51(12): 914-21.
96. Arai S, Mangono M, Starr S, Spivack J. Optimization of conditions for detecting human cytomegalovirus DNA with nested PCR. In: Becker Y, Darai G, editors. *PCR: Protocols for*

Diagnosis of Human and Animal Viruses Disease. Philadelphia: Heidelberg : Springer-Verlag; 1995. p. 205-13.

97. Grove and Newell. (1950). The Human Eye. From Animal Biology [Online]. Available: <http://www.micrographia.com/tutorial/micbasic/micbpt02/micb0200/hu446gan.htm> [2003, September 15].
98. Pan L, Milligan L, Michaeli J, Cesarman E, Knowles DM. Polymerase chain reaction detection of Kaposi's sarcoma-associated herpesvirus-optimized protocols and their application to myeloma. *J Mol Diagn* 2001; 3(1): 32-8.
99. Limaye AP, Huang ML, Leisenring W, Stensland L, Corey L, Boeckh M. Cytomegalovirus (CMV) DNA load in plasma for the diagnosis of CMV disease before engraftment in hematopoietic stem-cell transplant recipients. *J Infect Dis* 2001; 183(3): 377-82.
100. Machida U, Kami M, Fukui T, Kazuyama Y, Kinoshita M, Tanaka Y, et al. Real-time automated PCR for early diagnosis and monitoring of cytomegalovirus infection after bone marrow transplantation. *J Clin Microbiol* 2000; 38(7): 2536-42.
101. Boriskin YS, Fuller K, Powles RL, Vipond IB, Rice PS, Booth JC, et al. Early detection of cytomegalovirus (CMV) infection in bone marrow transplant patients by reverse transcription-PCR for CMV spliced late gene UL21.5: a two site evaluation. *J Clin Virol* 2002; 24(1-2): 13-23.
102. Myerson D, Lingenfelter PA, Gleaves CA, Meyers JD, Bowden RA. Diagnosis of cytomegalovirus pneumonia by the polymerase chain reaction with archived frozen lung tissue and bronchoalveolar lavage fluid. *Am J Clin Pathol* 1993; 100(4): 407-13.
103. Brytting M, Sundqvist VA, Stalhandske P, Linde A, Wahren B. Cytomegalovirus DNA detection of an immediate early protein gene with nested primer oligonucleotides. *J Virol Methods* 1991; 32(2-3): 127-38.
104. Ando Y, Terao K, Narita M, Oguchi Y, Sata T, Iwasaki T. Quantitative analyses of cytomegalovirus genome in aqueous humor of patients with cytomegalovirus retinitis. *Jpn J Ophthalmol* 2002; 46(3): 254-60.
105. Bachmann B, Luke W, Hunsmann G. Improvement of PCR amplified DNA sequencing with the aid of detergents. *Nucleic Acids Res* 1990; 18(5): 1309.

106. Sarkar G, Kapelner S, Sommer SS. Formamide can dramatically improve the specificity of PCR. *Nucleic Acids Res* 1990; 18(24): 7465.
107. Burkardt HJ. Standardization and quality control of PCR analyses. *Clin Chem Lab Med* 2000; 38(2): 87-91.
108. Danise A, Cinque P, Vergani S, Candino M, Racca S, De Bona A, et al. Use of polymerase chain reaction assays of aqueous humor in the differential diagnosis of retinitis in patients infected with human immunodeficiency virus. *Clin Infect Dis* 1997; 24(6): 1100-6.
109. Fox GM, Crouse CA, Chuang EL, Pflugfelder SC, Cleary TJ, Nelson SJ, et al. Detection of herpesvirus DNA in vitreous and aqueous specimens by the polymerase chain reaction. *Arch Ophthalmol* 1991; 109(2):266-71.
110. de Boer JH, Verhagen C, Bruinenberg M, Rothova A, de Jong PT, Baarsma GS, et al. Serologic and polymerase chain reaction analysis of intraocular fluids in the diagnosis of infectious uveitis. *Am J Ophthalmol* 1996; 121(6): 650-8.
111. Garweg J, Fenner T, Schmitz H, Bohnke M. Detection of HCMV-DNA in aqueous humor for diagnostic confirmation of viral retinitis. *Fortschr Ophthalmol* 1991; 88(6): 735-9.
112. Fenner TE, Garweg J, Hufert FT, Boehnke M, Schmitz H. Diagnosis of human cytomegalovirus-induced retinitis in human immunodeficiency virus type 1-infected subjects by using the polymerase chain reaction. *J Clin Microbiol* 1991; 29(11): 2621-2.
113. Gerna G, Baldanti F, Sarasini A, Furione M, Percivalle E, Revello MG, et al. Effect of foscarnet induction treatment on quantitation of human cytomegalovirus (HCMV) DNA in peripheral blood polymorphonuclear leukocytes and aqueous humor of AIDS patients with HCMV retinitis. The Italian Foscarnet Study Group. *Antimicrob Agents Chemother* 1994; 38(1): 38-44.
114. Mitchell SM, Fox JD, Tedder RS, Gazzard BG, Lightman S. Vitreous fluid sampling and viral genome detection for the diagnosis of viral retinitis in patients with AIDS. *J Med Virol* 1994; 43(4): 336-40.
115. McCann JD, Margolis TP, Wong MG, Kuppermann BD, Luckie AP, Schwartz DM, et al. A sensitive and specific polymerase chain reaction-based assay for the diagnosis of cytomegalovirus retinitis. *Am J Ophthalmol* 1995; 120(2): 219-26.

116. Preiser W, Brauninger S, Schwerdtfeger R, Ayliffe U, Garson JA, Brink NS, et al. Evaluation of diagnostic methods for the detection of cytomegalovirus in recipients of allogeneic stem cell transplants. *J Clin Virol* 2001; 20(1-2): 59-70.
117. Boeckh M, Gallez-Hawkins GM, Myerson D, Zaia JA, Bowden RA. Plasma polymerase chain reaction for cytomegalovirus DNA after allogeneic marrow transplantation: comparison with polymerase chain reaction using peripheral blood leukocytes, pp65 antigenemia, and viral culture. *Transplantation* 1997; 64(1): 108-13.
118. Michaelides A, Facey D, Spelman D, Wesselingh S, Kotsimbos T. HCMV DNA detection and quantitation in the plasma and PBL of lung transplant recipients: COBAS Amplicor HCMV monitor test versus in-house quantitative HCMV PCR. *J Clin Virol* 2003; 00:1-10.
119. Blok MJ, Goossens VJ, Vanherle SJ, Top B, Tacke N, Middeldorp JM, et al. Diagnostic value of monitoring human cytomegalovirus late pp67 mRNA expression in renal-allograft recipients by nucleic acid sequence-based amplification. *J Clin Microbiol* 1998; 36(5): 1341-6.