

REFERENCES

- Ada, AO, Suzen, SH, Iscan, M. Polymorphisms of cytochrome P450 1A1, glutathione S-transferases M1 and T1 in a Turkish population. *Toxicol Lett.* 2004;151(1):311-5.
- Alexandrie, AK, Sundberg, MI, Seidegard, J, Tornling, G, et al. Genetic susceptibility to lung cancer with special emphasis on CYP1A1 and GSTM1: a study on host factors in relation to age at onset, gender and histological cancer types. *Carcinogenesis.*1994;15(9): 1785-90.
- Alexandrov, K, Rojas, M, Geneste, O, Castegnaro, M, et al. An improved fluorometric assay for dosimetry of benzo(a)pyrene diol-epoxide-DNA adducts in smokers' lung: comparisons with total bulky adducts and aryl hydrocarbon hydroxylase activity. *Cancer Res.*1992;52(22): 6248-53.
- Arand, M, Muhlbauer, R, Hengstler, J, Jager, E, et al. A multiplex polymerase chain reaction protocol for the simultaneous analysis of the glutathione S-transferase GSTM1 and GSTT1 polymorphisms. *Anal Biochem.*1996; 236(1): 184-6.
- Arnhold, J. Properties, functions, and secretion of human myeloperoxidase. *Biochemistry (Mosc).*2004; 69(1): 4-9.
- Babior, BM. Phagocytes and oxidative stress. *Am J Med.*2000; 109(1): 33-44.
- Beckman, G, Birgander, R, Sjalander, A, Saha, N, et al. Is p53 polymorphism maintained by natural selection? *Hum Hered.*1994; 44(5): 266-70.
- Benhamou, S, Lee, WJ, Alexandrie, AK, Boffetta, P, et al. Meta- and pooled analyses of the effects of glutathione S-transferase M1 polymorphisms and smoking on lung cancer risk. *Carcinogenesis.*2002; 23(8): 1343-50.
- Birgander, R, Sjalander, A, Rannug, A, Alexandrie, AK, et al. P53 polymorphisms and haplotypes in lung cancer. *Carcinogenesis.*1995; 16(9): 2233-6.
- Boiteux, S, Radicella, JP. The human OGG1 gene: structure, functions, and its implication in the process of carcinogenesis. *Arch Biochem Biophys.* 2000; 377(1): 1-8.

- Bonner, MR, Bennett, WP, Xiong, W, Lan, Q, et al. Radon, secondhand smoke, glutathione-S-transferase M1 and lung cancer among women. *Int J Cancer*. 2006; 119(6): 1462-7.
- Bourdes, V, Boffetta, P, Pisani, P. Environmental exposure to asbestos and risk of pleural mesothelioma: review and meta-analysis. *Eur J Epidemiol*. 2000; 16(5): 411-7.
- Buttice, G, Duterque-Coquillaud, M, Basuyaux, JP, Carrere, S, et al. Erg, an Ets-family member, differentially regulates human collagenase1 (MMP1) and stromelysin1 (MMP3) gene expression by physically interacting with the Fos/Jun complex. *Oncogene*. 1996; 13(11):2297-306.
- Cajas-Salazar, N, Sierra-Torres, CH, Salama, SA, Zwischenberger, JB, et al. Combined effect of MPO, GSTM1 and GSTT1 polymorphisms on chromosome aberrations and lung cancer risk. *Int J Hyg Environ Health*. 2003; 206(6): 473-83.
- Carleton, T, Garrett, S. *Cellular Cancer Markers*. Totowa, New Jersey: Human Press; 1995.
- Cascorbi, I, Henning, S, Brockmoller, J, Gephart, J, et al. Substantially reduced risk of cancer of the aerodigestive tract in subjects with variant-463A of the myeloperoxidase gene. *Cancer Res*. 2000; 60(3): 644-9.
- Castell, JV, Donato, MT, Gomez-Lechon, MJ. Metabolism and bioactivation of toxicants in the lung. The in vitro cellular approach. *Exp Toxicol Pathol*. 2005; 57 Suppl 1: 189-204.
- Chambers, AF, Matrisian, LM. Changing views of the role of matrix metalloproteinases in metastasis. *J Natl Cancer Inst*. 1997; 89(17): 1260-70.
- Chen, J. Canadian individual risks of radon-induced lung cancer for different exposure profiles. *Can J Public Health*. 2005; 96(5): 360-3.
- Cho, EY, Hildesheim, A, Chen, CJ, Hsu, MM, et al. Nasopharyngeal carcinoma and genetic polymorphisms of DNA repair enzymes XRCC1 and hOGG1. *Cancer Epidemiol Biomarkers Prev*. 2003; 12(10): 1100-4.
- Coggan, M, Whitbread, L, Whittington, A, Board, P. Structure and organization of the human theta-class glutathione S-transferase and D-dopachrome tautomerase gene complex. *Biochem J*. 1998; 334 (Pt 3): 617-23.

- Coleman, W, Tsongalis, G. *The Molecular Basis of Human Cancer*. Totowa, New Jersey: Human Press; 2002.
- Cosma, G, Crofts, F, Currie, D, Wirgin, I, et al. Racial differences in restriction fragment length polymorphisms and messenger RNA inducibility of the human CYP1A1 gene. *Cancer Epidemiol Biomarkers Prev*.1993; 2(1): 53-7.
- Crofts, F, Taioli, E, Trachman, J, Cosma, GN, et al. Functional significance of different human CYP1A1 genotypes. *Carcinogenesis*.1994; 15(12): 2961-3.
- Dally, H, Gassner, K, Jager, B, Schmezer, P, et al. Myeloperoxidase (MPO) genotype and lung cancer histologic types: the MPO -463 A allele is associated with reduced risk for small cell lung cancer in smokers. *Int J Cancer*.2002; 102(5): 530-5.
- Detterbeck, F, Rivera, M, Socinski, M, Rosenman, J. *Diagnostic and Treatment of Lung Cancer*. Philadelphia: Saunders; 2001.
- Dherin, C, Radicella, JP, Dizdaroglu, M, Boiteux, S. Excision of oxidatively damaged DNA bases by the human alpha-hOgg1 protein and the polymorphic alpha-hOgg1(Ser326Cys) protein which is frequently found in human populations. *Nucleic Acids Res*.1999; 27(20): 4001-7.
- Dherin C, RJ, Dizdaroglu M, Boiteux S. . Excision of oxidatively damaged DNA bases by the human alpha-hOgg1 protein and the polymorphic alpha-hOgg1 (Ser326Cys) protein which is frequently found in human populations. *Nucl Acids Res*.1999; 27: 4001-7.
- Dialyna, IA, Miyakis, S, Georgatou, N, Spandidos, DA. Genetic polymorphisms of CYP1A1, GSTM1 and GSTT1 genes and lung cancer risk. *Oncol Rep*.2003; 10(6): 1829-35.
- Egeblad, M, Werb, Z. New functions for the matrix metalloproteinases in cancer progression. *Nat Rev Cancer*.2002; 2(3): 161-74.
- el-Zein, R, Zwischenberger, JB, Wood, TG, Abdel-Rahman, SZ, et al. Combined genetic polymorphism and risk for development of lung cancer. *Mutat Res*. 1997; 381(2): 189-200.
- El-Zein, RA, Zwischenberger, JB, Abdel-Rahman, SZ, Sankar, AB, et al. Polymorphism of metabolizing genes and lung cancer histology: prevalence of CYP2E1 in adenocarcinoma. *Cancer Lett*.1997; 112(1): 71-8.

- Fan, R, Wu, MT, Miller, D, Wain, JC, et al. The p53 codon 72 polymorphism and lung cancer risk. *Cancer Epidemiol Biomarkers Prev.*2000; 9(10): 1037-42.
- Fang, S, Jin, X, Wang, R, Li, Y, et al. Polymorphisms in the MMP1 and MMP3 promoter and non-small cell lung carcinoma in North China. *Carcinogenesis.* 2005; 26(2): 481-6.
- Feyler, A, Voho, A, Bouchardy, C, Kuokkanen, K, et al. Point: myeloperoxidase - 463G -> a polymorphism and lung cancer risk. *Cancer Epidemiol Biomarkers Prev.*2002;11(12):1550-4.
- Field, RW, Krewski, D, Lubin, JH, Zielinski, JM, et al. An overview of the North American residential radon and lung cancer case-control studies. *J Toxicol Environ Health A.*2006; 69(7): 599-631.
- Ford, JG, Li, Y, O'Sullivan, MM, Demopoulos, R, et al. Glutathione S-transferase M1 polymorphism and lung cancer risk in African-Americans. *Carcinogenesis.* 2000; 21(11): 1971-5.
- Gao, YT, Blot, WJ, Zheng, W, Ershow, AG, et al. Lung cancer among Chinese women. *Int J Cancer.*1987; 40(5): 604-9.
- Garte, S, Gaspari, L, Alexandrie, AK, Ambrosone, C, et al. Metabolic gene polymorphism frequencies in control populations. *Cancer Epidemiol Biomarkers Prev.*2001; 10(12): 1239-48.
- Geisler, SA, Olshan, AF. GSTM1, GSTT1, and the risk of squamous cell carcinoma of the head and neck: a mini-HuGE review. *Am J Epidemiol.*2001; 154(2): 95-105.
- Harris, GSaCC. Cancer Risk and Low-Penetrance Susceptibility Genes in Gene-Environment Interactions. *Journal of Clinical Oncology.*2000; 18(11): 2309-15.
- Hayashi, S, Watanabe, J, Nakachi, K, Kawajiri, K. Genetic linkage of lung cancer-associated MspI polymorphisms with amino acid replacement in the heme binding region of the human cytochrome P450IA1 gene. *J Biochem (Tokyo).*1991; 110(3): 407-11.
- Hayashi, S, Watanabe, J, Kawajiri, K. High susceptibility to lung cancer analyzed in terms of combined genotypes of P450IA1 and Mu-class glutathione S-transferase genes. *Jpn J Cancer Res.*1992; 83(8): 866-70.

- Hewitt, RE, Leach, IH, Powe, DG, Clark, IM, et al. Distribution of collagenase and tissue inhibitor of metalloproteinases (TIMP) in colorectal tumours. *Int J Cancer*.1991; 49(5): 666-72.
- Hung, RJ, Brennan, P, Canzian, F, Szeszenia-Dabrowska, N, et al. Large-scale investigation of base excision repair genetic polymorphisms and lung cancer risk in a multicenter study. *J Natl Cancer Inst*.2005; 97(8): 567-76.
- Hung, RJ, Hall, J, Brennan, P, Boffetta, P. Genetic polymorphisms in the base excision repair pathway and cancer risk: a HuGE review. *Am J Epidemiol*. 2005 ; 162(10): 925-42.
- Husgafvel-Pursiainen, K. Genotoxicity of environmental tobacco smoke: a review. *Mutat Res*.2004; 567(2-3): 427-45.
- Imai, K, Dalal, S, Chen, E, Downey, R. Human collagenase (matrix metalloproteinase-1) expression in the lungs of patients with emphysema *Am J Respir Crit Care Med* 2001; 163: 786-91.
- Irrarrazabal, CE, Rojas, C, Aracena, R, Marquez, C, et al. Chilean pilot study on the risk of lung cancer associated with codon 72 polymorphism in the gene of protein p53. *Toxicol Lett*.2003;144(1):69-76.
- Jin, X, Wu, X, Roth, JA, Amos, CI, et al. Higher lung cancer risk for younger African-Americans with the Pro/Pro p53 genotype. *Carcinogenesis*. 1995; 16(9): 2205-8.
- Johns, MB, Jr., Paulus-Thomas, JE. Purification of human genomic DNA from whole blood using sodium perchlorate in place of phenol. *Anal Biochem*. 1989; 180(2): 276-8.
- Kanamori, Y, Matsushima, M, Minaguchi, T, Kobayashi, K, et al. Correlation between expression of the matrix metalloproteinase-1 gene in ovarian cancers and an insertion/deletion polymorphism in its promoter region. *Cancer Res*. 1999; 59(17): 4225-7.
- Kawajiri, K, Nakachi, K, Imai, K, Yoshii, A, et al. Identification of genetically high risk individuals to lung cancer by DNA polymorphisms of the cytochrome P450IA1 gene. *FEBS Lett*.1990; 263(1): 131-3.
- Kellermann, G, Shaw, CR, Luyten-Kellerman, M. Aryl hydrocarbon hydroxylase inducibility and bronchogenic carcinoma. *N Engl J Med*.1973; 289(18): 934-7.

- Kelsey, KT, Wiencke, JK, Ward, J, Bechtold, W, et al. Sister-chromatid exchanges, glutathione S- transferase theta deletion and cytogenetic sensitivity to diepoxybutane in lymphocytes from butadiene monomer production workers. *Mutat Res.*1995; 335(3): 267-73.
- Ketterer, B, Harris, JM, Talaska, G, Meyer, DJ, et al. The human glutathione S-transferase supergene family, its polymorphism, and its effects on susceptibility to lung cancer. *Environ Health Perspect.*1992; 98: 87-94.
- Kihara, M, Noda, K, Kihara, M. Distribution of GSTM1 null genotype in relation to gender, age and smoking status in Japanese lung cancer patients. *Pharmacogenetics.* 1995; 5 Spec No: S74-9.
- Kim, JW, Lee, CG, Park, YG, Kim, KS, et al. Combined analysis of germline polymorphisms of p53, GSTM1, GSTT1, CYP1A1, and CYP2E1: relation to the incidence rate of cervical carcinoma. *Cancer.*2000; 88(9): 2082-91.
- Kim, RB, O'Shea, D. Interindividual variability of chlorzoxazone 6-hydroxylation in men and women and its relationship to CYP2E1 genetic polymorphisms. *Clin Pharmacol Ther.*1995; 57(6): 645-55.
- Kiyohara, C, Otsu, A, Shirakawa, T, Fukuda, S, et al. Genetic polymorphisms and lung cancer susceptibility: a review. *Lung Cancer.*2002; 37(3): 241-56.
- Kiyohara, C, Wakai, K, Mikami, H, Sido, K, et al. Risk modification by CYP1A1 and GSTM1 polymorphisms in the association of environmental tobacco smoke and lung cancer: a case-control study in Japanese nonsmoking women. *Int J Cancer.*2003; 107(1): 139-44.
- Kiyohara, C, Yoshimasu, K, Takayama, K, Nakanishi, Y. NQO1, MPO, and the risk of lung cancer: a HuGE review. *Genet Med.*2005; 7(7): 463-78.
- Ko, YC, Cheng, LS, Lee, CH, Huang, JJ, et al. Chinese food cooking and lung cancer in women nonsmokers. *Am J Epidemiol.*2000; 151(2): 140-7.
- Kohno, T, Shinmura, K, Tosaka, M, Tani, M, et al. Genetic polymorphisms and alternative splicing of the hOGG1 gene, that is involved in the repair of 8-hydroxyguanine in damaged DNA. *Oncogene.*1998; 16(25): 3219-25.
- Kohno, T, Kunitoh, H, Toyama, K, Yamamoto, S, et al. Association of the OGG1-Ser326Cys polymorphism with lung adenocarcinoma risk. *Cancer Sci.*2006; 97(8):724-8.

- Kongruttanachok, N, Sukdikul, S, Setavarin, S, Kerekhjanarong, V, et al. Cytochrome P450 2E1 polymorphism and nasopharyngeal carcinoma development in Thailand: a correlative study. *BMC Cancer*.2001; 1:4.
- Lai, C, Shields, PG. The role of interindividual variation in human carcinogenesis. *J Nutr*.1999; 129 (2S Suppl):552S-5S.
- Lam, WK. Lung cancer in Asian women-the environment and genes. *Respirology*. 2005; 10(4): 408-17.
- Landi, MT, Bertazzi, PA, Shields, PG, Clark, G, et al. Association between CYP1A1 genotype, mRNA expression and enzymatic activity in humans. *Pharmacogenetics*. 1994; 4(5): 242-6.
- Landi, S. Mammalian class theta GST and differential susceptibility to carcinogens: a review. *Mutat Res*.2000; 463(3): 247-83.
- Larsen, JE, Colosimo, ML, Yang, IA, Bowman, R, et al. CYP1A1 Ile462Val and MPO G-463A interact to increase risk of adenocarcinoma but not squamous cell carcinoma of the lung. *Carcinogenesis*.2006; 27(3): 525-32.
- Le Marchand, L, Sivaraman, L, Pierce, L, Seifried, A, et al. Associations of CYP1A1, GSTM1, and CYP2E1 polymorphisms with lung cancer suggest cell type specificities to tobacco carcinogens. *Cancer Res*.1998; 58(21): 4858-63.
- Le Marchand, L, Seifried, A, Lum, A, Wilkens, LR. Association of the myeloperoxidase -463G->a polymorphism with lung cancer risk. *Cancer Epidemiol Biomarkers Prev*.2000; 9(2): 181-4.
- Le Marchand, L, Donlon, T, Lum-Jones, A, Seifried, A, et al. Association of the hOGG1 Ser326Cys polymorphism with lung cancer risk. *Cancer Epidemiol Biomarkers Prev*.2002; 11(4): 409-12.
- Lee, AJ, Hodges, NJ, Chipman, JK. Interindividual variability in response to sodium dichromate-induced oxidative DNA damage: role of the Ser326Cys polymorphism in the DNA-repair protein of 8-oxo-7,8-dihydro-2'-deoxyguanosine DNA glycosylase 1. *Cancer Epidemiol Biomarkers Prev*. 2005; 14(2): 497-505.
- Liang, G, Pu, Y, Yin, L. Rapid detection of single nucleotide polymorphisms related with lung cancer susceptibility of Chinese population. *Cancer Lett*.2005; 223(2): 265-74.

- Liang, GY, Pu, YP, Yin, LH. [Studies of the genes related to lung cancer susceptibility in Nanjing Han population, China.]. *Yi Chuan*.2004; 26(5): 584-8.
- Liu, Y, Borchert, GL, Surazynski, A, Phang, JM. Proline oxidase, a p53-induced gene, targets COX-2/PGE(2) signaling to induce apoptosis and inhibit tumor growth in colorectal cancers. *Oncogene*.2008.
- London, SJ, Lehman, TA, Taylor, JA. Myeloperoxidase genetic polymorphism and lung cancer risk. *Cancer Res*.1997; 57(22):5001-3.
- London, SJ, Yuan, JM, Coetzee, GA, Gao, YT, et al. CYP1A1 I462V genetic polymorphism and lung cancer risk in a cohort of men in Shanghai, China. *Cancer Epidemiol Biomarkers Prev*.2000; 9(9): 987-91.
- Lu, W, Xing, D, Qi, J, Tan, W, et al. Genetic polymorphism in myeloperoxidase but not GSTM1 is associated with risk of lung squamous cell carcinoma in a Chinese population. *Int J Cancer*. 2002; 102(3): 275-9.
- Ma, JX, Zhang, KL, Liu, X, Ma, YL, et al. Concurrent expression of aryl hydrocarbon receptor and CYP1A1 but not CYP1A1 MspI polymorphism is correlated with gastric cancers raised in Dalian, China. *Cancer Lett*.2006; 240(2): 253-60.
- Marchand, LL, Wilkinson, GR, Wilkens, LR. Genetic and dietary predictors of CYP2E1 activity: a phenotyping study in Hawaii Japanese using chlorzoxazone. *Cancer Epidemiol Biomarkers Prev*.1999; 8(6): 495-500.
- Matsuzoe, D, Hideshima, T, Iwasaki, A, Yoneda, S, et al. Glutathione S-transferase mu1 null genotype is associated with K-ras gene mutation in lung adenocarcinoma among smokers. *Carcinogenesis*.2001; 22(8): 1327-30.
- Menzel, DB. The toxicity of air pollution in experimental animals and humans: the role of oxidative stress. *Toxicol Lett*.1994; 72(1-3): 269-77.
- Misra, RR, Tangrea, JA, Virtamo, J, Ratnasinghe, D, et al. Variation in the promoter region of the myeloperoxidase gene is not directly related to lung cancer risk among male smokers in Finland. *Cancer Lett*.2001; 164(2): 161-7.
- Mitacek, EJ, Brunnemann, KD, Polednak, AP, Hoffmann, D, et al. Composition of popular tobacco products in Thailand, and its relevance to disease prevention. *Prev Med*.1991; 20(6): 764-73.

- Moller, P, Folkmann, JK, Forchhammer, L, Brauner, EV, et al. Air pollution, oxidative damage to DNA, and carcinogenesis. *Cancer Lett.*2008; 266(1): 84-97.
- Mooney, LA, Bell, DA, Santella, RM, Van Bennekum, AM, et al. Contribution of genetic and nutritional factors to DNA damage in heavy smokers. *Carcinogenesis.* 1997; 18(3): 503-9.
- Murata, M, Tagawa, M, Kimura, M, Kimura, H, et al. Analysis of a germ line polymorphism of the p53 gene in lung cancer patients; discrete results with smoking history. *Carcinogenesis.* 1996; 17(2): 261-4.
- Murata, M, Watanabe, M, Yamanaka, M, Kubota, Y, et al. Genetic polymorphisms in cytochrome P450 (CYP) 1A1, CYP1A2, CYP2E1, glutathione S-transferase (GST) M1 and GSTT1 and susceptibility to prostate cancer in the Japanese population. *Cancer Lett.*2001; 165(2): 171-7.
- Murray, GI, Duncan, ME, O'Neil, P, McKay, JA, et al. Matrix metalloproteinase-1 is associated with poor prognosis in oesophageal cancer. *J Pathol.*1998; 185(3): 256-61.
- Murray, SL, Grubman, SA, Perrone, RD, Rojkind, M, et al. Matrix metalloproteinase activity in human intrahepatic biliary epithelial cell lines from patients with autosomal dominant polycystic kidney disease. *Connect Tissue Res.* 1996; 33(4): 249-56.
- Nakachi, K, Imai, K, Hayashi, S, Watanabe, J, et al. Genetic susceptibility to squamous cell carcinoma of the lung in relation to cigarette smoking dose. *Cancer Res.*1991; 51(19): 5177-80.
- Nakachi, K, Imai, K, Hayashi, S, Kawajiri, K. Polymorphisms of the CYP1 A1 and glutathione S-transferase genes associated with susceptibility to lung cancer in relation to cigarette dose in a Japanese population. *Cancer Res.*1993;53: 2994-9.
- Nakachi, K, Imai, K, Hayashi, S, Kawajiri, K. Polymorphisms of the CYP1A1 and glutathione S-transferase genes associated with susceptibility to lung cancer in relation to cigarette dose in a Japanese population. *Cancer Res.*1993; 53(13): 2994-9.
- Nakachi, K, Limtrakul, P, Sonklin, P, Sonklin, O, et al. Risk factors for lung cancer among Northern Thai women: epidemiological, nutritional, serological, and bacteriological surveys of residents in high- and low-incidence areas. *Jpn J Cancer Res.*1999; 90(11): 1187-95.

- Nazar-Stewart, V, Vaughan, TL, Stapleton, P, Van Loo, J, et al. A population-based study of glutathione S-transferase M1, T1 and P1 genotypes and risk for lung cancer. *Lung Cancer*. 2003; 40(3):247-58.
- Nebert, DW, McKinnon, RA, Puga, A. Human drug-metabolizing enzyme polymorphisms: effects on risk of toxicity and cancer. *DNA Cell Biol*. 1996; 15(4):273-80.
- Notani, P. Global variation in cancer incidence and mortality. *Curr Sci* 2001; 81 (5): 465-74.
- Okada, T, Kawashima, K, Fukushi, S, Minakuchi, T, et al. Association between a cytochrome P450 CYP1A1 genotype and incidence of lung cancer. *Pharmacogenetics*. 1994; 4(6): 333-40.
- Park, GT, Lee, OY, Kwon, SJ, Lee, CG, et al. Analysis of CYP2E1 polymorphism for the determination of genetic susceptibility to gastric cancer in Koreans. *J Gastroenterol Hepatol*. 2003; 18(11): 1257-63.
- Park, JH, Park, JM, Kim, EJ, Cha, SI, et al. Myeloperoxidase -463G>A polymorphism and risk of primary lung cancer in a Korean population. *Cancer Detect Prev*. 2006; 30(3): 257-61.
- Parkin, DM, Pisani, P, Ferlay, J. Estimates of the worldwide incidence of eighteen major cancers in 1985. *Int J Cancer*. 1993; 54(4): 594-606.
- Parkin, DM, Bray, F, Ferlay, J, Pisani, P. Global cancer statistics, 2002. *CA Cancer J Clin*. 2005 ; 55(2):74-108.
- Pauk, N, Kubik, A, Zatloukal, P, Krepela, E. Lung cancer in women. *Lung Cancer*. 2005; 48(1): 1-9.
- Paz-Elizur, T, Krupsky, M, Blumenstein, S, Elinger, D, et al. DNA repair activity for oxidative damage and risk of lung cancer. *J Natl Cancer Inst*. 2003; 95(17): 1312-9.
- Pearson, WR, Vorachek, WR, Xu, SJ, Berger, R, et al. Identification of class-mu glutathione transferase genes GSTM1-GSTM5 on human chromosome 1p13. *Am J Hum Genet*. 1993; 53(1):220-33.
- Pemble, S, Schroeder, KR, Spencer, SR, Meyer, DJ, et al. Human glutathione S-transferase theta (GSTT1): cDNA cloning and the characterization of a genetic polymorphism. *Biochem J*. 1994; 300 (Pt 1):271-6.

- Persson, I, Johansson, I, Bergling, H, Dahl, ML, et al. Genetic polymorphism of cytochrome P4502E1 in a Swedish population. Relationship to incidence of lung cancer. *FEBS Lett.*1993 ; 319(3):207-11.
- Persson, I, Johansson, I, Lou, YC, Yue, QY, et al. Genetic polymorphism of xenobiotic metabolizing enzymes among Chinese lung cancer patients. *Int J Cancer.*1999; 81(3):325-9.
- Petersen, DD, McKinney, CE, Ikeya, K, Smith, HH, et al. Human CYP1A1 gene: cosegregation of the enzyme inducibility phenotype and an RFLP. *Am J Hum Genet.*1991; 48(4):720-5.
- Piedrafita, FJ, Molander, RB, Vansant, G, Orlova, EA, et al. An Alu element in the myeloperoxidase promoter contains a composite SP1-thyroid hormone-retinoic acid response element. *J Biol Chem.*1996; 271(24):14412-20.
- Pisani, P, Srivatanakul, P, Randerson-Moor, J, Vipasrinimit, S, et al. GSTM1 and CYP1A1 polymorphisms, tobacco, air pollution, and lung cancer: a study in rural Thailand. *Cancer Epidemiol Biomarkers Prev.*2006; 15(4):667-74.
- Quinones, L, Lucas, D, Godoy, J, Caceres, D, et al. CYP1A1, CYP2E1 and GSTM1 genetic polymorphisms. The effect of single and combined genotypes on lung cancer susceptibility in Chilean people. *Cancer Lett.*2001; 174(1):35-44.
- Rebbeck, TR. Molecular epidemiology of the human glutathione S-transferase genotypes GSTM1 and GSTT1 in cancer susceptibility. *Cancer Epidemiol Biomarkers Prev.*1997; 6(9):733-43.
- Rutter, JL, Mitchell, TI, Buttice, G, Meyers, J, et al. A single nucleotide polymorphism in the matrix metalloproteinase-1 promoter creates an Ets binding site and augments transcription. *Cancer Res.*1998; 58(23):5321-5.
- Schabath, MB, Spitz, MR, Delclos, GL, Gunn, GB, et al. Association between asbestos exposure, cigarette smoking, myeloperoxidase (MPO) genotypes, and lung cancer risk. *Am J Ind Med.*2002; 42(1):29-37.
- Schroder, KR, Wiebel, FA, Reich, S, Dannappel, D, et al. Glutathione-S-transferase (GST) theta polymorphism influences background SCE rate. *Arch Toxicol.* 1995; 69(7):505-7.
- Seidegard, J, Vorachek, WR, Pero, RW, Pearson, WR. Hereditary differences in the expression of the human glutathione transferase active on trans-stilbene oxide are due to a gene deletion. *Proc Natl Acad Sci U S A.*1988; 85(19):7293-7.

Sgambato, A, Campisi, B, Zupa, A, Bochicchio, A, et al. Glutathione S-transferase (GST) polymorphisms as risk factors for cancer in a highly homogeneous population from southern Italy. *Anticancer Res.*2002; 22(6B):3647-52.

Shields, PG, Caporaso, NE, Falk, RT, Sugimura, H, et al. Lung cancer, race, and a CYP1A1 genetic polymorphism. *Cancer Epidemiol Biomarkers Prev.*1993 ; 2(5):481-5.

Shields, PG, Harris, CC. Cancer risk and low-penetrance susceptibility genes in gene-environment interactions. *J Clin Oncol.*2000; 18(11):2309-15.

Shimura, K, Yokota, J. The OGG1 gene encodes a repair enzyme for oxidatively damaged DNA and is involved in human carcinogenesis. *Antioxid Redox Signal.*2001; 3(4):597-609.

Simarak, S, de Jong, UW, Breslow, N, Dahl, CJ, et al. Cancer of the oral cavity, pharynx/larynx and lung in North Thailand: case-control study and analysis of cigar smoke. *Br J Cancer.*1977 ; 36(1):130-40.

Singto, Y. p53 Codon 72 Polymorphism and Selected Risk Factors in Cervical Cancer Patients at Srinagarind Hospital. Khon kaen: Khon kaen; 2004.

Sobti RC, SS, Joshi A, Jindal SK, Janmeja A. Genetic polymorphism of the CYP1A1, CYP2E1, GSTM1 and GSTT1 genes and lung cancer susceptibility in a North Indian population. *Mol Cell Biochem* 2004.2004; 266:1-9.

Sriplung, H, Sontipong, S, Martin, N, Wiangnon, S, et al. Cancer incidence in Thailand, 1995-1997. *Asian Pac J Cancer Prev.*2005; 6(3):276-81.

Su, L, Zhou, W, Park, S, Wain, JC, et al. Matrix metalloproteinase-1 promoter polymorphism and lung cancer risk. *Cancer Epidemiol Biomarkers Prev.*2005; 14(3):567-70.

Sugimura, H, Kohno, T, Wakai, K, Nagura, K, et al. hOGG1 Ser326Cys polymorphism and lung cancer susceptibility. *Cancer Epidemiol Biomarkers Prev.*1999; 8(8):669-74.

Suttajit, M, Mitacek, D, Martin, N. Toxicological study of traditional tobacco smoking related to high incidence of lung cancer in Chiang Mai. Abstract of Fourth Cancer Research Symposium, Faculty of Medicine, Chiang Mai University.1994:33-4.

- Sweeney, C, Nazar-Stewart, V, Stapleton, PL, Eaton, DL, et al. Glutathione S-transferase M1, T1, and P1 polymorphisms and survival among lung cancer patients. *Cancer Epidemiol Biomarkers Prev.*2003; 12(6):527-33.
- Szymanowska, A, Jassem, E, Dziadziuszko, R, Borg, A, et al. Increased risk of non-small cell lung cancer and frequency of somatic TP53 gene mutations in Pro72 carriers of TP53 Arg72Pro polymorphism. *Lung Cancer.*2006; 52(1):9-14.
- Tagawa, M, Murata, M, Kimura, H. Prognostic value of mutations and a germ line polymorphism of the p53 gene in non-small cell lung carcinoma: association with clinicopathological features. *Cancer Lett.*1998; 128(1):93-9.
- Tefre, T, Ryberg, D, Haugen, A, Nebert, DW, et al. Human CYP1A1 (cytochrome P(1)450) gene: lack of association between the Msp I restriction fragment length polymorphism and incidence of lung cancer in a Norwegian population. *Pharmacogenetics.*1991;1(1):20-5.
- Thaeomor, A. P53 codon 72 polymorphism, human papillomavirus infection and risk factors on pre-invasive cervical cancer of patients at Srinagarind hospital. Thailand., MoPH. Thailand. Public Health Statistics A.D.2003.Bangkok; 2004.
- Thomas, M, Kalita, A, Labrecque, S, Pim, D, et al. Two polymorphic variants of wild-type p53 differ biochemically and biologically. *Mol Cell Biol.*1999; 19(2):1092-100.
- Tiwawech, D, Srivatanakul, P, Karaluk, A, Ishida, T. The p53 codon 72 polymorphism in Thai nasopharyngeal carcinoma. *Cancer Lett.*2003; 198(1):69-75.
- To-Figueras, J, Gene, M, Gomez-Catalan, J, Galan, C, et al. Glutathione-S-Transferase M1 and codon 72 p53 polymorphisms in a northwestern Mediterranean population and their relation to lung cancer susceptibility. *Cancer Epidemiol Biomarkers Prev.*1996; 5(5):337-42.
- Tsuda, T, Mino, Y, Babazono, A, Shigemi, J, et al. A case-control study of lung cancer in relation to silica exposure and silicosis in a rural area in Japan. *Ann Epidemiol.*2002; 12(5):288-94.
- Uematsu, F, Kikuchi, H, Motomiya, M, Abe, T, et al. Association between restriction fragment length polymorphism of the human cytochrome P450IIE1 gene and susceptibility to lung cancer. *Jpn J Cancer Res.*1991; 82(3):254-6.

- Uematsu, F, Kikuchi, H, Motomiya, M, Abe, T, et al. Human cytochrome P450IIE1 gene: DraI polymorphism and susceptibility to cancer. *Tohoku J Exp Med.* 1992; 168(2):113-7.
- Uematsu, F, Ikawa, S, Kikuchi, H, Sagami, I, et al. Restriction fragment length polymorphism of the human CYP2E1 (cytochrome P450IIE1) gene and susceptibility to lung cancer: possible relevance to low smoking exposure. *Pharmacogenetics.* 1994; 4(2):58-63.
- van der Vaart, H, Postma, DS, Timens, W, ten Hacken, NH. Acute effects of cigarette smoke on inflammation and oxidative stress: a review. *Thorax.* 2004; 59(8):713-21.
- Van Schooten, FJ, Boots, AW, Knaapen, AM, Godschalk, RW, et al. Myeloperoxidase (MPO) -463G->A reduces MPO activity and DNA adduct levels in bronchoalveolar lavages of smokers. *Cancer Epidemiol Biomarkers Prev.* 2004;13(5):828-33.
- Vatanasapt, V, Sriamporn, S, Vatanasapt, P. Cancer control in Thailand. *Jpn J Clin Oncol.* 2002;32 Suppl:S82-91.
- Villard, PH, Seree, EM, Re, JL, De Meo, M, et al. Effects of tobacco smoke on the gene expression of the Cyp1a, Cyp2b, Cyp2e, and Cyp3a subfamilies in mouse liver and lung: relation to single strand breaks of DNA. *Toxicol Appl Pharmacol.* 1998; 148(2):195-204.
- Vinitketkumnuen, U, Kalayanamitra, K, Chewonarin, T, Kamens, R. Particulate matter, PM 10 & PM 2.5 levels, and airborne mutagenicity in Chiang Mai, Thailand. *Mutat Res.* 2002; 519 (1-2):121-31.
- Wang, J, Deng, Y, Li, L, Kuriki, K, et al. Association of GSTM1, CYP1A1 and CYP2E1 genetic polymorphisms with susceptibility to lung adenocarcinoma: a case-control study in Chinese population. *Cancer Sci.* 2003; 94(5):448-52.
- Wang, SL, Lee, H, Chen, KW, Tsai, KJ, et al. Cytochrome P4502E1 genetic polymorphisms and lung cancer in a Taiwanese population. *Lung Cancer.* 1999; 26(1):27-34.
- Wang, YC, Chen, CY, Chen, SK, Cherng, SH, et al. High frequency of deletion mutations in p53 gene from squamous cell lung cancer patients in Taiwan. *Cancer Res.* 1998; 58(2):328-33.

- Wang, YC, Chen, CY, Chen, SK, Chang, YY, et al. p53 codon 72 polymorphism in Taiwanese lung cancer patients: association with lung cancer susceptibility and prognosis. *Clin Cancer Res.*1999; 5(1):129-34.
- Warwick, A, Sarhanis, P, Redman, C, Pemble, S, et al. Theta class glutathione S-transferase GSTT1 genotypes and susceptibility to cervical neoplasia: interactions with GSTM1, CYP2D6 and smoking. *Carcinogenesis.*1994; 15(12):2841-5.
- Weiss, JM, Goode, EL, Ladiges, WC, Ulrich, CM. Polymorphic variation in hOGG1 and risk of cancer: a review of the functional and epidemiologic literature. *Mol Carcinog.*2005; 42(3):127-41.
- Wenham, RM, Calingaert, B, Ali, S, McClean, K, et al. Matrix metalloproteinase-1 gene promoter polymorphism and risk of ovarian cancer. *J Soc Gynecol Investig.*2003; 10(6):381-7.
- Weston, A, Godbold, JH. Polymorphisms of H-ras-1 and p53 in breast cancer and lung cancer: a meta-analysis. *Environ Health Perspect.*1997;1 05 Suppl 4: 919-26.
- Whittington, A, Vichai, V, Webb, G, Baker, R, et al. Gene structure, expression and chromosomal localization of murine theta class glutathione transferase mGSTT1-1. *Biochem J.*1999; 337 (Pt 1):141-51.
- Whyatt, RM, Bell, DA, Jedrychowski, W, Santella, RM, et al. Polycyclic aromatic hydrocarbon-DNA adducts in human placenta and modulation by CYP1A1 induction and genotype. *Carcinogenesis.*1998; 19(8):1389-92.
- Wiwatanadate, P, Voravons, R, Mahawana, T, wiwatanadate, M, et al. Lung Cancer Prevalence and Indoor Radon in Saraphi District, Chiang Mai, Thailand. *Intern.Med.*2001; 17 (1).
- Wu, X, Amos, CI, Kemp, BL, Shi, H, et al. Cytochrome P450 2E1 DraI polymorphisms in lung cancer in minority populations. *Cancer Epidemiol Biomarkers Prev.*1998; 7(1):13-8.
- Wyatt, CA, Coon, CI, Gibson, JJ, Brinckerhoff, CE. Potential for the 2G single nucleotide polymorphism in the promoter of matrix metalloproteinase to enhance gene expression in normal stromal cells. *Cancer Res.*2002; 62(24):7200-2.

- Yamane, A, Kohno, T, Ito, K, Sunaga, N, et al. Differential ability of polymorphic OGG1 proteins to suppress mutagenesis induced by 8-hydroxyguanine in human cell in vivo. *Carcinogenesis*. 2004; 25(9):1689-94.
- Yamazaki, H, Inui, Y, Yun, CH, Guengerich, FP, et al. Cytochrome P450 2E1 and 2A6 enzymes as major catalysts for metabolic activation of N-nitrosodialkylamines and tobacco-related nitrosamines in human liver microsomes. *Carcinogenesis*.1992; 13(10):1789-94.
- Yang, M, Choi, Y, Hwangbo, B, Lee, JS. Combined effects of genetic polymorphisms in six selected genes on lung cancer susceptibility. *Lung Cancer*.2007; 57(2):135-42.
- Yoon, KA, Kim, JH, Gil, HJ, Hwang, H, et al. CYP1B1, CYP1A1, MPO, and GSTP1 polymorphisms and lung cancer risk in never-smoking Korean women. *Lung Cancer*.2008; 60(1):40-6.
- Zhang, ZY, Fasco, MJ, Huang, L, Guengerich, FP, et al. Characterization of purified human recombinant cytochrome P4501A1-Ile462 and -Val462: assessment of a role for the rare allele in carcinogenesis. *Cancer Res*.1996; 56(17):3926-33.
- Zhu, Y, Spitz, MR, Lei, L, Mills, GB, et al. A single nucleotide polymorphism in the matrix metalloproteinase-1 promoter enhances lung cancer susceptibility. *Cancer Res*.2001; 61(21):7825-9.