

## REFERENCES

1. Chiu LZ, Schilling BK. A primer on weightlifting: From sport to sports training. *J Strength Cond Res.* 2005;27:42-8.
2. Garhammer J. Power production by Olympic weightlifters. *Med Sci Sports Exerc.* 1980;12:54-60.
3. Gisslén K, Öhberg L, Alfredson H. Is the chronic painful tendinosis tendon a strong tendon?: A case study involving an Olympic weightlifter with chronic painful jumper's knee. *Knee Surg Sports Traumatol Arthrosc.* 2006;14:897-902.
4. Calhoun G, Fry AC. Injury rates and profiles of elite competitive weightlifters. *J Athl Train.* 1999;34:232-8.
5. Neuro-Musculoskeletal and Pain Research Unit. Analysis of injuries and rehabilitation for excellence of Thai national weightlifters. Chiangmai: Chiang Mai University; 2007.
6. Draghi F, Danesino GM, Coscia D, Precerutti M, Pagani C. Overload syndromes of the knee in adolescents: Sonographic findings. *J Ultras.* 2008;11:151-7.
7. Jensen K, Di Fabio RP. Evaluation of eccentric exercise in treatment of patellar tendinitis. *Phys Ther.* 1989;69:211-6.
8. Visnes H, Bahr R. The evolution of eccentric training as treatment for patellar tendinopathy (jumper's knee): A critical review of exercise programmes. *Br J Sports Med.* 2007;41:217-23.

9. Cook JL, Khan KM, Purdam CR. Conservative treatment of patellar tendinopathy. *Phys Ther Sport*. 2001;2:54-65.
10. Khan KM, Maffulli N, B.D. C, Cook JL, Taunton JE. Patellar tendinopathy: Some aspects of basic science and clinical management. *Br J Sports Med*. 1998;32:346-55.
11. Kountouris A, Cook J. Rehabilitation of Achilles and patellar tendinopathies. *Best Pract Res Clin Rheumatol*. 2007;21:295-316.
12. Connell AT. Concepts for assessment and treatment of anterior knee pain related to altered spinal and pelvic biomechanics: A case report. *Man Ther*. 2008;23:560-3.
13. Panni AS, Tartarone M, Maffulli N. Patellar tendinopathy in athletes, outcome of nonoperative and operative management. *Am J Sports Med*. 2000;28:392-7.
14. Perez VE. Patellofemoral rehabilitation. *Operat Tech Orthop*. 2007;17:257-64.
15. Coleman S, Briffa K, Conroy H, Prince R, Carroll G, McQuade J. Short and medium-term effects of an education self-management program for individuals with osteoarthritis of the knee, designed and delivered by health professionals: A quality assurance study. *BMC Muscoskel Disord*. 2008;9:117-24.
16. Coleman S, Briffa NK, Carroll G, Inderjeeth C, Cook N, McQuade J. Effects of self-management, education and specific exercises, delivered by health professionals, in patients with osteoarthritis of the knee. *BMC Muscoskel Disord*. 2008;9:133-9.

17. Fitzgerald GK, Oatis C. Role of physical therapy in management of knee osteoarthritis. *Curr Opin Rheumatol*. 2004;16:143-7.
18. Hurley MV, Walsh NE, Mitchell HL, Pimm TJ, Patel A, Williamson E, et al. Clinical effectiveness of a rehabilitation program integrating exercise, self-management, and active coping strategies for chronic knee pain: A cluster randomized trial. *Arthritis Rheum*. 2007;57:1211-9.
19. Maurer BT, Stern AG, Kinossian B, Cook KD, Schumacher HR. Osteoarthritis of the knee: Isokinetic quadriceps exercise versus an educational intervention. *Arch Phys Med Rehabil*. 1999;80:1293-9.
20. Warsi A, LaValley MP, Wang PS, Avorn J, Solomon DH. Arthritis self-management education programs: A meta-analysis of the effect on pain and disability. *Arthritis Rheum*. 2003;48:2207-13.
21. Forssell M. The back school. *Spine*. 1981;6:104-6.
22. Adirima TA, Barouha A. Common orthopaedic injuries in young athletes. *Curr Paediatr*. 2006;16:205-10.
23. Llopis E, Padrón M. Anterior knee pain. *Eur J Radiol*. 2007;62:27-43.
24. Sharma P, Maffulli N. Tendon injury and tendinopathy: Healing and repair. *J Bone Joint Surg Am*. 2005;87:187-202.
25. Warden SJ, Brukner P. Patellar tendinopathy. *Clin Sports Med*. 2003;22:743-9.
26. Renstrom P, Woo S. Tendinopathy: A major medical problem in sport. In: Woo S, Renstrom P, Arnoczky S, editors. *Tendinopathy in Athletes*. Hoboken: John Wiley & Sons; 2007. p. 1-9.

27. Cook JL, Khan KM, Maffulli N, Purdam C. Overuse tendinosis, not tendinitis. *Phys Sports Med.* 2000;28:31-46
28. Giffin JR, Stanish WD. Overuse tendonitis and rehabilitation. *Can Fam Physician.* 1993;39:1762-9.
29. Gisslén K, Alfredson H. Neovascularisation and pain in jumper's knee: A prospective clinical and sonographic study in elite junior volleyball players. *Br J Sports Med.* 2005;39:423-8.
30. Peace KAL, Lee JC, Healy J. Imaging the infrapatellar tendon in the elite athlete. *Clin Radiol.* 2006;61:570-8.
31. Almekinders LC, Vellema JH, Weinhold PS. Strain patterns in the patellar tendon and the implications for patellar tendinopathy. *Knee Surg, Sports Traumatol, Arthrosc.* 2002;10 2-5.
32. Loudon JK, Wiesner D, Goist-Foley HL, Asjes C, Loudon KL. Intrarater reliability of functional performance tests for subjects with patellofemoral pain syndrome. *J Athl Train.* 2002;37:256-1.
33. Juhn MS. Patellofemoral pain syndrome: A review and guidelines for treatment. *Am Fam Physician.* 1999;60:2012-22.
34. Cabral CMN, Melim AMO, Sacco ICN, Marques AP. Physical therapy in patellofemoral syndrome patients: Comparison open and closed kinetic chain exercises. *Acta Ortop Bras.* 2008;16:180-5.
35. Gerbino PG. Adolescent anterior knee pain. *Oper Tech Sports Med.* 2006;14:203-11.

36. Escamilla RF, Zheng N, MacLeod TD, Edwards WB, Hreljac A, Fleisig GS, et al. Patellofemoral compressive force and stress during the forward and side lunges with and without a stride. *Clin Biomech.* 2008;23:1026-37.
37. Waryasz G, McDermott AY. Patellofemoral pain syndrome (PFPS): A systematic review of anatomy and potential risk factors. *Dyn Med.* 2008;7:1-14.
38. France RC. *Introduction To Sports Medicine & Athletic Training.* New York: Thomson Delmar Learning; 2003.
39. Medlar RC, Lyne ED. Sinding-Larsen-Johansson disease: Its etiology and natural history. *J Bone Joint Surg Am.* 1978;60:1113-6.
40. Khaund R, Flynn SH. Iliotibial band syndrome: A common source of knee pain. *Am Fam Physician.* 2005;71:1545-50.
41. Ellis R, Hing W, Reid D. Iliotibial band friction syndrome: A systematic review. *Man Ther.* 2007;12:200-8.
42. Bily W, Trimmel L, Mödlin M, Kaider A, Kern H. Training program and additional eccentric muscle stimulation for patellofemoral pain syndrome: A pilot study. *Arch Phys Med Rehabil.* 2008;89:1230-6.
43. Boling MC, Bolgla LA, Mattacola CG, Uhl TL, Hosey RG. Outcomes of a weight-bearing rehabilitation program for patients diagnosed with patellofemoral pain syndrome. *Arch Phys Med Rehabil.* 2006;87:1428-35.
44. Kongsgaard M, Aagaard P, Roikjaer S, Olsen D, Jensen M. Decline eccentric squats increases patellar tendon loading compared to standard eccentric squats. *Clin Biomech.* 2006;21:748-54.

45. Ebumea J, Bannister G. The McConnell regimen versus isometric quadriceps exercises in the management of anterior knee pain: A randomised prospective controlled trial. *Knee*. 1996;3:151-3.
46. Cannell LJ, Taunton JE, Clement DB, Smith C, Khan KM. A randomised clinical trial of the efficacy of drop squats or leg extension/leg curl exercises to treat clinically diagnosed jumper's knee in athletes: Pilot study. *Br J Sports Med*. 2001;35:60-4.
47. Bahr R, Fossan B, Løken S. Surgical treatment compared with eccentric training for patellar tendinopathy (jumper's knee): A randomized, controlled trial. *J Bone Joint Surg Am*. 2006;88:1689-98.
48. Purdam CR, Johnsson P, Alfredson H, Lorentzon R, Cook JL, Khan KM. A pilot study of the eccentric decline squat in the management of painful chronic patellar tendinopathy. *Br J Sports Med*. 2004;38:395-7.
49. Potach DH, Chu DA. Plyometric Training. In: Baechle TR, Earle RW, editors. *Essentials of Strength Training and Conditioning*. 2<sup>nd</sup> ed. Champaign: Human Kinetics; 2000.
50. Chimera NJ, Swanik KA, Swanik CB, Straub SJ. Effects of plyometric training on muscle-activation strategies and performance in female athletes. *J Athl Train*. 2004;39:24-31.
51. Vescovi JD. Plyometric, Speed, and Agility Exercise Prescription. In: Chandler TJ, Brown LE, editors. *Conditioning for Strength and Human Performance*. Philadelphia: Wolters Kluwer Health/Lippincott Williams & Wilkins; 2008. p. 306-45.

52. Fleck SJ, Kraemer WJ. Designing Resistance Training Programs. 3<sup>rd</sup> ed. Champaign: Human Kinetics; 2004.
53. McNeely E. Introduction to plyometrics: Converting strength to power. NSCA PTJ. 2007;6:19-22.
54. Baechle TR, Earle RW. Essentials of Strength Training and Conditioning. 2<sup>nd</sup> ed. Champaign: Human Kinetics; 2000.
55. Faigenbaum AD, McFarland JM, Keiper FB, Tevlin W, Ratamess NA, Kang J, et al. Effects of a short-term plyometric and resistance training program on fitness performance in boys age 12 to 15 years. J Sports Sci Med. 2007;6:519-25.
56. Roddy E, Zhang W, Doherty M. Aerobic walking or strengthening exercise for osteoarthritis of the knee?: A systematic review. Ann Rheum Dis. 2005;64:544-8.
57. Bird SB, Dickson EW. Clinically significant changes in pain along the visual analog scale. Ann Emerg Med. 2001;38:639-43.
58. Perry JD. Exercise, injury and chronic inflammatory lesions. Br Med Bull. 1992;48:668-82.
59. Jonsson P, Alfredson H. Superior results with eccentric compared to concentric quadriceps training in patients with jumper's knee: A prospective randomised study. Br J Sports Med. 2005;39:847-50.
60. Young MA, Cook JL, Purdam CR, Kiss ZS, Alfredson H. Eccentric decline squat protocol offers superior results at 12 months compared with traditional

- eccentric protocol for patellar tendinopathy in volleyball players. *Br J Sports Med.* 2005;39:102-5.
61. Visnes H, Hoksrud A, Cook J, Bahr R. No effect of eccentric training on jumper's knee in volleyball players during the competitive season: A randomized clinical trial. *Clin J Sport Med.* 2005;15:227-34.
62. Langberg H, Ellingsgaard H, Madsen T, Jansson J, Magnusson SP, Aagaard P, et al. Eccentric rehabilitation exercise increases peritendinous type I collagen synthesis in humans with Achilles tendinosis. *Scand J Med Sci Sports.* 2007;17:61-6.
63. Mafi N, Lorentzon R, Alfredson H. Superior short-term results with eccentric calf muscle training compared to concentric training in a randomized prospective multicenter study on patients with chronic Achilles tendinosis. *Knee Surg, Sports Traumatol, Arthrosc.* 2001;9:42-7.
64. Wasielewski NJ, Kotsko KM. Does eccentric exercise reduce pain and improve strength in physically active adults with symptomatic lower extremity tendinosis?: A systematic review. *J Athl Train.* 2007;42:409-21.
65. Pufe T, Petersen W, Kurz B, Tsokos M, Tillmann B, Mentlein R. Mechanical factors influence the expression of endostatin--an inhibitor of angiogenesis--in tendons. *J Orthop Res.* 2003;21:610-6.
66. Shalabi A, Kristoffersen-Wiberg M, Aspelin P, Movin T. Immediate Achilles tendon response after strength training evaluated by MRI. *Med Sci Sports Exerc.* 2004;36:1841-6.



67. Portney LG, Wakins MP. Foundations of Clinical Research: Applications to Practice. 2<sup>nd</sup> ed. New Jersey: Prentice Hall Health; 2000.
68. Frohm A, Saartok T, Halvorsen K, Renstrom P. Eccentric treatment for patellar tendinopathy: A prospective randomised short-term pilot study of two rehabilitation protocols. *Br J Sports Med.* 2007;41:451-7
69. Lord BA, Parsell B. Measurement of pain in the prehospital setting using a visual analog scale. *Prehosp Disaster Med.* 2003;18:353-8.
70. Visentini PJ, Khan KM, Cook JL, Kiss ZS, Harcourt PR, Wark JD. The VISA score: An index of severity of symptoms in patients with jumper's knee (patellar tendinosis). *J Sci Med Sport.* 1998;1:22-8.