

REFERENCES

1. Järup L, Åkesson A. Current status of cadmium as an environmental health problem. *Toxicol Appl Pharmacol* 2009;238(3):201-8.
2. Järup L, Rogenfelt A, Elinder C, Nogawa K, Kjellström T. Biological half-time of cadmium in the blood of workers after cessation of exposure. *Scand J Work Environ Health* 1983;9(4):327-31.
3. Kido T, Nordberg G. Cadmium-induced renal effects in the general environment. In: *Clinical Nephrotoxins*. Dordrecht: Kluwer Academic Publishers; 1998. p. 345-61.
4. Jin T, Nordberg G, Ye T, Bo M, Wang H, Zhu G, et al. Osteoporosis and renal dysfunction in a general population exposed to cadmium in China. *Environ Res* 2004;96(3):353-9.
5. Kasuya M. Recent epidemiological studies on itai-itai disease as a chronic cadmium poisoning in Japan. *Water Sci Technol* 2000;42(7-8):147-54.
6. Horiguchi H, Teranishi H, Niiya K, Aoshima K, Katoh T, Sakuragawa N, et al. Hypoproduction of erythropoietin contributes to anemia in chronic cadmium intoxication: clinical study on Itai-itai disease in Japan. *Arch Toxicol* 1994;68(10):632-6.
7. Kido T, Nogawa K, Honda R, Tsuritani I, Ishizaki M, Yamada Y, et al. The association between renal dysfunction and osteopenia in environmental cadmium-exposed subjects. *Environ Res* 1990;51(1):71-82.

8. Kasuya M, Aoshima K, Katoh T, Teranishi H, Horiguchi H, Kitagawa M, et al., editors. Natural history of Itai-itai disease: A long-term observation on the clinical and laboratory findings in patients with Itai-itai disease. Seventh International Cadmium Conference; 1992; New Orleans. Cadmium Association.
9. Nogawa K, Honda R, Kido T, Tsuritani I, Yamada Y, Ishizaki M, et al. A dose-response analysis of cadmium in the general environment with special reference to total cadmium intake limit. Environ Res 1989;48(1):7-16.
10. Nakashima K, Kobayashi E, Nogawa K, Kido T, Honda R. Concentration of cadmium in rice and urinary indicators of renal dysfunction. Occup Environ Med 1997;54(10):750-5.
11. Kido T, Nogawa K. Dose-response relationship between total cadmium intake and β_2 -microglobulinuria using logistic regression analysis. Toxicol Lett 1993;69(2):113-20.
12. Åkesson A, Bjellerup P, Lundh T, Lidfeldt J, Nerbrand C, Samsioe G, et al. Cadmium-Induced Effects on Bone in a Population-Based Study of Women. Environ Health Perspect 2006;114(6):830-4.
13. Järup L, Hellströmd L, Alfvén T, Carlsson MD, Grubb A, Persson B, et al. Low level exposure to cadmium and early kidney damage: the OSCAR study. Occup Environ Med 2000;57(10):668-72.
14. Sartor F, Rondia D, Claeys F, Staessen J, Lauwers R, Bernard A, et al. Impact of environmental cadmium pollution on cadmium exposure and body burden. Arch Environ Health 1992;47(5):347-53.

15. Staessen JA, Amery A, Bernard A, Bruaux P, Buchet JP, Claeys F, et al. Effects of exposure to cadmium on calcium metabolism: a population study. *Br J Ind Med* 1991;48:710-4.
16. Wang H, Zhu G, Shi Y, Weng S, Jin T, Kong Q, et al. Influence of environmental cadmium exposure on forearm bone density. *J Bone Miner Res* 2003;18(3):553-60.
17. Wu X, Jin T, Wang Z, Ye T, Kong Q, Nordberg G. Urinary calcium as a biomarker of renal dysfunction in a general population exposed to cadmium. *J Occup Environ Med* 2001;43(10):898-904.
18. Jin T, Nordberg G, Wu X, Ye T, Kong Q, Wang Z, et al. Urinary N-Acetyl- β -D - glucosaminidase isoenzymes as biomarker of renal dysfunction caused by cadmium in a general population. *Environ Res* 1999;81(2):167-73.
19. Padungtod C, Swadiwudhipong W, Nishijo M, Ruangyuttikarn W, Inud T, editors. *Health Risk Management for Cadmium Contamination in Thailand : Are Challenges Overcome?* ; 2006; Budapest, Hungary
20. Swadiwudhipong W, Limpatanachote P, Mahasakpan P, Somyos K, Padungtod C. Cadmium-exposed population in MaeSot district, Tak province: 1. Prevalence of high urinary cadmium levels in the adults. *J Med Assoc Thai* 2007;90:143-8.
21. Satarug S, Baker JR, Urbenjapol S, Haswell-Elkins M, Reilly PEB, Williams DJ, et al. A global perspective on cadmium pollution and toxicity in non-occupationally exposed population. *Toxicol Lett* 2003;137(1-2):65-83.

22. Limpatanachote P, Swaddiwudhipong W, Mahasakpan P, Krinratun S. Cadmium-exposed population in mae sot district tak province- 2. prevalence of renal dysfunction in the adults. *J Med Assoc Thai* 2009;92(10):1345-53.
23. Teeyakasem W, Nishijo M, Honda R, Satarug S, Swaddiwudhipong W, Ruangyuttikarn W. Monitoring of cadmium toxicity in a Thai population with high-level environmental exposure. *Toxicol Lett* 2007;169:185-95.
24. Nogawa K, Tsuritani I, Kido T, Honda R, Ishizaki M, Yamada Y. Serum vitamin D metabolites in cadmium-exposed persons with renal damage. *Int Arch Occup Environ Health* 1990;62(3):189-93.
25. Kido T, Honda R, Yamada Y, Tsuritani I, Ishizaki M, Nogawa K. α 1-Microglobulin determination in urine for the early detection of renal tubular dysfunctions caused by exposure to cadmium. *Toxicol Lett* 1985;24(2-3):195-201.
26. Bhattacharyya MH, Wilson AK, Rajan SS, Jonah M. Biochemical pathways in cadmium toxicity. In: Zalups RK, Koropatnick DJ, editors. *Molecular biology and toxicology of metals*. London: Taylor & Francis; 2000. p. 34-74.
27. Bridges CC, Zalups RK. Molecular and ionic mimicry and the transport of toxic metals. *Toxicol Appl Pharmacol* 2005;204(3):274-308.
28. Ferguson MA, Vaidya VS, Bonventre JV. Biomarkers of nephrotoxic acute kidney injury. *Toxicology* 2008;245(3):182-93.
29. Kobayashi E, Suwazono Y, Dochii M, Honda R, Nishijo M, Kido T, et al. Estimation of benchmark doses as threshold levels of urinary cadmium, based on excretion of β 2-microglobulin in cadmium-polluted and non-polluted regions in Japan. *Toxicol Lett* 2008;179(2):108-12.

30. Kido T, Honda R, Tsuritani I, Yamaya H, Ishizaki M, Yamada Y, et al. Progress of renal dysfunction in inhabitants environmentally exposed to cadmium. *Arch Environ Health* 1988;43(3):213-7.
31. Kido T, Nogawa K, Yamada Y, Honda R, Tsuritani I, Ishizaki M, et al. Osteopenia in inhabitants with renal dysfunction induced by exposure to environmental cadmium. *Int Arch Occup Environ Health* 1989;61(4):271-6.
32. Horiguchi H, Oguma E, Sasaki S, Miyamoto K, Ikeda Y, Machida M, et al. Environmental exposure to cadmium at a level insufficient to induce renal tubular dysfunction does not affect bone density among female Japanese farmers. *Environ Res* 2005;97(1):83-92.
33. Nogawa K, Tsuritani I, Kido T, Honda R, Yamada Y, Ishizaki M. Mechanism for bone disease found in inhabitants environmentally exposed to cadmium: decreased serum 1 α , 25-dihydroxyvitamin D level. *Int Arch Occup Environ Health* 1987;59(1):21-30.
34. Kido T, Honda R, Tsuritani I, Ishizaki M, Yamada Y, Nogawa K, et al. Assessment of cadmium-induced osteopenia by measurement of serum bone Gla protein, parathyroid hormone, and 1 α ,25-dihydroxyvitamin D. *J Appl Toxicol* 1991;11(3):161-6.
35. Engstrom A, Skerfving S, Lidfeldt J, Burgaz A, Lundh T, Samsioe Gr, et al. Cadmium-induced bone effect is not mediated via low serum 1,25-dihydroxy vitamin D. *Environ Res* 2009;109(2):188-92.
36. Aoshima K, Fan J, Cai Y, Katoh T, Teranishi H, Kasuya M. Assessment of bone metabolism in cadmium-induced renal tubular dysfunction by measurements of biochemical markers. *Toxicol Lett* 2003;136(3):183-92.

37. Omote S, Kido T, Nishijo M, Nakagawa H, Suwazono Y, Kobayashi E, et al. Urinary Type I Collagen Cross-Linked N-Telopeptides in Inhabitants 18 Years after Cessation of Exposure to Cadmium in Japan. *Bull Environ Contam Toxicol* 2006;76(2):187-94.
38. Schutte R, Nawrot TS, Richart T, Thijs L, Vanderschueren D, Kuznetsova T, et al. Bone resorption and environmental exposure to cadmium in women: a population study. *Environ Health Perspect* 2008;116(6):777-83.
39. Kido T, Honda R, Tsuritani I, Ishizaki M, Yamada Y, Nakagawa H. Serum levels of bone gla-protein in inhabitants exposed to environmental cadmium. . *Arch Environ Health* 1991;46(1):475-81.
40. Tsuritani I, Honda R, Ishizaki M, Yamada Y, Aoshima K, Kasuya M. Serum bone-type alkaline phosphatase activity in women living in a cadmium-polluted area. *Toxicol Lett* 1994;71(3):209-16.
41. Honda R, Tsuritani I, Noborisaka Y, Suzuki H, Ishizaki M, Yamada Y. Urinary cadmium excretion is correlated with calcaneal bone mass in Japanese women living in an urban area. *Environ Res* 2003;91(2):63-70.
42. Berns JS. Anemia due to Chronic Kidney Disease in the Elderly. In: Balducci L, Ershler WB, Bennett JM, editors. *Anemia in the Elderly*: Springer US; 2007. p. 61-73.
43. McClellan W, Aronoff SL, Bolton WK, Hood S, Lorber DL, Tang KL, et al. The prevalence of anemia in patients with chronic kidney disease. *Current Medical Research and Opinion* 2004;20(9):1501-10.
44. McFarlane SI, Chen S-C, Whaley-Connell AT, Sowers JR, Vassalotti JA, Salifu MO, et al. Prevalence and Associations of Anemia of CKD: Kidney Early

Evaluation Program (KEEP) and National Health and Nutrition Examination Survey (NHANES) 1999-2004. American Journal of Kidney Diseases 2008;51(4, Supplement 2):S46-S55.

45. Moriguchi J, Ezaki T, Tsukahara T, Furuki K, Fukui Y, Okamoto S, et al. Comparative evaluation of four urinary tubular dysfunction markers, with special references to the effects of aging and correction for creatinine concentration. *Toxicol Lett* 2003;143(3):279-90.
46. Tsukahara T, Ezaki T, Moriguchi J, Furuki K, Fukui Y, Ukai H, et al. No significant effect of iron deficiency on cadmium body burden or kidney dysfunction among women in the general population in Japan. *Int Arch Occup Environ Health* 2003;76(4):275-81.
47. Barrouillet MP, Ohayon-Courts C, Dubus I, L'Azou B, Nguyen Ba C. Influence of cadmium speciation for the evaluation of in vitro cadmium toxicity on LLC-PK₁ cells. *Toxicol In Vitro* 2001;15(4-5):525-9.
48. Belyaeva EA, Dymkowska D, Wieckowski MR, Wojtczak L. Reactive oxygen species produced by the mitochondrial respiratory chain are involved in Cd²⁺-induced injury of rat ascites hepatoma AS-30D cells. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* 2006;1757(12):1568-74.
49. Angle C, Thomas D, Swanson S. Osteotoxicity of cadmium and lead in HOS TE 85 and ROS 17/2.8 cells: relation to metallothionein induction and mitochondrial binding. *BioMetals* 1993;6(3):179-84.

50. Belyaeva EA, Dymkowska D, Wieckowski MR, Wojtczak L. Mitochondria as an important target in heavy metal toxicity in rat hepatoma AS-30D cells. *Toxicol Appl Pharmacol* 2008;231(1):34-42.
51. Gebhardt R. Prevention of cadmium-induced toxicity in liver-derived cells by the combination preparation Hepeel®. *Environ Toxicol Pharmacol* 2009;27(3):402-9.
52. Seibel MJ. Biochemical markers of bone turnover part I : biochemistry and variability. *Clin Biochem Rev* 2005;26:97-122.
53. Hannon RA, Eastell R. Bone markers and current laboratory assays. *Cancer Treatment Reviews* 2006;32(Supplement 1):7-14.
54. Grimaud E, Soubigou L, Couillaud S, Coipeau P, Moreau A, Passuti N, et al. Receptor Activator of Nuclear Factor κ B Ligand (RANKL)/Osteoprotegerin (OPG) Ratio Is Increased in Severe Osteolysis. *Am J Pathol* 2003;163(5):2021-31.
55. Humphrey EL, Williams JHH, Davie MWJ, Marshall MJ. Effects of dissociated glucocorticoids on OPG and RANKL in osteoblastic cells. *Bone* 2006;38(5):652-61.
56. Simmons RW, Pongsakul P, Saiyasitpanich D, Klinphoklap S. Elevated Levels of Cadmium and Zinc in Paddy Soils and Elevated Levels of Cadmium in Rice Grain Downstream of a Zinc Mineralized Area in Thailand: Implications for Public Health. *Environ Geochem Health* 2005;27(5):501-11.

57. Department of Provincial Administration MoI. รายงานสถิติจำนวนประชากร และบ้าน รายจังหวัด รายอำเภอ และรายตำบล ณ เดือน ธันวาคม พ.ศ. 2549 Bangkok; 2006 [updated 2006; cited 2010 22 June]; Available from: http://www.dopa.go.th/xstat/p4963_02.html.
58. Zarcinas B, Pongsakul P, McLaughlin M, Cozens G. Heavy metals in soils and crops in Southeast Asia 2. Thailand. Environ Geochem Health 2004;26(3):359-71.
59. Nogawa K, Kido T. Itai-Itai Disease and Health Effects of Cadmium. In: Chang LW, Magos L, Suzuki T, editors. Toxicology of Metals. Florida: CRC. Press.; 1996. p. 353-69.
60. Ogawa T, Kobayashi E, Okubo Y, Suwazono Y, Kido T, Nogawa K. Relationship among prevalence of patients with Itai-itai disease, prevalence of abnormal urinary findings, and cadmium concentrations in rice of individual hamlets in the Jinzu River basin, Toyama prefecture of Japan. Int J Environ Health Res 2004;14(4):243 - 52.
61. Nishijo M, Nagakawa H, Morikawa Y, Tabata M, Miura K, Kawano S, et al. Clinical courses and prognosis of itai-itai disease patients induced by chronic cadmium exposure. In: Collery PH, Corbella J, Domingo JL, Etienne JC, Llobet JM, editors. Metal Ions in Biology and Medicine. Paris: John Libbey Eurotext; 1996. p. 608-10.
62. Nriagu JO. Cadmium in the environment, Part II: Health effects. New York: John Wiley & Sons,; 1981.

63. Hayashi Y, Kobayashi E, Okubo Y, Suwazono Y, Kido T, Nogawa K. Excretion levels of urinary calcium and phosphorus among the inhabitants of Cd-polluted Kakehashi River basin of Japan. *Biol Trace Elem Res* 2003;91(1):45-55.
64. Nishijo M, Nakagawa H, Tabata M. Environmental exposure level and blood and urinary cadmium concentrations. *Environ Sci* 1995;3:125-35.
65. Nogawa K, Yamada Y, Honda R, Ishizaki M, Tsuritani I, Kawano S, et al. The relationship between itai-itai disease among inhabitants of the jinzu river basin and cadmium in rice. *Toxicol Lett* 1983;17(3-4):263-6.
66. Nordberg G, Jin T, Bernard A, Fierens S, Buchet JP, Ye T, et al. Low Bone Density and Renal Dysfunction Following Environmental Cadmium Exposure in China. *Ambio* 2002;31(6):478-81.
67. Alfvén T, Elinder C-G, Hellström L, Lagarde F, Järup L. Cadmium exposure and distal forearm fractures. *J Bone Miner Res* 2004;19(6):900-5.
68. Järup L, Alfvén, T., Persson, B., Toss, G., Elinder, C.G. Cadmium may be a risk factor for osteoporosis. *Occup Environ Med* 1998;55(7):435-9.
69. Staessen JA, Roels HA, Emelianov D, Kuznetsova T, Thijs L, Vangronsveld J, et al. Environmental exposure to cadmium, forearm bone density, and risk of fractures: prospective population study. *The Lancet* 1999;353(9159):1140-4.
70. Aylett BJ. The chemistry and bioinorganic chemistry of cadmium. In: Webb M, editor. *The Chemistry, biochemistry, and biology of cadmium*. New York . : Elsevier/North Holland Biomedical Press in Amsterdam; 1979. p. 1-44.
71. Peterson PJ, Alloway BJ. Cadmium in soils and vegetation. In: Webb M, editor. *The Chemistry, biochemistry, and biology of cadmium*. New York . : Elsevier/North Holland Biomedical Press in Amsterdam; 1979. p. 44-89.

72. Zalups RK, Ahmad S. Molecular handling of cadmium in transporting epithelia. *Toxicol Appl Pharmacol* 2003;186(3):163-88.
73. Raspotnicky B. Distribution and speciation of cadmium in natural waters. In: Nriagu JO, editor. Cadmium in the environment part 1: Ecological cycling Brisbane: John Wiley & Sons. Inc.; 1980.
74. Giesy JPJ. Distribution and speciation of cadmium in natural waters. In: Nriagu JO, editor. Cadmium in the environment part 1: Ecological cycling Brisbane: John Wiley & Sons. Inc.; 1980. p. 237-55.
75. Satarug S, Moore MR. Adverse Health Effects of Chronic Exposure to Low-Level Cadmium in Foodstuffs and Cigarette Smoke *Environ Health Perspect* 2004;112(10):1099-103.
76. Endo T, Kimura O, Sakata M. Bidirectional transport of cadmium across apical membrane of renal epithelial cell lines via H⁺-antiporter and inorganic anion exchanger. *Toxicology* 1998;131(2-3):183-92.
77. Endo T, Kimura O, Sakata M. Cadmium Uptake from Apical Membrane of LLC-PK Cells via Inorganic Anion Exchanger. *Pharmacol Toxicol* 1998;82(5):230-5.
78. Souza V, Bucio L, Gutiérrez-Ruiz MC. Cadmium uptake by a human hepatic cell line (WRL-68 cells). *Toxicology* 1997;120(3):215-20.
79. Faroone O, Ashizawa A, Wright S, Tucker P, Jenkins K, Ingberman L, et al. Toxicological profile for cadmium (Draft). Atlanta: Agency for Toxic Substances and Disease Registry; 2008.
80. Watanabe T, Zhang ZW, Moon CS, Shimbo S, Nakatsuka H, Matsuda-Inoguchi N, et al. Cadmium exposure of women in general populations in Japan during

- 1991-1997 compared with 1977-1981. *Int Arch Occup Environ Health* 2000;73(1):26-34.
81. ATSDR. Toxicological profile for cadmium. In: Registry AfTSaD, editor. Atlanta; 1999.
 82. Satarug S, Ujjin P, Vanavanitkul Y, Baker JR, Moore MR. Influence of body iron store status and cigarette smoking on cadmium body burden of healthy Thai women and men. *Toxicol Lett* 2004;148(3):177-85.
 83. Järup L, Berglund M, Elinder C, Nordberg G, Vahter M. Health effects of cadmium exposure a review of literature and a risk estimate. *Scand J Work Environ Health* 1998;24(suppl 1):1-52.
 84. WHO. Evaluation of certain food additives and contaminants. Thirty-third Report of the Joint FAO/WHO Expert Committee on Food Additives. Geneva: World Health Organization.; 1989. p. 28-31.
 85. Järup L, Persson B, Elinder C. Blood cadmium as an indicator of dose in a long-term follow-up of workers previously exposed to cadmium. *Scand J Work Environ Health* 1997;23(1):31-6.
 86. Bertin G, Averbeck D. Cadmium: cellular effects, modifications of biomolecules, modulation of DNA repair and genotoxic consequences (a review). *Biochimie* 2006;88(11):1549-59.
 87. Brzóska MM, Moniuszko-Jakoniuk J. Interactions between cadmium and zinc in the organism. *Food Chem Toxicol* 2001;39(10):967-80.
 88. Martelli A, Rousselet E, Dycke C, Bouron A, Moulis JM. Cadmium toxicity in animal cells by interference with essential metals. *Biochimie* 2006;88(11):1807-14.

89. Prozialeck WC, Lamar PC, Lynch SM. Cadmium alters the localization of N-cadherin, E-cadherin, and β -catenin in the proximal tubule epithelium. *Toxicol Appl Pharmacol* 2003;189(3):180-95.
90. Valko M, Morris H, Cronin MT. Metals, toxicity and oxidative stress. *Curr Med Chem* 2005;12(10):1161-208.
91. Long GJ. Cadmium perturbs calcium homeostasis in rat osteosarcoma (ros 17/2.8) cells; a possible role for protein kinase c. *Toxicol Lett* 1997;91(2):91-7.
92. Anke M, Latunde-Dada O, Arnhold W, Glei M, Anke S, Hartmann E, editors. The influence of age, sex and cadmium exposure on the ash, calcium, phosphorus, trace element and ultra trace element content in skeleton, kidneys and liver of humans. International conference on itai-itai disease, environmental cadmium pollution and countermeasures; 1998; Kanazawa. Eiko Laboratory.
93. Bhattacharyya MH. Cadmium osteotoxicity in experimental animals: Mechanisms and relationship to human exposures. *Toxicol Appl Pharmacol* 2009;238(3):258-65.
94. Mackie EJ. Osteoblasts: novel roles in orchestration of skeletal architecture. *Int J Biochem Cell B* 2003;35(9):1301-5.
95. Sims NA, Gooi JH. Bone remodeling: Multiple cellular interactions required for coupling of bone formation and resorption. *Semin Cell Dev Biol* 2008;19(5):444-51.
96. Teitelbaum SL. Bone Resorption by Osteoclasts. *Science* 2000;289(5484):1504-8.
97. Chen X, Zhu G, Gu S, Jin T, Shao C. Effects of cadmium on osteoblasts and osteoclasts in vitro. *Environ Toxicol Pharmacol* 2009;28(2):232-6.

98. Coonse KG, Coonts AJ, Morrison EV, Heggland SJ. Cadmium induces apoptosis in the human osteoblast-like cell line Saos-2. . J Toxicol Env Health 2007;70:575-81.
99. Lévesque M, Martineau C, Jumarie C, Moreau R. Characterization of cadmium uptake and cytotoxicity in human osteoblast-like MG-63 cells. Toxicol Appl Pharmacol 2008;231(3):308-17.
100. Kondoh M, Ogasawara S, Araragi S, Higashimoto M, Sato M. Cytochrome C release from mitochondria induced by cadmium. Health Sci 2001;47:78-82.
101. Chen L, Liu L, Huang S. Cadmium activates the mitogen-activated protein kinase (MAPK) pathway via induction of reactive oxygen species and inhibition of protein phosphatases 2A and 5. Free Radical Bio Med 2008;45:1035-44.
102. Li Y, Lim S-C. Cadmium-induced apoptosis of hepatocytes is not associated with death receptor-related caspase-dependent pathways in the rat. Environ Toxicol Pharmacol 2007;24:231–8.
103. Wilson AK, Cerny EA, Smith BD, Wagh A, Bhattacharyya MH. Effects of Cadmium on Osteoclast Formation and Activityin Vitro. Toxicol Appl Pharmacol 1996;140(2):451-60.
104. Nogawa K, Kobayashi E, Yamada Y, Honda R, Kido T, Tsuritani I, et al. Parathyroid hormone concentration in the serum of people with cadmium-induced renal damage. Int Arch Occup Environ Health 1984;54(3):187-93.
105. Tsuritani I, Honda R, Ishizaki M, Yamada Y. Impairment of vitamin D metabolism due to environmental cadmium exposure, and possible relevance to sex-related differences in vulnerability to the bone damage. J Toxicol Environ Health 1992;37:519-33.

106. Kido T, Nogawa K, Hochi Y, Hayano M, Honda R, Tsuritani I, et al. The renal handling of calcium and phosphorus in environmental cadmium-exposed subjects with renal dysfunction. *J Appl Toxicol* 1993;13(1):43-7.
107. Kobayashi E, Suwazono Y, Honda R, Dochii M, Nishijo M, Kido T, et al. Serial Follow-Up Study on Renal Handling of Calcium and Phosphorus After Soil Replacement in Cd-Polluted Rice Paddies Estimated Using a General Linear Mixed Model. *Biol Trace Elem Res* 2009;129(1):261-9.
108. Hiratsuka H, Katsuta O, Toyota N, Tsuchitani M, Umemura T, Marumo F. Chronic Cadmium Exposure-Induced Renal Anemia in Ovariectomized Rats. *Toxicol Appl Pharmacol* 1996;137(2):228-36.
109. Horiguchi H, Sato M, Konno N, Fukushima M. Long-term cadmium exposure induces anemia in rats through hypoinduction of erythropoietin in the kidneys. *Arch Toxicol* 1996;71(1):11-9.
110. Horiguchi H, Aoshima K, Oguma E, Sasaki S, Miyamoto K, Hosoi Y, et al. Latest status of cadmium accumulation and its effects on kidneys, bone, and erythropoiesis in inhabitants of the formerly cadmium-polluted Jinzu River Basin in Toyama, Japan, after restoration of rice paddies. *Int Arch Occup Environ Health* 2010.
111. Cumming RG, Mitchell P, Craig JC, Knight JF. Renal impairment and anaemia in a population-based study of older people. *Intern Med J* 2004;34(1-2):20-3.
112. Kido T, Sunaga K, Nishijo M, Nakagawa H, Kobayashi E, Nogawa K. The relation of individual cadmium concentration in urine with total cadmium intake in Kakehashi River basin, Japan. *Toxicol Lett* 2004;152(1):57-61.

113. Kido T, Nogawa K, Ohmichi M, Honda R, Tsuritani I, Ishizaki M, et al. Significance of urinary cadmium concentration in a Japanese population environmentally exposed to cadmium. *Arch Environ Health* 1992;47(3):196-202.
114. Nishijo M, Satarug S, Honda R, Tsuritani I, Aoshima K. The gender differences in health effects of environmental cadmium exposure and potential mechanisms. *Mol Cell Biochem* 2004;255(1):87-92.
115. Mueller PW, Price RG, Finn WF. New approaches for detecting thresholds of human nephrotoxicity using cadmium as an example. *Environ Health Perspect* 1998;106(5):227-30.
116. Schardijn GHC, Statis Van Eps LW. β 2-microglobulin: Its significance in the evaluation of renal function. *Kidney Int* 1987;32:635-41.
117. Price RG. Measurement of N-acetyl-beta-glucosaminidase and its isoenzymes in urine methods and clinical applications. *Eur J Clin Chem Clin Biochem* 1992;30(10):693-705.
118. Moriguchi J, Inoue Y, Kamiyama S, Horiguchi M, Murata K, Sakuragi S, et al. N-Acetyl- β -d-glucosaminidase (NAG) as the most sensitive marker of tubular dysfunction for monitoring residents in non-polluted areas. *Toxicol Lett* 2009;190(1):1-8.
119. Han WK, Bailly V, Abichandani R, Thadhani R, Bonventre JV. Kidney Injury Molecule-1 (KIM-1): A novel biomarker for human renal proximal tubule injury. *Kidney Int* 2002;62(1):237-44.

120. Vaidya VS, Ramirez V, Ichimura T, Bobadilla NA, Bonventre JV. Urinary kidney injury molecule-1: a sensitive quantitative biomarker for early detection of kidney tubular injury. *Am J Physiol Renal Physiol* 2006;290(2):F517-29.
121. Ichimura T, Hung CC, Yang SA, Stevens JL, Bonventre JV. Kidney injury molecule-1: a tissue and urinary biomarker for nephrotoxicant-induced renal injury. *Am J Physiol Renal Physiol* 2004;286(3):F552-63.
122. Panyamoon A NK, Nishijo M, Swaddiwudhipong W, Ruangyuttikarn W. . Detection of Urinary Kidney Injury Molecule-1 in a Chronic Cadmium Exposed Population, Mae Sot District, Tak Province. The 2nd National Conference in Toxicology; 2009 17-18 December 2009; Miracle Grand, Bangkok. 2009. p. 72-80.
123. Chantrel F, Agin A, Offner M, Koehl C, Moulin B, Hannedouche T. Comparison of cystatin C versus creatinine for detection of mild renal failure. *Clin Nephrol* 2000;54(5):374-81.
124. Mussap M, Vestra MD, Fioretto P, Saller A, Varagnolo M, Nosadini R, et al. Cystatin C is a more sensitive marker than creatinine for the estimation of GFR in type 2 diabetic patients. *Kidney Int* 2002;61(4):1453-61.
125. Uzun H, Ozmen Keles M, Ataman R, Aydin S, Kalender B, Uslu E, et al. Serum cystatin C level as a potentially good marker for impaired kidney function. *Clin Biochem* 2005;38(9):792-8.
126. Hoek FJ, Kemperman FAW, Krediet RT. A comparison between cystatin C, plasma creatinine and the Cockcroft and Gault formula for the estimation of glomerular filtration rate. *Nephrol Dial Transplant* 2003;18(10):2024-31.

127. Finney H, Newman DJ, Price CP. Adult reference ranges for serum cystatin C, creatinine and predicted creatinine clearance. *Ann Clin Biochem* 2000;37(1):49-59.
128. Gerbes AL, Gülberg V, Bilzer M, Vogeser M. Evaluation of serum cystatin C concentration as a marker of renal function in patients with cirrhosis of the liver. *Gut* 2002;50(1):106-10.
129. Roos JF, Doust J, Tett SE, Kirkpatrick CMJ. Diagnostic accuracy of cystatin C compared to serum creatinine for the estimation of renal dysfunction in adults and children--A meta-analysis. *Clin Biochem* 2007;40(5-6):383-91.
130. Papp AEd, Bone HG, Caulfield MP, Kagan R, Buinewicz A, Chen E, et al. A cross-sectional study of bone turnover markers in healthy premenopausal women. *Bone* 2007;40(5):1222-30.
131. Ebeling PR, Åkesson K. Role of chemical marker in the management of osteoporosis. *Best Pract Res Clin Rheumatol* 2001;15(3):385-400.
132. Delmas PD, Hardy P, Garnero P, Dain MP. Monitoring individual response to hormone replacement therapy with bone markers. *Bone* 2000;26(6):553-60.
133. Vasikaran SD, Glendenning P, Morris HA. The Role of Biochemical Markers of Bone Turnover in Osteoporosis Management in Clinical Practice. *Clin Biochem Rev* 2006;27(3):119-21.
134. Nishizawa Y, Nakamura T, Ohta H, Kushida K, Gorai I, Shiraki M, et al. Guidelines for the use of biochemical markers of bone turnover in osteoporosis (2004). *J Bone Miner Metab* 2005;23(2):97-104.

135. Rubinacci A, Melzi R, Zampino M, Soldarini A, Villa I. Total and Free Deoxypyridinoline after Acute Osteoclast Activity Inhibition. *Clin Chem* 1999;45(9):1510-6.
136. Paijitpraprakorn A, Udomthanatheera K, Udomthanatheera O, Mornthong T, Pungsuk S, Kongmetheekulrat C, et al. Source of cadmium contamination in soil in Amphoe Mae Sot, Tak Province. Bangkok: Department of Primary Industries and Mines, Ministry of Industry; 2006.
137. Glantz SA. How to test for trends. In: Glantz SA, editor. *Primer of Biostatistics*. 4 ed. Toronto: The McGraw-Hill Inc; 1997. p. 213-81.
138. Beaty RD, Kerber JD. *Concepts, Instrumentation and Techniques in Atomic Absorption Spectrophotometry*. Connecticut: Perkin Elmer, Inc.; 2002.
139. Biesalski H-K, Erhardt JG. Diagnosis of nutritional anemia-laboratory assessment of iron status. In: Kraemer K, Zimmermann MB, editors. *Nutritional anemia*. Basel: Sight and Life; 2007. p. 37-43.
140. Sugaya S, Nakanishi H, Tanzawa H, Sugita K, Kita K, Suzuki N. Down-regulation of SMT3A gene expression in association with DNA synthesis induction after X-ray irradiation in nevoid basal cell carcinoma syndrome (NBCCS) cells. *Mutat Res-Fund Mol M* 2005;578(1-2):327-32.
141. Bari CD, Dell'Accio F, Karystinou A, Guillot PV, Fisk NM, Jones EA, et al. A biomarker-based mathematical model to predict bone-forming potency of human synovial and periosteal mesenchymal stem cells. *Arthritis Rheum* 2008;58(1):240-50.

142. Monjo M, Lamolle SF, Lyngstadaas SP, Rønold HJ, Ellingsen JE. In vivo expression of osteogenic markers and bone mineral density at the surface of fluoride-modified titanium implants. *Biomaterials* 2008;29(28):3771-80.
143. Tabb MM, Sun A, Zhou C, Grønn F, Errandi J, Romero K, et al. Vitamin K₂ Regulation of Bone Homeostasis Is Mediated by the Steroid and Xenobiotic Receptor SXR. *J Biol Chem* 2003;278(45):43919-27.
144. Findlay D, Chehade M, Tsangari H, Neale S, Hay S, Hopwood B, et al. Circulating RANKL is inversely related to RANKL mRNA levels in bone in osteoarthritic males. *Arthritis Res Ther* 2008;10(1):R2.
145. Sirivarasai J, Kaojaren S, Wanakul W, Srisomerang P. Non-occupational determinants of cadmium and lead in blood and urine among a general population in Thailand. *Southeast Asian J Trop Med Public Health* 2002;33(1):180-7.
146. Suwazono Y, Kobayashi E, Okubo Y, Nogawa K, Kido T, Nakagawa H. Renal Effects of Cadmium Exposure in Cadmium Nonpolluted Areas in Japan. *Environ Res* 2000;84(1):44-55.
147. Kraemer K, Zimmermann MB. Nutritional anemia. Basel: Sight and Life; 2007.
148. Horiguchi H, Oguma E, Kayama F. Cadmium and cisplatin damage erythropoietin-producing proximal renal tubular cells. *Arch Toxicol* 2006;80(10):680-6.
149. Miyamoto T, Kobayashi E, Suwazono Y, Uetani M, Nishijo M, Miura K, et al. Changes in Urinary Excretions of Cadmium, Copper, and Zinc in Japanese

Workers. Archives of Environmental and Occupational Health 2006;61(5):215-22.

150. Shaikh ZA, Kido T, Kito H, Honda R, Nogawa K. Prevalence of metallothioneinuria among the population living in the Kakehashi river basin in Japan -- an epidemiological study. Toxicology 1990;64(1):59-69.
151. Aoshima K, Iwata K, Kasuya M. Environmental exposure to cadmium and effects on human health. Part 2. Bone and mineral metabolism in inhabitants of the cadmium-polluted Jinzu River basin in Toyama Prefecture. Jpn J Hyg 1988;43:864-71.
152. Yajima A, Ogawa Y, Takahashi HE, Tominaga Y, Inou T, Otsubo O. Changes of bone remodeling immediately after parathyroidectomy for secondary hyperparathyroidism. American Journal of Kidney Diseases 2003;42(4):729-38.
153. Coen G, Mantella D, Calabria S, Sardella D, Manni M, Fassino V, et al. Urinary deoxypyridinoline excretion for the evaluation of bone turnover in chronic renal failure. Am J Nephrol 2000;20:283-90.
154. Kobayashi E, Suwazono Y, Uetani M, Inaba T, Oishi M, Kido T, et al. Estimation of benchmark dose as the threshold levels of urinary cadmium, based on excretion of total protein, β_2 -microglobulin, and N-acetyl- β_2 -D-glucosaminidase in cadmium nonpolluted regions in Japan. Environ Res 2006;101(3):401-6.
155. Drüeke TB, Lacour B. Disorders of calcium, phosphate and magnesium metabolism. In: ed JR, editor. Comprehensive clinical nephrology. London: Mosby; 2003. p. 123-40.

156. Aoshima K, Katoh T, Teranishi H, Horiguchi H, Kasuya M. Abnormalities of calcium, phosphorus and vitamin D metabolism with proximal renal tubular dysfunction in subjects environmentally exposed to cadmium. *Nippon Eiseigaku Zasshi* 1993;47:1009-20.
157. Prayurahong B, Tungtrongchitr R, Chanjanakijskul S, Lertchavanakul A, Supawan V, Pongpaew P, et al. Vitamin B12, folic acid and haematological status in elderly Thais. *J Med Assoc Thai* 1993;76(2):71-8.
158. Limpatanachote P. Prevalence of renal dysfunction among persons with high urinary cadmium, Mae Sot district, Tak province. *Sawanpracharak Medical Journal* 2007;4:604-21.
159. Chittinandana A, Chailimpamontree W, Chaloeiphat P. Prevalence of Chronic Kidney Disease in Thai Adult Population. *J Med Assoc Thai* 2006;89(Suppl 2):S112-S20.
160. Sumboonnanonda A, Malasit P, Tanphaichitr VS, Ong-ajyooth S, Petrarat S, Vongjirad A. Renal tubular dysfunction in α -thalassemia. *Pediatric Nephrology* 2003;18(3):257-60.
161. Cetin T, Oktenli C, Ozgurtas T, Yenicesu M, Sanisoglu SY, Oguz Y, et al. Renal tubular dysfunction in β -thalassemia minor. *American Journal of Kidney Diseases* 2003;42(6):1164-8.
162. Horiguchi H. Anemia Induced by Cadmium Intoxication. *Nippon Eiseigaku Zasshi (Japanese Journal of Hygiene)* 2007;62(3):888-904.
163. ผ่องศรี_ savvyสม, จันทร์สว่าง_ทวีวนน_. โครงการพัฒนาการให้คำปรึกษาและดูแลโรคชาลัสซีเมียในสตรีตั้งครรภ์. ตาก: กลุ่มงานสุส蒂-นรีเวชกรรม โรงพยาบาลตากสิน; 2008.

164. Nishijo M, Tawara K, Honda R, Kuriwaki J-i, Nakagawa H, Tanebe K, et al. Cadmium and nutritional intake in pregnant Japanese women. *Toxicol Lett* 2004;148(3):171-6.
165. Swaddiwudhipong W, Mahasakpan P, Limpatanachote P, Krinratun S. Correlations of urinary cadmium with hypertension and diabetes in persons living in cadmium-contaminated villages in northwestern Thailand: A population study. *Environ Res* 2010;110(6):612-6.
166. Honda R, Swaddiwudhipong W, Nishijo M, Mahasakpan P, Teeyakasem W, Ruangyuttikarn W, et al. Cadmium induced renal dysfunction among residents of rice farming area downstream from a zinc-mineralized belt in Thailand. *Toxicol Lett* 2010;198(1):26-32.
167. Finn WF, Porter GA. Urinary biomarkers and nephrotoxicity. In: De Broe MEP, G. A. Bennett, W. M. Verpoorten, G. A. , editor. *Clinical nephrotoxins*. Boston: Kluwer academic publishers; 1998. p. 439-63.
168. Cho WCS. Proteomics Technologies and Challenges. *Genomics, Proteomics & Bioinformatics* 2007;5(2):77-85.
169. Bachi A, Bonaldi T. Quantitative proteomics as a new piece of the systems biology puzzle. *Journal of Proteomics* 2008;71(3):357-67.
170. Zhang X, Fang A, Riley CP, Wang M, Regnier FE, Buck C. Multi-dimensional liquid chromatography in proteomics--A review. *Anal Chim Acta*;664(2):101-13.

171. L'Azou B, Dubus I, Ohayon-Courts C, Labouyrie J-P, Perez L, Pouvreau C, et al. Cadmium induces direct morphological changes in mesangial cell culture. *Toxicology* 2002;179(3):233-45.
172. Long GJ. The Effect of Cadmium on Cytosolic Free Calcium, Protein Kinase C, and Collagen Synthesis in Rat Osteosarcoma (ROS 17/2.8) Cells. *Toxicol Appl Pharmacol* 1997;143(1):189-95.
173. Miyahara T, Takata M, Miyata M, Nagai M, Sugure A, Kozuka H, et al. Cadmium stimulates osteoclast-like multinucleated cell formation in mouse bone marrow cell cultures. *Bull Environ Contam Toxicol* 1991;47(2):283-7.
174. Li Z, Zhou Z, Saunders MM, Donahue HJ. Modulation of connexin43 alters expression of osteoblastic differentiation markers. *Am J Physiol Cell Physiol* 2006;290(4):C1248-55.
175. Ponader S, Brandt H, Vairaktaris E, von Wilmowsky C, Nkenke E, Schlegel KA, et al. In vitro response of hFOB cells to pamidronate modified sodium silicate coated cellulose scaffolds. *Colloids Surf B Biointerfaces* 2008;64(2):275-83.
176. Enjuanes A, Ruiz-Gaspà S, Peris P, Ozalla D, Álvarez L, Combalia A, et al. The effect of the alendronate on OPG/RANKL system in differentiated primary human osteoblasts. *Endocrine* 2010;37(1):180-6.
177. Zhang D, Yang YQ, Li XT, Fu MK. The expression of osteoprotegerin and the receptor activator of nuclear factor kappa B ligand in human periodontal ligament cells cultured with and without 1 α ,25-dihydroxyvitamin D3. *Arch Oral Biol* 2004;49(1):71-6.

178. Silvestrini G, Ballanti P, Sebastiani M, Leopizzi M, Di Vito M, Bonucci E. OPG and RANKL mRNA and protein expressions in the primary and secondary metaphyseal trabecular bone of PTH-treated rats are independent of that of SOST. *J Mol Histol* 2008;39(2):237-42.

179. Khosla S. Minireview: The OPG/RANKL/RANK System. *Endocrinology* 2001;142(12):5050-5.

180. Tat SK, Amiable N, Pelletier J-P, Boileau C, Lajeunesse D, Duval N, et al. Modulation of OPG, RANK and RANKL by human chondrocytes and their implication during osteoarthritis. *Rheumatology* 2009;48(12):1482-90.