

## BIBLIOGRAPHY

- Battagel JM. The Aetiological Factors in Class III Malocclusion. *Eur J Orthod* 1993; 15: 347-70.
- Bergman RT. Cephalometric Soft Tissue Facial Analysis. *Am J Orthod Dentofac Orthop* 1999; 116: 373-89.
- Bilodeau, JE. Vertical Considerations in Diagnosis and Treatment. A Surgical Orthodontic Case Report. *Am J Orthod Dentofac Orthop* 1995; 107: 91-100.
- Bjork A. Some Biological Aspects of Prognathism and Occlusion of the Teeth. *Acta Odont Scand* 1950; 9: 1-40.
- Burstone CJ, James RB, Legan H, Murphy GA, Norton LA. Cephalometrics for Orthognathic Surgery. *J Oral Surg* 1978; 36: 269-77.
- Chaiworawitkul M. A Comparison of the Cephalometric Profiles of Young Adults from Thailand and the U.K. [Thesis]. Sheffield: University of Sheffield, 1998.
- Chang HP, Kinoshita Z, Kawamoto T. A Study of the Growth Changes in Facial Configuration. *Eur J Orthod* 1993; 15: 493-501.
- Chareonvicha W. The Morphological Change During developing Class III Malocclusion in Female: a Cross-Sectional Study [Thesis]. Bangkok: Mahidol University, 1992.
- Chatkupt S, Sirinawin I, Patanaporn V, Bundech A, Apisariyakul S, Jotikasthira D, et al. Cephalometric Study in Northern Thai People. Research submitted to the Faculty of Dentistry, Chiang Mai: Chiang Mai University, 1987.
- Dechkunakorn S, Chaiwat J, Sawaengkit P, Anuwongnukor N. Facial Heights and Dentoalveolar Heights in Thai Adults with Normal Occlusion. *J Dent Assoc Thai* 1994; 44: 142-48.
- Dibbets JMH, Nolte K. Effect of Magnification on Lateral Cephalometric Studies. *Am J Orthod Dentofac Orthop* 2002; 122: 196-201.

- Droel R, Isaacson RJ. Some Relationships between the Glenoid Fossa Position Discrepancies. Am J Orthod 1972; 61: 64-78.
- El-Mangoury NH, Mostafa YA. Epidemiologic Panorama of Malocclusion. Angle Orthod 1990; 60: 207-14.
- Ellis E, McNamara JA Jr. Components of Adults Class III Malocclusion. J Oral Maxillofac Surg 1984; 42: 295-305.
- Enlow DH, Hans MG. Essentials of Facial Growth. Philadelphia: W.B.Saunders, 1996: 129-33.
- Frost DE, Fonseca RJ, Turvey TA, Hall DJ. Cephalometric Diagnosis and Surgical-Orthodontic Correction of Apertognathia. Am J Orthod 1980; 78: 657-69.
- Guyer EC, Ellis EE, McNamara JA Jr. Components of Class III Malocclusion in Juveniles and Adolescents. Angle Orthod 1986; 56: 7-30.
- Holdaway RA. A Soft Tissue Cephalometric Analysis and its Use in Orthodontic Treatment Planning. Part I. Am J Orthod 1983; 84: 1-28.
- Hopkin GB, Houston WJB, James GA. The Cranial Base as an Aetiological Factor in Malocclusion. Angle Orthod 1968; 38: 250-5.
- Houston WJB. The Analysis of Errors in Orthodontic Measurements. Am J Orthod 1983; 83: 382-90.
- Hwang HS, Kim WS, McNamara JA Jr. A Comparative Study of Two Methods of Quantifying the Soft Tissue Profile. Angle Orthod 2000; 70: 200-7.
- Isaacson JR, Isaacson RJ, Speidel TM, Waring RW. Extreme Variations in Vertical Facial Growth and Associated Variation in Skeletal and Dental Relationship. Angle Orthod 1971; 41: 219-29.
- Jacobson A, Evans WG, Preston CB, Sadowsky PL. Mandibular Prognathism. Am J Orthod 1974; 66: 140-71.
- Jacobson A. The "Wits" Appraisal of Jaw Disharmony. Am J Orthod 1975; 67: 125-38.
- Jarabak JR, Fizzell JA. Technique and Treatment with Lightwire Edgewise Appliance. 2<sup>nd</sup> ed. St.Louis: Mosby, 1972.

- Jarvinen S. Relation of the Wits Appraisal to the ANB Angle: A Statistical Appraisal. Am J Orthod Dentofac Orthop 1988; 94: 432-5.
- Jotikasthira D. Variation in Dentofacial Morphology in Thai and Norwegian Adults with Ideal Occlusion [Thesis]. Bergen: University of Bergen, 1989.
- Kahl-Nieke B, Schwarze CW. Treatment and Postretention Changes in Dental Arch Width Dimensions- a Long-Term Evaluation of Influencing Cofactors. Am J Orthod Dentofac Orthop 1996; 109: 368-78.
- Kerr WJS, TenHave TR. Mandibular Position in Class III Malocclusion. Br J Orthod 1988; 15: 241-5.
- Kilicoglu H, Kirlic Y. Profile Changes in Patients with Class III Malocclusions after Delaire Mask Therapy. Am J Orthod Dentofac Orthop 1998; 113: 453-62.
- Lee RT. Arch Width and Form: A Review. Am J Orthod Dentofac Orthop 1999; 115: 305-13.
- Lew KK, Fong WC. Horizontal Skeletal Typing in an Ethnic Chinese Population with True Class III Malocclusion. Br J Orthod 1993; 20: 19-23.
- Little RM, Reidal RA, Artun J. An Evaluation in Mandibular Anterior Alignment from 10-20 Years Postretention. Am J Orthod Dentofac Orthop 1988; 93: 423-8.
- Little RM, Wallen TR, Riedel RA. Stability and Retention of Mandibular Anterior Alignment - Four Premolar Extraction Cases Treated by Traditional Edgewise Orthodontics. Am J Orthod 1981; 80: 349-65.
- McClintock Robison J, Rinchuse DJ, Zullo TG. Relationship of Skeletal Pattern and Nasal Form. Am J Orthod 1986; 89: 499-506.
- Merz ML, Isaacson RJ, Germane N, Rubenstein LK. Tooth Diameters and Arch Perimeters in a Black and a White Population. Am J Orthod Dentofac Orthop 1991; 100: 53-8.
- Nabangxang H. Height, Head Circumference and Weight of Chiang Mai Children Age 0-18 Years. Chiang Mai: The Department of Preventive and Social Medicine, The Faculty of Medicine, Chiang Mai University, 1978.

- Nojima K, McLaughlin RP, Isshiki Y, Sinclair PM. A Comparative Study of Caucasian and Japanese Mandibular Clinical Arch Forms. *Angle Orthod* 2001; 71: 195-200.
- Nummikoski P, Langlais RP, Welander U. Dental and Mandibular Arch Widths in Three Ethnic Groups in Texas: A Radiographic Study. *Oral Surg Oral Med Oral Pathol* 1988; 65: 609-17.
- Patanaporn V. Cephalometric Study of Northern Thais after Maturity according to Ricketts Analysis. *CM Dent J* 1996; 17: 68-81.
- Proffit WR, Fields HW Jr. Malocclusion and Dentofacial Deformity in Contemporary Society. In: Rudolph P, editor. *Contemporary Orthodontics*. 3<sup>rd</sup> ed. St.Louis: Mosby, 1999: 9-13.
- Raberin M, Brunner F. Dimensions and Form of Dental Arches in Subjects with Normal Occlusions. *Am J Orthod Dentofac Orthop* 1993; 104: 67-72.
- Rakosi T, Jonas I, Gaber TM. *Color Atlas of Dental Medicine: Orthodontic-Diagnosis*. New York: Georg Thieme Verlag, 1993: 207-18.
- Ridell A, Soremark R, Lundberg M. Roentgen-Cephalometric Analysis of the Jaws in Subjects with and without Mandibular Protrusion. *Acta Odont Scand* 1971; 29: 103-21.
- Scheideman GB, Bell WH, Legan HL, Finn RA, Reisch JS. Cephalometric Analysis of Dentofacial Normals. *Am J Orthod* 1980; 78: 404-20.
- Sorathesn K. Craniofacial Norm for Thai in Combined Orthodontic Surgical Procedure. *J Dent Assoc Thai* 1988; 38: 190-201.
- Steiner CC. Cephalometric for You and Me. *Am J Orthod* 1953; 39: 729-55.
- Steiner CC. Cephalometric in Clinical Practice. *Angle Orthod* 1959; 29: 8-29.
- Steiner CC. The Use of Cephalometric as an Aid to Planning and Assessing Orthodontic Treatment. *Am J Orthod* 1960; 46: 721-35.
- Suchato W, Chaiwat J. Cephalometric Evaluation of the Dentofacial Complex of Thai Adults. *J Dent Assoc Thai* 1984; 34: 233-43.

- Suchato W, Chaiwat J. Prevalency of Malocclusion in Young Adult. Mahidol Dent J 1981; 2: 133-43.
- Swasdi-ampairaks S. Lateral Cephalometric Analysis of Anterior Open Bite and Normal Subjects in Adults [Thesis]. Chiang Mai: Chiang Mai University, 1997.
- Tahmina K, Tanaka E, Tanne K. Craniofacial Morphology in Orthodontically Treated Patients of Class III Malocclusion with Stable and Unstable Treatment Outcomes. Am J Orthod Dentofac Orthop 2000; 117: 681-90.
- Toms AP. Class III Malocclusion: A Cephalometric Study of Saudi Arabians. Br J Orthod 1989; 16: 201-6.
- Triwattanawong W. Craniofacial Values of Anterior Deepbite Samples in Northern Thai Adults [Thesis]. Chiang Mai: Chiang Mai University, 2001.
- Zar JH. Biostatistical Analysis. 3<sup>rd</sup> ed. NJ: Prentic-Hall, 1996: 133.

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่  
Copyright © by Chiang Mai University  
All rights reserved