

TABLE OF CONTENTS


	Page
ACKNOWLEDGEMENTS	iii
ABSTRACT (THAI)	v
ABSTRACT (ENGLISH)	vii
LIST OF TABLES	xi
LIST OF ILLUSTRATIONS	xii
CHAPTER I INTRODUCTION	
1.1 Statement of the problem and objectives	1
1.2 Anticipated benefits	3
CHAPTER II LITERATURE REVIEW	
2.1 Open bite	4
2.2 Orthodontic molar intrusion	5
2.3 Clinical application of miniscrew implants for molar intrusion	7
2.4 Assessments of orthodontic molar intrusion	8

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright © by Chiang Mai University
All rights reserved

	Page
2.5 Assessments of miniscrew implant stability	12
2.6 Gingival crevicular fluid (GCF), Peri-miniscrew implant crevicular fluid (PMICF) and Enzyme-linked immunosorbent assay (ELISA)	16
CHAPTER III MATERIALS AND METHODS	
3.1 Materials	21
3.2 Methods	23
CHAPTER IV RESULTS	31
CHAPTER V DISCUSSION	43
CHAPTER VI CONCLUSIONS	48
BIBLIOGRAPHY	50
APPENDIX	59
CIRRICULUM VITAE	73

LIST OF TABLES

TABLE		Page
4.1	Age distribution and gender of the subjects in this study	31



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved

LIST OF ILLUSTRATIONS

Figure	Page
2.1 ELISA method	19
3.1 Orthodontic appliances used in the study	23
3.2 Diagram of the experimental design	24
3.3 Gingival crevicular fluid and peri-miniscrew implant crevicular fluid collection	25
3.4 Clinical mobility assessment of miniscrew implants	25
3.5 Miniscrew implant placement procedure	26
3.6 Procedures for connecting two Sentalloy closed coil springs to miniscrew implant head	28
4.1 Boxplot graphs of medians of CS (WF6 epitope) levels during the unloaded (2 weeks) and the loaded (12 weeks) periods of all groups	33
4.2 Boxplot graphs of medians of CS (WF6 epitope) levels during the unloaded period (2 weeks) and each one-week interval of the loaded period (12 weeks) around experimental molars	34

Figure	Page
4.3 Boxplot graphs of medians of CS (WF6 epitope) levels during the unloaded period (2 weeks) and each one-week interval of the loaded period (12 weeks) around control right mandibular first molars	35
4.4 Boxplot graphs of medians of CS (WF6 epitope) levels during the unloaded period (2 weeks) and each one-week interval of the loaded period (12 weeks) around control right maxillary second molars	36
4.5 Boxplot graphs of medians of CS (WF6 epitope) levels during the unloaded period (2 weeks) and each one-week interval of the loaded period (12 weeks) around miniscrew implants	37
4.6 Boxplot graphs of medians of CS (WF6 epitope) levels during the unloaded period (2 weeks) and each two-week interval of the loaded period (12 weeks) around experimental molars	38
4.7 Boxplot graphs of medians of CS (WF6 epitope) levels during the unloaded period (2 weeks) and each two-week interval of the loaded period (12 weeks) around control right mandibular first molars	39

Figure	Page
4.8 Boxplot graphs of medians of CS (WF6 epitope) levels during the unloaded period (2 weeks) and each two-week interval of the loaded period (12 weeks) around control right maxillary second molars	40
4.9 Boxplot graphs of medians of CS (WF6 epitope) levels during the unloaded period (2 weeks) and each two-week interval of the loaded period (12 weeks) around miniscrew implants	41