

BIBLIOGRAPHY

- Akimenko MA, Johnson SL, Westerfield M, Ekker M (1995). Differential induction of four *msx* homeobox genes during fin development and regeneration in zebrafish. *Development* 121(2):347-57.
- Alappat S, Zhang ZY, Chen YP (2003). *Msx* homeobox gene family and craniofacial development. *Cell Res* 13(6):429-42.
- Andersen H, Matthiessen M (1967). Histochemistry of the early development of the human central face and nasal cavity with special reference to the movements and fusion of the palatine processes. *Acta Anat (Basel)* 68(4):473-508.
- Andersson E, Tryggvason U, Deng Q, Friling S, Alekseenko Z, Robert B, et al. (2006). Identification of intrinsic determinants of midbrain dopamine neurons. *Cell* 124(2):393-405.
- Arte S, Nieminen P, Apajalahti S, Haavikko K, Thesleff I, Pirinen S (2001). Characteristics of incisor-premolar hypodontia in families. *J Dent Res* 80(5):1445-50.
- Beatty TH, Wang H, Hetmanski JB, Fan YT, Zeiger JS, Liang KY, et al. (2001). A case-control study of nonsyndromic oral clefts in Maryland. *Ann Epidemiol* 11(6):434-42.
- Bei M, Maas R (1998). FGFs and BMP4 induce both *Msx1*-independent and *Msx1*-dependent signaling pathways in early tooth development. *Development* 125(21):4325-33.

- Bei M, Kratochwil K, Maas RL (2000). BMP4 rescues a non-cell-autonomous function of Msx1 in tooth development. *Development* 127(21):4711-8.
- Bendall AJ, Rincon-Limas DE, Botas J, Abate-Shen C (1998). Protein complex formation between Msx1 and Lhx2 homeoproteins is incompatible with DNA binding activity. *Differentiation* 63(3):151-7.
- Bendall AJ, Ding J, Hu G, Shen MM, Abate-Shen C (1999). Msx1 antagonizes the myogenic activity of Pax3 in migrating limb muscle precursors. *Development* 126(22):4965-76.
- Blin-Wakkach C, Lezot F, Ghoul-Mazgar S, Hotton D, Monteiro S, Teillaud C, et al. (2001). Endogenous Msx1 antisense transcript: in vivo and in vitro evidences, structure, and potential involvement in skeleton development in mammals. *Proc Natl Acad Sci U S A* 98(13):7336-41.
- Bonano M, Tribulo C, De Calisto J, Marchant L, Sanchez SS, Mayor R, et al. (2008). A new role for the Endothelin-1/Endothelin-A receptor signaling during early neural crest specification. *Dev Biol* 323(1):114-29.
- Cam Y, Neumann MR, Oliver L, Raulais D, Janet T, Ruch JV (1992). Immunolocalization of acidic and basic fibroblast growth factors during mouse odontogenesis. *Int J Dev Biol* 36(3):381-9.
- Campbell P, Neil T, Wake PN (1989). Horner's syndrome caused by an intercostal chest drain. *Thorax* 44(4):305-6.
- Carlson BM. *Human embryology and developmental biology*, 2004.
- Catron KM, Iler N, Abate C (1993). Nucleotides flanking a conserved TAAT core dictate the DNA binding specificity of three murine homeodomain proteins. *Mol Cell Biol* 13(4):2354-65.

- Catron KM, Zhang H, Marshall SC, Inostroza JA, Wilson JM, Abate C (1995). Transcriptional repression by Msx-1 does not require homeodomain DNA-binding sites. *Mol Cell Biol* 15(2):861-71.
- Catron KM, Wang H, Hu G, Shen MM, Abate-Shen C (1996). Comparison of MSX-1 and MSX-2 suggests a molecular basis for functional redundancy. *Mech Dev* 55(2):185-99.
- Charite J, McFadden DG, Merlo G, Levi G, Clouthier DE, Yanagisawa M, et al. (2001). Role of Dlx6 in regulation of an endothelin-1-dependent, dHAND branchial arch enhancer. *Genes Dev* 15(22):3039-49.
- Chen Y, Bei M, Woo I, Satokata I, Maas R (1996). Msx1 controls inductive signaling in mammalian tooth morphogenesis. *Development* 122(10):3035-44.
- Chen Y, Zhao X (1998). Shaping limbs by apoptosis. *J Exp Zool* 282(6):691-702.
- Chishti MS, Muhammad D, Haider M, Ahmad W (2006). A novel missense mutation in MSX1 underlies autosomal recessive oligodontia with associated dental anomalies in Pakistani families. *J Hum Genet* 51(10):872-8.
- Chitty LS, Dennis N, Baraitser M (1996). Hidrotic ectodermal dysplasia of hair, teeth, and nails: case reports and review. *J Med Genet* 33(8):707-10.
- Clouthier DE, Hosoda K, Richardson JA, Williams SC, Yanagisawa H, Kuwaki T, et al. (1998). Cranial and cardiac neural crest defects in endothelin-A receptor-deficient mice. *Development* 125(5):813-24.
- Cobourne MT (1999). The genetic control of early odontogenesis. *Br J Orthod* 26(1):21-8.
- Coelho CN, Upholt WB, Kosher RA (1993). The expression pattern of the chicken homeobox-containing gene GHox-7 in developing polydactylous limb buds

suggests its involvement in apical ectodermal ridge-directed outgrowth of limb mesoderm and in programmed cell death. *Differentiation* 52(2):129-37.

Das P, Stockton DW, Bauer C, Shaffer LG, D'Souza RN, Wright T, et al. (2002). Haploinsufficiency of PAX9 is associated with autosomal dominant hypodontia. *Hum Genet* 110(4):371-6.

de Jong DM, Hislop NR, Hayward DC, Reece-Hoyes JS, Pontynen PC, Ball EE, et al. (2006). Components of both major axial patterning systems of the Bilateria are differentially expressed along the primary axis of a 'radiate' animal, the anthozoan cnidarian *Acropora millepora*. *Dev Biol* 298(2):632-43.

De Muynck S, Schollen E, Matthijs G, Verdonck A, Devriendt K, Carels C (2004). A novel MSX1 mutation in hypodontia. *Am J Med Genet A* 128A(4):401-3..

Ekker M, Akimenko MA, Allende ML, Smith R, Drouin G, Langille RM, et al. (1997). Relationships among msx gene structure and function in zebrafish and other vertebrates. *Mol Biol Evol* 14(10):1008-22

Ferguson CA, Tucker AS, Sharpe PT (2000). Temporospatial cell interactions regulating mandibular and maxillary arch patterning. *Development* 127(2):403-12.

Finnerty JR, Martindale MQ (1997). Homeoboxes in sea anemones (Cnidaria:Anthozoa): a PCR-based survey of *Nematostella vectensis* and *Metridium senile*. *Biol Bull* 193(1):62-76.

Finnerty JR, Mazza ME, Jezewski PA (2009). Domain duplication, divergence, and loss events in vertebrate Msx paralogs reveal phylogenomically informed disease markers. *BMC Evol Biol* 9(18):1-23.

- Galle S, Yanze N, Seipel K (2005). The homeobox gene *Msx* in development and transdifferentiation of jellyfish striated muscle. *Int J Dev Biol* 49(8):961-7.
- Gene Expression in Tooth.
<http://bite-it.helsinki.fi/> (12 September 2009)
- Gorlin RJ, Cohen NM, Hannekam R. syndromes of the had and neck. Oxford University Press, New York 2001.
- Grahnén H (1956). Hypodontia in the permanent dentition: a clinical and genetical investigation. *Odont Revy* 7(Suppl 3):1-100.
- Gritli-Linde A (2007). Molecular control of secondary palate development. *Dev Biol* 301(2):309-26.
- Gupta V, Bei M (2006). Modification of *Msx1* by SUMO-1. *Biochem Biophys Res Commun* 345(1):74-7.
- Hardcastle Z, Mo R, Hui CC, Sharpe PT (1998). The *Shh* signalling pathway in tooth development: defects in *Gli2* and *Gli3* mutants. *Development* 125(15):2803-11.
- Hay RT (2005). SUMO: a history of modification. *Mol Cell* 18(1):1-12.
- Hewitt JE, Clark LN, Ivens A, Williamson R (1991). Structure and sequence of the human homeobox gene *HOX7*. *Genomics* 11(3):670-8.
- Hill RE, Jones PF, Rees AR, Sime CM, Justice MJ, Copeland NG, et al. (1989). A new family of mouse homeo box-containing genes: molecular structure, chromosomal location, and developmental expression of *Hox-7.1*. *Genes Dev* 3(1):26-37.
- Hodges SJ, Harley KE (1999). Witkop tooth and nail syndrome: report of two cases in a family. *Int J Paediatr Dent* 9(3):207-11.

- Hovde S, Abate-Shen C, Geiger JH (2001). Crystal structure of the Msx-1 homeodomain/DNA complex. *Biochemistry* 40(40):12013-21.
- Hu G, Lee H, Price SM, Shen MM, Abate-Shen C (2001). Msx homeobox genes inhibit differentiation through upregulation of cyclin D1. *Development* 128(12):2373-84.
- Hu G, Vastardis H, Bendall AJ, Wang Z, Logan M, Zhang H, et al. (1998). Haploinsufficiency of MSX1: a mechanism for selective tooth agenesis. *Mol Cell Biol* 18(10):6044-51.
- Hwang SJ, Beaty TH, McIntosh I, Hefferon T, Panny SR (1998). Association between homeobox-containing gene MSX1 and the occurrence of limb deficiency. *Am J Med Genet* 75(4):419-23.
- Ivey K, Tyson B, Ukidwe P, McFadden DG, Levi G, Olson EN, et al. (2003). Galphaq and Galpha11 proteins mediate endothelin-1 signaling in neural crest-derived pharyngeal arch mesenchyme. *Dev Biol* 255(2):230-7.
- Jernvall J, Thesleff I (2000). Reiterative signaling and patterning during mammalian tooth morphogenesis. *Mech Dev* 92(1):19-29.
- Jezewski PA, Vieira AR, Nishimura C, Ludwig B, Johnson M, O'Brien SE, et al. (2003). Complete sequencing shows a role for MSX1 in non-syndromic cleft lip and palate. *J Med Genet* 40(6):399-407.
- Johnson ES, Gupta AA (2001). An E3-like factor that promotes SUMO conjugation to the yeast septins. *Cell* 106(6):735-44.
- Jowett AK, Vainio S, Ferguson MW, Sharpe PT, Thesleff I (1993). Epithelial mesenchymal interactions are required for msx 1 and msx 2 gene expression in the developing murine molar tooth. *Development* 117(2):461-70.

- Jugessur A, Lie RT, Wilcox AJ, Murray JC, Taylor JA, Saugstad OD, et al. (2003). Variants of developmental genes (TGFA, TGFB3, and MSX1) and their associations with orofacial clefts: a case-parent triad analysis. *Genet Epidemiol* 24(3):230-9.
- Jugessur A, Murray JC (2005). Orofacial clefting: recent insights into a complex trait. *Curr Opin Genet Dev* 15(3):270-8.
- Jumlongras D, Bei M, Stimson JM, Wang WF, DePalma SR, Seidman CE, et al. (2001). A nonsense mutation in MSX1 causes Witkop syndrome. *Am J Hum Genet* 69(1):67-74.
- Kettunen P, Thesleff I (1998). Expression and function of FGFs-4, -8, and -9 suggest functional redundancy and repetitive use as epithelial signals during tooth morphogenesis. *Dev Dyn* 211(3):256-68.
- Kim JW, Simmer JP, Lin BP, Hu JC (2006). Novel MSX1 frameshift causes autosomal-dominant oligodontia. *J Dent Res* 85(3):267-71.
- Koillinen H, Lahermo P, Rautio J, Hukki J, Peyrard-Janvid M, Kere J (2005). A genome-wide scan of non-syndromic cleft palate only (CPO) in Finnish multiplex families. *J Med Genet* 42(2):177-84.
- Kolenc-Fuse FJ (2004). Tooth agenesis: in search of mutations behind failed dental development. *Med Oral Patol Oral Cir Bucal* 9(5):390-5; 385-90.
- Lan Y, Ovitt CE, Cho ES, Maltby KM, Wang Q, Jiang R (2004). Odd-skipped related 2 (Osr2) encodes a key intrinsic regulator of secondary palate growth and morphogenesis. *Development* 131(13):3207-16.

- Lee H, Habas R, Abate-Shen C (2004). MSX1 cooperates with histone H1b for inhibition of transcription and myogenesis. *Science* 304(5677):1675-8.
- Lee H, Quinn JC, Prasanth KV, Swiss VA, Economides KD, Camacho MM, et al. (2006). PIAS1 confers DNA-binding specificity on the Msx1 homeoprotein. *Genes Dev* 20(7):784-94.
- Lee SM, Danielian PS, Fritsch B, McMahon AP (1997). Evidence that FGF8 signalling from the midbrain-hindbrain junction regulates growth and polarity in the developing midbrain. *Development* 124(5):959-69.
- Lidral AC, Moreno LM (2005). Progress toward discerning the genetics of cleft lip. *Curr Opin Pediatr* 17(6):731-9.
- Lidral AC, Romitti PA, Basart AM, Doetschman T, Leysens NJ, Daack-Hirsch S, et al. (1998). Association of MSX1 and TGFB3 with nonsyndromic clefting in humans. *Am J Hum Genet* 63(2):557-68.
- Lidral AC, Reising BC (2002). The role of MSX1 in human tooth agenesis. *J Dent Res* 81(4):274-8.
- Ma L, Golden S, Wu L, Maxson R (1996). The molecular basis of Boston-type craniosynostosis: the Pro148-->His mutation in the N-terminal arm of the MSX2 homeodomain stabilizes DNA binding without altering nucleotide sequence preferences. *Hum Mol Genet* 5(12):1915-20.
- Ma L, Liu J, Wu T, Plikus M, Jiang TX, Bi Q, et al. (2003). 'Cyclic alopecia' in Msx2 mutants: defects in hair cycling and hair shaft differentiation. *Development* 130(2):379-89.
- Mackenzie A, Leeming GL, Jowett AK, Ferguson MW, Sharpe PT (1991). The homeobox gene Hox 7.1 has specific regional and temporal expression patterns

during early murine craniofacial embryogenesis, especially tooth development in vivo and in vitro. *Development* 111(2):269-85.

Mackenzie A, Ferguson MW, Sharpe PT (1992). Expression patterns of the homeobox gene, Hox-8, in the mouse embryo suggest a role in specifying tooth initiation and shape. *Development* 115(2):403-20.

Morgan R, In der Rieden P, Hooiveld MH, Durston AJ (2000). Identifying HOX paralog groups by the PBX-binding region. *Trends Genet* 16(2):66-7.

Mostowska A, Kobiela A, Trzeciak WH (2003). Molecular basis of non-syndromic tooth agenesis: mutations of MSX1 and PAX9 reflect their role in patterning human dentition. *Eur J Oral Sci* 111(5):365-70.

Mostowska A, Biedziak B, Trzeciak WH (2006). A novel c.581C>T transition localized in a highly conserved homeobox sequence of MSX1: is it responsible for oligodontia? *J Appl Genet* 47(2):159-64.

Neubuser A, Peters H, Balling R, Martin GR (1997). Antagonistic interactions between FGF and BMP signaling pathways: a mechanism for positioning the sites of tooth formation. *Cell* 90(2):247-55.

Nieminen P, Kotilainen J, Aalto Y, Knuutila S, Pirinen S, Thesleff I (2003). *MSX1* gene is deleted in Wolf-Hirschhorn syndrome patients with oligodontia. *J Dent Res* 82(12):1013-7.

Nugent P, Greene RM (1998). *MSX-1* gene expression and regulation in embryonic palatal tissue. *In Vitro Cell Dev Biol Anim* 34(10):831-5.

Odelberg SJ, Kollhoff A, Keating MT (2000). Dedifferentiation of mammalian myotubes induced by msx1. *Cell* 103(7):1099-109.

- Ogawa T, Kapadia H, Feng JQ, Raghow R, Peters H, D'Souza RN (2006). Functional consequences of interactions between Pax9 and Msx1 genes in normal and abnormal tooth development. *J Biol Chem* 281(27):18363-9.
- Padanilam BJ, Stadler HS, Mills KA, McLeod LB, Solursh M, Lee B, et al. (1992). Characterization of the human HOX 7 cDNA and identification of polymorphic markers. *Hum Mol Genet* 1(6):407-10.
- Pauws E, Stanier P (2007). FGF signalling and SUMO modification: new players in the aetiology of cleft lip and/or palate. *Trends Genet* 23(12):631-40.
- Pawlowska E, Janik-Papis K, Wisniewska-Jarosinska M, Szczepanska J, Blasiak J (2009). Mutations in the human homeobox MSX1 gene in the congenital lack of permanent teeth. *Tohoku J Exp Med* 217(4):307-12.
- Peltenburg LT, Murre C (1996). Engrailed and Hox homeodomain proteins contain a related Pbx interaction motif that recognizes a common structure present in Pbx. *EMBO J* 15(13):3385-93.
- Peters H, Neubuser A, Kratochwil K, Balling R (1998). Pax9-deficient mice lack pharyngeal pouch derivatives and teeth and exhibit craniofacial and limb abnormalities. *Genes Dev* 12(17):2735-47.
- Peters H, Balling R (1999). Teeth. Where and how to make them. *Trends Genet* 15(2):59-65.
- Rave-Harel N, Miller NL, Givens ML, Mellon PL (2005). The Groucho-related gene family regulates the gonadotropin-releasing hormone gene through interaction with the homeodomain proteins MSX1 and OCT1. *J Biol Chem* 280(35):30975-83.

- Rice R, Spencer-Dene B, Connor EC, Gritli-Linde A, McMahon AP, Dickson C, et al. (2004). Disruption of Fgf10/Fgfr2b-coordinated epithelial-mesenchymal interactions causes cleft palate. *J Clin Invest* 113(12):1692-700.
- Richelle-Maurer E, Boury-Esnault N, Itskovich VB, Manuel M, Pomponi SA, Van de Vyver G, et al. (2006). Conservation and phylogeny of a novel family of non-Hox genes of the Antp class in Demospongiae (porifera). *J Mol Evol* 63(2):222-30.
- Richman JM, Herbert M, Matovinovic E, Walin J (1997). Effect of fibroblast growth factors on outgrowth of facial mesenchyme. *Dev Biol* 189(1):135-47.
- Robert B, Sassoon D, Jacq B, Gehring W, Buckingham M (1989). Hox-7, a mouse homeobox gene with a novel pattern of expression during embryogenesis. *EMBO J* 8(1):91-100.
- Ryan JF, Burton PM, Mazza ME, Kwong GK, Mullikin JC, Finnerty JR (2006). The cnidarian-bilaterian ancestor possessed at least 56 homeoboxes: evidence from the starlet sea anemone, *Nematostella vectensis*. *Genome Biol* 7(7):R64.
- Sarkar L, Cobourne M, Naylor S, Smalley M, Dale T, Sharpe PT (2000). Wnt/Shh interactions regulate ectodermal boundary formation during mammalian tooth development. *Proc Natl Acad Sci U S A* 97(9):4520-4.
- Satokata I, Maas R (1994). Msx1 deficient mice exhibit cleft palate and abnormalities of craniofacial and tooth development. *Nat Genet* 6(4):348-56.
- Satokata I, Ma L, Ohshima H, Bei M, Woo I, Nishizawa K, et al. (2000). Msx2 deficiency in mice causes pleiotropic defects in bone growth and ectodermal organ formation. *Nat Genet* 24(4):391-5.

- Schummer M, Scheurlen I, Schaller C, Galliot B (1992). HOM/HOX homeobox genes are present in hydra (*Chlorohydra viridissima*) and are differentially expressed during regeneration. *EMBO J* 11(5):1815-23.
- Seielstad M, Bekele E, Ibrahim M, Toure A, Traore M (1999). A view of modern human origins from Y chromosome microsatellite variation. *Genome Res* 9(6):558-67.
- Seimiya M, Ishiguro H, Miura K, Watanabe Y, Kurosawa Y (1994). Homeobox containing genes in the most primitive metazoa, the sponges. *Eur J Biochem* 221(1):219-25.
- Sharpe PT (1995). Homeobox genes and orofacial development. *Connect Tissue Res* 32(1-4):17-25.
- Shetty S, Takahashi T, Matsui H, Ayengar R, Raghov R (1999). Transcriptional autorepression of Msx1 gene is mediated by interactions of Msx1 protein with a multi-protein transcriptional complex containing TATA-binding protein, Sp1 and cAMP-response-element-binding protein-binding protein (CBP/p300). *Biochem J* 339 (Pt 3)(751-8.
- Slayton RL, Williams L, Murray JC, Wheeler JJ, Lidral AC, Nishimura CJ (2003). Genetic association studies of cleft lip and/or palate with hypodontia outside the cleft region. *Cleft Palate Craniofac J* 40(3):274-9.
- Smith ST, Jaynes JB (1996). A conserved region of engrailed, shared among all en-, gsc-, Nk1-, Nk2- and msh-class homeoproteins, mediates active transcriptional repression in vivo. *Development* 122(10):3141-50.

- South ST, Whitby H, Battaglia A, Carey JC, Brothman AR (2008). Comprehensive analysis of Wolf-Hirschhorn syndrome using array CGH indicates a high prevalence of translocations. *Eur J Hum Genet* 16(1):45-52.
- Stockton DW, Das P, Goldenberg M, D'Souza RN, Patel PI (2000). Mutation of PAX9 is associated with oligodontia. *Nat Genet* 24(1):18-9.
- Su MW, Suzuki HR, Solursh M, Ramirez F (1991). Progressively restricted expression of a new homeobox-containing gene during *Xenopus laevis* embryogenesis. *Development* 111(4):1179-87.
- Suazo J, Santos JL, Jara L, Blanco R (2010). Parent-of-origin effects for *MSX1* in a Chilean population with nonsyndromic cleft lip/palate. *Am J Med Genet A* 152A(8):2011-6.
- Suzuki Y, Jezewski PA, Machida J, Watanabe Y, Shi M, Cooper ME, et al. (2004). In a Vietnamese population, *MSX1* variants contribute to cleft lip and palate. *Genet Med* 6(3):117-25.
- Tabin CJ (1991). Retinoids, homeoboxes, and growth factors: toward molecular models for limb development. *Cell* 66(2):199-217.
- Takahashi H, Kamiya A, Ishiguro A, Suzuki AC, Saitou N, Toyoda A, et al. (2008). Conservation and diversification of *Msx* protein in metazoan evolution. *Mol Biol Evol* 25(1):69-82.
- Takahashi T, Guron C, Shetty S, Matsui H, Raghov R (1997). A minimal murine *Msx-1* gene promoter. Organization of its cis-regulatory motifs and their role in transcriptional activation in cells in culture and in transgenic mice. *J Biol Chem* 272(36):22667-78.

Takahashi Y, Le Douarin N (1990). cDNA cloning of a quail homeobox gene and its expression in neural crest-derived mesenchyme and lateral plate mesoderm. *Proc Natl Acad Sci U S A* 87(19):7482-6.

The amino acid repository, Image library of biological macromolecules.

<http://www.imb-jena.com> (9 September 2010)

Thesleff I (2003). Developmental biology and building a tooth. *Quintessence Int* 34(8):613-20.

Thomas T, Kurihara H, Yamagishi H, Kurihara Y, Yazaki Y, Olson EN, et al. (1998). A signaling cascade involving endothelin-1, dHAND and msx1 regulates development of neural-crest-derived branchial arch mesenchyme. *Development* 125(16):3005-14.

Tongkobpetch S, Siriwan P, Shotelersuk V (2006). MSX1 mutations contribute to nonsyndromic cleft lip in a Thai population. *J Hum Genet* 51(8):671-6.

Trasler DG (1968). Pathogenesis of cleft lip and its relation to embryonic face shape in A-J and C57BL mice. *Teratology* 1(1):33-49.

Trumpp A, Depew MJ, Rubenstein JL, Bishop JM, Martin GR (1999). Cre-mediated gene inactivation demonstrates that FGF8 is required for cell survival and patterning of the first branchial arch. *Genes Dev* 13(23):3136-48.

Tucker A, Sharpe P (2004). The cutting-edge of mammalian development; how the embryo makes teeth. *Nat Rev Genet* 5(7):499-508.

Tucker AS, Matthews KL, Sharpe PT (1998). Transformation of tooth type induced by inhibition of BMP signaling. *Science* 282(5391):1136-8.

Tucker AS, Sharpe PT (1999). Molecular genetics of tooth morphogenesis and patterning: the right shape in the right place. *J Dent Res* 78(4):826-34.


- Tucker AS, Yamada G, Grigoriou M, Pachnis V, Sharpe PT (1999). Fgf-8 determines rostral-caudal polarity in the first branchial arch. *Development* 126(1):51-61.
- Tupler R, Bortotto L, Buhler EM, Alkan M, Malik NJ, Bosch-Al Jadooa N, et al. (1992). Paternal origin of the de novo deleted chromosome 4 in Wolf-Hirschhorn syndrome. *J Med Genet* 29(1):53-5.
- Tureckova J, Sahlberg C, Aberg T, Ruch JV, Thesleff I, Peterkova R (1995). Comparison of expression of the *msx-1*, *msx-2*, *BMP-2* and *BMP-4* genes in the mouse upper diastemal and molar tooth primordia. *Int J Dev Biol* 39(3):459-68.
- van den Boogaard MJ, Dorland M, Beemer FA, van Amstel HK (2000). *MSX1* mutation is associated with orofacial clefting and tooth agenesis in humans. *Nat Genet* 24(4):342-3.
- van den Boogaard MJ, de Costa D, Krapels IP, Liu F, van Duijn C, Sinke RJ, et al. (2008). The *MSX1* allele 4 homozygous child exposed to smoking at periconception is most sensitive in developing nonsyndromic orofacial clefts. *Hum Genet* 124(5):525-34.
- Vastardis H, Karimbux N, Guthua SW, Seidman JG, Seidman CE (1996). A human *MSX1* homeodomain missense mutation causes selective tooth agenesis. *Nat Genet* 13(4):417-21.
- Vastardis H (2000). The genetics of human tooth agenesis: new discoveries for understanding dental anomalies. *Am J Orthod Dentofacial Orthop* 117(6):650-6.
- Vieira AR (2003). Oral clefts and syndromic forms of tooth agenesis as models for genetics of isolated tooth agenesis. *J Dent Res* 82(3):162-5.

- Vieira AR, Romitti PA, Orioli IM, Castilla EE (2003). Inheritance of cleft palate in South America: evidence for a major locus recessive. *Orthod Craniofac Res* 6(2):83-7.
- Vieira AR, Avila JR, Daack-Hirsch S, Dragan E, Felix TM, Rahimov F, et al. (2005). Medical sequencing of candidate genes for nonsyndromic cleft lip and palate. *PLoS Genet* 1(6):e64.
- Vogel A, Roberts-Clarke D, Niswander L (1995). Effect of FGF on gene expression in chick limb bud cells in vivo and in vitro. *Dev Biol* 171(2):507-20.
- Witkop C, editor (1965). *Genetic disease of the oral cavity*. New York: McGraw-Hill.
- Wright TJ, Ricke DO, Denison K, Abmayr S, Cotter PD, Hirschhorn K, et al. (1997). A transcript map of the newly defined 165 kb Wolf-Hirschhorn syndrome critical region. *Hum Mol Genet* 6(2):317-24.
- Xuan K, Jin F, Liu YL, Yuan LT, Wen LY, Yang FS, et al. (2008). Identification of a novel missense mutation of MSX1 gene in Chinese family with autosomal-dominant oligodontia. *Arch Oral Biol* 53(8):773-9.
- Yamashiro T, Tummers M, Thesleff I (2003). Expression of bone morphogenetic proteins and Msx genes during root formation. *J Dent Res* 82(3):172-6.
- Yanagisawa H, Yanagisawa M, Kapur RP, Richardson JA, Williams SC, Clouthier DE, et al. (1998). Dual genetic pathways of endothelin-mediated intercellular signaling revealed by targeted disruption of endothelin converting enzyme-1 gene. *Development* 125(5):825-36.
- Yanagisawa H, Clouthier DE, Richardson JA, Charite J, Olson EN (2003). Targeted deletion of a branchial arch-specific enhancer reveals a role of dHAND in craniofacial development. *Development* 130(6):1069-78.

- Yu L, Gu S, Alappat S, Song Y, Yan M, Zhang X, et al. (2005). Shox2-deficient mice exhibit a rare type of incomplete clefting of the secondary palate. *Development* 132(19):4397-406.
- Zhang H, Catron KM, Abate-Shen C (1996). A role for the Msx-1 homeodomain in transcriptional regulation: residues in the N-terminal arm mediate TATA binding protein interaction and transcriptional repression. *Proc Natl Acad Sci U S A* 93(5):1764-9.
- Zhang H, Hu G, Wang H, Sciavolino P, Iler N, Shen MM, et al. (1997). Heterodimerization of Msx and Dlx homeoproteins results in functional antagonism. *Mol Cell Biol* 17(5):2920-32.
- Zhang Y, Zhao X, Hu Y, St Amand T, Zhang M, Ramamurthy R, et al. (1999). Msx1 is required for the induction of Patched by Sonic hedgehog in the mammalian tooth germ. *Dev Dyn* 215(1):45-53.
- Zhang Y, Zhang Z, Zhao X, Yu X, Hu Y, Geronimo B, et al. (2000). A new function of BMP4: dual role for BMP4 in regulation of Sonic hedgehog expression in the mouse tooth germ. *Development* 127(7):1431-43.
- Zhang Z, Song Y, Zhao X, Zhang X, Fermin C, Chen Y (2002). Rescue of cleft palate in Msx1-deficient mice by transgenic Bmp4 reveals a network of BMP and Shh signaling in the regulation of mammalian palatogenesis. *Development* 129(17):4135-46.
- Zhang Z, Zhang X, Avniel WA, Song Y, Jones SM, Jones TA, et al. (2003). Malleal processus brevis is dispensable for normal hearing in mice. *Dev Dyn* 227(1):69-77.

Zhao X, Zhang Z, Song Y, Zhang X, Zhang Y, Hu Y, et al. (2000). Transgenically ectopic expression of Bmp4 to the Msx1 mutant dental mesenchyme restores downstream gene expression but represses Shh and Bmp2 in the enamel knot of wild type tooth germ. *Mech Dev* 99(1-2):29-38.

Zhou YL, Lei Y, Snead ML (2000). Functional antagonism between Msx2 and CCAAT/enhancer-binding protein alpha in regulating the mouse amelogenin gene expression is mediated by protein-protein interaction. *J Biol Chem* 275(37):29066-75.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved