

BIBLIOGRAPHY

- Adaimy L, Chouery E, Megarbane H, Mroueh S, Delague V, Nicolas E, et al. (2007). Mutation in WNT10A is associated with an autosomal recessive ectodermal dysplasia: the odonto-onycho-dermal dysplasia. *Am J Hum Genet* 81(4):821-8.
- Andl T, Reddy ST, Gaddapara T, Millar SE (2002). WNT signals are required for the initiation of hair follicle development. *Dev Cell* 2(5):643-53.
- Bal E, Baala L, Cluzeau C, El Kerch F, Ouldim K, Hadj-Rabia S, et al. (2007). Autosomal dominant anhidrotic ectodermal dysplasias at the EDARADD locus. *Hum Mutat* 28(7):703-9.
- Bayes M, Hartung AJ, Ezer S, Pispa J, Thesleff I, Srivastava AK, et al. (1998). The anhidrotic ectodermal dysplasia gene (EDA) undergoes alternative splicing and encodes ectodysplasin-A with deletion mutations in collagenous repeats. *Hum Mol Genet* 7(11):1661-9.
- Bohring A, Stamm T, Spaich C, Haase C, Spree K, Hehr U, et al. (2009). WNT10A mutations are a frequent cause of a broad spectrum of ectodermal dysplasias with sex-biased manifestation pattern in heterozygotes. *Am J Hum Genet* 85(1):97-105.
- Botchkarev VA, Fessing MY (2005). Edar signaling in the control of hair follicle development. *J Investig Dermatol Symp Proc* 10(3):247-51.
- Brockschmidt FF, Hillmer AM, Eigelshoven S, Hanneken S, Heilmann S, Barth S, et al. (2008). Fine mapping of the human AR/EDA2R locus in androgenetic alopecia. *Br J Dermatol* 162(4):899-903

- Brosh R, Sarig R, Natan EB, Molchadsky A, Madar S, Bornstein C, et al. (2010). p53-dependent transcriptional regulation of EDA2R and its involvement in chemotherapy-induced hair loss. *FEBS Lett* 584(11):2473-7.
- Callahan N, Modesto A, Meira R, Seymen F, Patir A, Vieira AR (2009). Axis inhibition protein 2 (AXIN2) polymorphisms and tooth agenesis. *Arch Oral Biol* 54(1):45-9.
- Cambiaghi S, Restano L, Paakkonen K, Caputo R, Kere J (2000). Clinical findings in mosaic carriers of hypohidrotic ectodermal dysplasia. *Arch Dermatol* 136(2):217-24.
- Chassaing N, Bourthoumieu S, Cossee M, Calvas P, Vincent MC (2006). Mutations in EDAR account for one-quarter of non-ED1-related hypohidrotic ectodermal dysplasia. *Hum Mutat* 27(3):255-9.
- Chassaing N, Cluzeau C, Bal E, Guigue P, Vincent MC, Viot G, et al. (2010). Mutations in EDARADD account for a small proportion of hypohidrotic ectodermal dysplasia cases. *Br J Dermatol* 162(5):1044-8.
- Chen Y, Molloy SS, Thomas L, Gambee J, Bachinger HP, Ferguson B, et al. (2001). Mutations within a furin consensus sequence block proteolytic release of ectodysplasin-A and cause X-linked hypohidrotic ectodermal dysplasia. *Proc Natl Acad Sci U S A* 98(13):7218-23.
- Chu EY, Hens J, Andl T, Kairo A, Yamaguchi TP, Brisken C, et al. (2004). Canonical WNT signaling promotes mammary placode development and is essential for initiation of mammary gland morphogenesis. *Development* 131(19):4819-29.
- Courtney JM, Blackburn J, Sharpe PT (2005). The Ectodysplasin and NFkappaB signalling pathways in odontogenesis. *Arch Oral Biol* 50(2):159-63.

- Crawford PJ, Aldred MJ, Clarke A (1991). Clinical and radiographic dental findings in X linked hypohidrotic ectodermal dysplasia. *J Med Genet* 28(3):181-5.
- Cui CY, Hashimoto T, Grivennikov SI, Piao Y, Nedospasov SA, Schlessinger D (2006). Ectodysplasin regulates the lymphotoxin-beta pathway for hair differentiation. *Proc Natl Acad Sci U S A* 103(24):9142-7.
- Darwin C (1875). The variation of animals and plants under domestication, 2nd ed. London: John Murray. Vol. II, p. 319.
- Doffinger R, Smahi A, Bessia C, Geissmann F, Feinberg J, Durandy A, et al. (2001). X-linked anhidrotic ectodermal dysplasia with immunodeficiency is caused by impaired NF-kappaB signaling. *Nat Genet* 27(3):277-85.
- Durmowicz MC, Cui CY, Schlessinger D (2002). The EDA gene is a target of, but does not regulate Wnt signaling. *Gene* 285(1-2):203-11.
- Ezer S, Bayes M, Elomaa O, Schlessinger D, Kere J (1999). Ectodysplasin is a collagenous trimeric type II membrane protein with a tumor necrosis factor-like domain and co-localizes with cytoskeletal structures at lateral and apical surfaces of cells. *Hum Mol Genet* 8(11):2079-86.
- Elomaa O, Pulkkinen K, Hannelius U, Mikkola M, Saarialho-Kere U, Kere J (2001). Ectodysplasin is released by proteolytic shedding and binds to the EDAR protein. *Hum Mol Genet* 10(9):953-62.
- Fan H, Ye X, Shi L, Yin W, Hua B, Song G, et al. (2008). Mutations in the EDA gene are responsible for X-linked hypohidrotic ectodermal dysplasia and hypodontia in Chinese kindreds. *Eur J Oral Sci* 116(5):412-7.

- Ferguson BM, Brockdorff N, Formstone E, Ngyuen T, Kronmiller JE, Zonana J (1997). Cloning of Tabby, the murine homolog of the human EDA gene: evidence for a membrane-associated protein with a short collagenous domain. *Hum Mol Genet* 6(9):1589-94.
- Fliniaux I, Mikkola ML, Lefebvre S, Thesleff I (2008). Identification of dkk4 as a target of Eda-A1/Edar pathway reveals an unexpected role of ectodysplasin as inhibitor of Wnt signalling in ectodermal placodes. *Dev Biol* 320(1):60-71.
- Han D, Gong Y, Wu H, Zhang X, Yan M, Wang X, et al. (2008). Novel EDA mutation resulting in X-linked non-syndromic hypodontia and the pattern of EDA-associated isolated tooth agenesis. *Eur J Med Genet* 51(6):536-46.
- Hardy MH (1992). The secret life of the hair follicle. *Trends Genet* 8(2):55-61.
- Hashiguchi T, Yotsumoto S, Kanzaki T (2003). Mutations in the ED1 gene in Japanese families with X-linked hypohidrotic ectodermal dysplasia. *Exp Dermatol* 12(4):518-22.
- Headon DJ, Overbeek PA (1999). Involvement of a novel Tnf receptor homologue in hair follicle induction. *Nat Genet* 22(4):370-4.
- Headon DJ, Emmal SA, Ferguson BM, Tucker AS, Justice MJ, Sharpe PT, et al. (2001). Gene defect in ectodermal dysplasia implicates a death domain adapter in development. *Nature* 414(6866):913-6.
- Hens JR, Wysolmerski JJ (2005). Key stages of mammary gland development: molecular mechanisms involved in the formation of the embryonic mammary gland. *Breast Cancer Res* 7(5):220-4.

- Hofmann K (1999). The modular nature of apoptotic signaling proteins. *Cell Mol Life Sci* 55(8-9):1113-28.
- Israel A (2000). The IKK complex: an integrator of all signals that activate NF-kappaB? *Trends Cell Biol* 10(4):129-33.
- Itin PH, Fistarol SK (2004). Ectodermal dysplasias. *Am J Med Genet C Semin Med Genet* 131C(1):45-51.
- Jernvall J, Thesleff I (2000). Reiterative signaling and patterning during mammalian tooth morphogenesis. *Mech Dev* 92(1):19-29.
- Kangas AT, Evans AR, Thesleff I, Jernvall J (2004). Nonindependence of mammalian dental characters. *Nature* 432(7014):211-4.
- Kantaputra P, Sripathomsawat W (2010). WNT10A and isolated hypodontia. *Am J Med Genet A* 155A(5):1119-22.
- Kapadia H, Frazier-Bowers S, Ogawa T, D'Souza RN (2006). Molecular characterization of a novel PAX9 missense mutation causing posterior tooth agenesis. *Eur J Hum Genet* 14(4):403-9.
- Kere J, Srivastava AK, Montonen O, Zonana J, Thomas N, Ferguson B, et al. (1996). X-linked anhidrotic (hypohidrotic) ectodermal dysplasia is caused by mutation in a novel transmembrane protein. *Nat Genet* 13(4):409-16.
- Klein ML, Nieminen P, Lammi L, Niebuhr E, Kreiborg S (2005). Novel mutation of the initiation codon of PAX9 causes oligodontia. *J Dent Res* 84(1):43-7.
- Kobielsk K, Kobielsk A, Limon J, Trzeciak WH (1998). Mutation in the regulatory region of the EDA gene coincides with the symptoms of anhidrotic ectodermal dysplasia. *Acta Biochim Pol* 45(1):245-50.

- Kobielsk K, Kobielska A, Roszkiewicz J, Wierzba J, Limon J, Trzeciak WH (2001). Mutations in the EDA gene in three unrelated families reveal no apparent correlation between phenotype and genotype in the patients with an X-linked anhidrotic ectodermal dysplasia. *Am J Med Genet* 100(3):191-7.
- Kumar A, Eby MT, Sinha S, Jasmin A, Chaudhary PM (2001). The ectodermal dysplasia receptor activates the nuclear factor-kappaB, JNK, and cell death pathways and binds to ectodysplasin A. *J Biol Chem* 276(4):2668-77.
- Lamartine J (2003). Towards a new classification of ectodermal dysplasias. *Clin Exp Dermatol* 28(4):351-5.
- Laurikkala J, Mikkola M, Mustonen T, Aberg T, Koponen P, Pispa J, et al. (2001). TNF signaling via the ligand-receptor pair ectodysplasin and edar controls the function of epithelial signaling centers and is regulated by Wnt and activin during tooth organogenesis. *Dev Biol* 229(2):443-55.
- Laurikkala J, Pispa J, Jung HS, Nieminen P, Mikkola M, Wang X, et al. (2002). Regulation of hair follicle development by the TNF signal ectodysplasin and its receptor Edar. *Development* 129(10):2541-53.
- Li Q, Van Antwerp D, Mercurio F, Lee KF, Verma IM (1999). Severe liver degeneration in mice lacking the IkappaB kinase 2 gene. *Science* 284(5412):321-5.
- Li S, Li J, Cheng J, Zhou B, Tong X, Dong X, et al. (2008). Non-syndromic tooth agenesis in two Chinese families associated with novel missense mutations in the TNF domain of EDA (ectodysplasin A). *PLoS One* 3(6):e2396.

- Lexner MO, Bardow A, Hertz JM, Nielsen LA, Kreiborg S (2007). Anomalies of tooth formation in hypohidrotic ectodermal dysplasia. *Int J Paediatr Dent* 17(1):10-8.
- Lexner MO, Bardow A, Juncker I, Jensen LG, Almer L, Kreiborg S, et al. (2008). X-linked hypohidrotic ectodermal dysplasia. Genetic and dental findings in 67 Danish patients from 19 families. *Clin Genet* 74(3):252-9.
- Makris C, Godfrey VL, Krahn-Senftleben G, Takahashi T, Roberts JL, Schwarz T, et al. (2000). Female mice heterozygous for IKK gamma/NEMO deficiencies develop a dermatopathy similar to the human X-linked disorder incontinentia pigmenti. *Mol Cell* 5(6):969-79.
- Mikkola ML, Thesleff I (2003). Ectodysplasin signaling in development. *Cytokine Growth Factor Rev* 14(3-4):211-24.
- Mikkola ML (2009). Molecular aspects of hypohidrotic ectodermal dysplasia. *Am J Med Genet A* 149A(9):2031-6.
- Mikkola ML, Millar SE (2006). The mammary bud as a skin appendage: unique and shared aspects of development. *J Mammary Gland Biol Neoplasia* 11(3-4):187-203.
- Mikkola ML, Pispa J, Pekkanen M, Paulin L, Nieminen P, Kere J, et al. (1999). Ectodysplasin, a protein required for epithelial morphogenesis, is a novel TNF homologue and promotes cell-matrix adhesion. *Mech Dev* 88(2):133-46.
- Mill P, Mo R, Fu H, Grachtchouk M, Kim PC, Dlugosz AA, et al. (2003). Sonic hedgehog-dependent activation of Gli2 is essential for embryonic hair follicle development. *Genes Dev* 17(2):282-94.

- Millar SE (2002). Molecular mechanisms regulating hair follicle development. *J Invest Dermatol* 118(2):216-25.
- Monreal AW, Zonana J, Ferguson B (1998). Identification of a new splice form of the EDA1 gene permits detection of nearly all X-linked hypohidrotic ectodermal dysplasia mutations. *Am J Hum Genet* 63(2):380-9.
- Monreal AW, Ferguson BM, Headon DJ, Street SL, Overbeek PA, Zonana J (1999). Mutations in the human homologue of mouse dl cause autosomal recessive and dominant hypohidrotic ectodermal dysplasia. *Nat Genet* 22(4):366-9.
- Morlon A, Munnich A, Smahi A (2005). TAB2, TRAF6 and TAK1 are involved in NF-kappaB activation induced by the TNF-receptor, Edar and its adaptator Edaradd. *Hum Mol Genet* 14(23):3751-7.
- Mostowska A, Biedziak B, Trzeciak WH (2006). A novel c.581C>T transition localized in a highly conserved homeobox sequence of MSX1: is it responsible for oligodontia? *J Appl Genet* 47(2):159-64.
- Mou C, Jackson B, Schneider P, Overbeek PA, Headon DJ (2006). Generation of the primary hair follicle pattern. *Proc Natl Acad Sci U S A* 103(24):9075-80.
- Munoz F, Lestringant G, Sybert V, Frydman M, Alswaini A, Frossard PM, et al. (1997). Definitive evidence for an autosomal recessive form of hypohidrotic ectodermal dysplasia clinically indistinguishable from the more common X-linked disorder. *Am J Hum Genet* 61(1):94-100.
- Mustonen T, Pispa J, Mikkola ML, Pummila M, Kangas AT, Pakkasjarvi L, et al. (2003). Stimulation of ectodermal organ development by Ectodysplasin-A1. *Dev Biol* 259(1):123-36.

- Mustonen T, Ilmonen M, Pummila M, Kangas AT, Laurikkala J, Jaatinen R, et al. (2004). Ectodysplasin A1 promotes placodal cell fate during early morphogenesis of ectodermal appendages. *Development* 131(20):4907-19.
- Naito A, Yoshida H, Nishioka E, Satoh M, Azuma S, Yamamoto T, et al. (2002). TRAF6-deficient mice display hypohidrotic ectodermal dysplasia. *Proc Natl Acad Sci U S A* 99(13):8766-71.
- Ohazama A, Courtney JM, Sharpe PT (2003). Expression of TNF-receptor-associated factor genes in murine tooth development. *Gene Expr Patterns* 3(2):127-9.
- Paakkonen K, Cambiaghi S, Novelli G, Ouzts LV, Penttinen M, Kere J, et al. (2001). The mutation spectrum of the EDA gene in X-linked anhidrotic ectodermal dysplasia. *Hum Mutat* 17(4):349.
- Pantalacci S, Chaumot A, Benoit G, Sadier A, Delsuc F, Douzery EJ, et al. (2008). Conserved features and evolutionary shifts of the EDA signaling pathway involved in vertebrate skin appendage development. *Mol Biol Evol* 25(5):912-28.
- Perou CM, Perchellet A, Jago T, Pryor R, Kaplan J, Justice MJ (1997). Comparative mapping in the beige-satin region of mouse chromosome 13. *Genomics* 39(2):136-46.
- Pinheiro M, Freire-Maia N (1994). Ectodermal dysplasias: a clinical classification and a causal review. *Am J Med Genet* 53(2):153-62.
- Pispa J, Jung HS, Jernvall J, Kettunen P, Mustonen T, Tabata MJ, et al. (1999). Cusp patterning defect in Tabby mouse teeth and its partial rescue by FGF. *Dev Biol* 216(2):521-34.

- Pispa J, Thesleff I (2003). Mechanisms of ectodermal organogenesis. *Dev Biol* 262(2):195-205.
- Priolo M, Lagana C (2001). Ectodermal dysplasias: a new clinical-genetic classification. *J Med Genet* 38(9):579-85.
- Prodi DA, Pirastu N, Maninchetta G, Sassu A, Picciau A, Palmas MA, et al. (2008). EDA2R is associated with androgenetic alopecia. *J Invest Dermatol* 128(9):2268-70.
- Pummila M, Fliniaux I, Jaatinen R, James MJ, Laurikkala J, Schneider P, et al. (2007). Ectodysplasin has a dual role in ectodermal organogenesis: inhibition of Bmp activity and induction of Shh expression. *Development* 134(1):117-25.
- Rasool M, Schuster J, Aslam M, Tariq M, Ahmad I, Ali A, et al. (2008). A novel missense mutation in the EDA gene associated with X-linked recessive isolated hypodontia. *J Hum Genet* 53(10):894-8.
- Schmidt-Ullrich R, Paus R (2005). Molecular principles of hair follicle induction and morphogenesis. *Bioessays* 27(3):247-61.
- Schmidt-Ullrich R, Tobin DJ, Lenhard D, Schneider P, Paus R, Scheidereit C (2006). NF-kappaB transmits Eda A1/EdaR signalling to activate Shh and cyclin D1 expression, and controls post-initiation hair placode down growth. *Development* 133(6):1045-57.
- Schneider P, Street SL, Gaide O, Hertig S, Tardivel A, Tschoop J, et al. (2001). Mutations leading to X-linked hypohidrotic ectodermal dysplasia affect three major functional domains in the tumor necrosis factor family member ectodysplasin-A. *J Biol Chem* 276(22):18819-27.

- Shimomura Y, Sato N, Miyashita A, Hashimoto T, Ito M, Kuwano R (2004). A rare case of hypohidrotic ectodermal dysplasia caused by compound heterozygous mutations in the EDAR gene. *J Invest Dermatol* 123(4):649-55.
- Sofaer JA (1981). Hypodontia and sweat pore counts in detecting carriers of X-linked hypohidrotic ectodermal dysplasia. *Br Dent J* 151(10):327-30.
- Song S, Han D, Qu H, Gong Y, Wu H, Zhang X, et al. (2009). EDA gene mutations underlie non-syndromic oligodontia. *J Dent Res* 88(2):126-31.
- Srivastava AK, Pispa J, Hartung AJ, Du Y, Ezer S, Jenks T, et al. (1997). The Tabby phenotype is caused by mutation in a mouse homologue of the EDA gene that reveals novel mouse and human exons and encodes a protein (ectodysplasin-A) with collagenous domains. *Proc Natl Acad Sci U S A* 94(24):13069-74.
- Stevenson AC, Kerr CB (1967). On the distribution of frequencies of mutation to genes determining harmful traits in man. *Mutat Res* 4(3):339-52.
- Takeda K, Takeuchi O, Tsujimura T, Itami S, Adachi O, Kawai T, et al. (1999). Limb and skin abnormalities in mice lacking IKKalpha. *Science* 284(5412):313-6.
- Tanikawa C, Furukawa Y, Yoshida N, Arakawa H, Nakamura Y, Matsuda K (2009). XEDAR as a putative colorectal tumor suppressor that mediates p53-regulated anoikis pathway. *Oncogene* 28(34):3081-92.
- Tao R, Jin B, Guo SZ, Qing W, Feng GY, Brooks DG, et al. (2006). A novel missense mutation of the EDA gene in a Mongolian family with congenital hypodontia. *J Hum Genet* 51(5):498-502.
- Tarpey P, Pemberton TJ, Stockton DW, Das P, Ninis V, Edkins S, et al. (2007). A novel Gln358Glu mutation in ectodysplasin A associated with X-linked dominant incisor hypodontia. *Am J Med Genet A* 143(4):390-4.

- Thesleff I, Mikkola M (2002). The role of growth factors in tooth development. *Int Rev Cytol* 217:93-135.
- Thurnam J (1848). Two cases in which the skin, hair and teeth were very imperfectly developed. *Med Chir Trans* 3:71-82.
- Tucker AS, Headon DJ, Schneider P, Ferguson BM, Overbeek P, Tschopp J, et al. (2000). Edar/Eda interactions regulate enamel knot formation in tooth morphogenesis. *Development* 127(21):4691-700.
- Tucker A, Sharpe P (2004). The cutting-edge of mammalian development; how the embryo makes teeth. *Nat Rev Genet* 5(7):499-508.
- van den Boogaard MJ, Dorland M, Beemer FA, van Amstel HK (2000). MSX1 mutation is associated with orofacial clefting and tooth agenesis in humans. *Nat Genet* 24(4):342-3.
- van der Hout AH, Oudesluijs GG, Venema A, Verheij JB, Mol BG, Rump P, et al. (2008). Mutation screening of the Ectodysplasin-A receptor gene EDAR in hypohidrotic ectodermal dysplasia. *Eur J Hum Genet* 16(6):673-9.
- Vincent MC, Biancalana V, Ginisty D, Mandel JL, Calvas P (2001). Mutational spectrum of the ED1 gene in X-linked hypohidrotic ectodermal dysplasia. *Eur J Hum Genet* 9(5):355-63.
- Veltmaat JM, Mailleux AA, Thiery JP, Bellusci S (2003). Mouse embryonic mammogenesis as a model for the molecular regulation of pattern formation. *Differentiation* 71(1):1-17.
- Veltmaat JM, Van Veelen W, Thiery JP, Bellusci S (2004). Identification of the mammary line in mouse by Wnt10b expression. *Dev Dyn* 229(2):349-56.

- Yamamoto Y, Kim DW, Kwak YT, Prajapati S, Verma U, Gaynor RB (2001). IKKgamma /NEMO facilitates the recruitment of the IkappaB proteins into the IkappaB kinase complex. *J Biol Chem* 276(39):36327-36.
- Yan M, Wang LC, Hymowitz SG, Schilbach S, Lee J, Goddard A, et al. (2000). Two-amino acid molecular switch in an epithelial morphogen that regulates binding to two distinct receptors. *Science* 290(5491):523-7.
- Yan M, Zhang Z, Brady JR, Schilbach S, Fairbrother WJ, Dixit VM (2002). Identification of a novel death domain-containing adaptor molecule for ectodysplasin-A receptor that is mutated in crinkled mice. *Curr Biol* 12(5):409-13.
- Zhang H, Quan C, Sun LD, Lv HL, Gao M, Zhou FS, et al. (2009). A novel frameshift mutation of the EDA1 gene in a Chinese Han family with X-linked hypohidrotic ectodermal dysplasia. *Clin Exp Dermatol* 34(1):74-6.
- Zhang J, Han D, Song S, Wang Y, Zhao H, Pan S, et al. (2011). Correlation between the phenotypes and genotypes of X-linked hypohidrotic ectodermal dysplasia and non-syndromic hypodontia caused by ectodysplasin-A mutations. *Eur J Med Genet* 54(4):e377-82.
- Zonana J (1993). Hypohidrotic (anhidrotic) ectodermal dysplasia: molecular genetic research and its clinical applications. *Semin Dermatol* 12(3):241-6.