

BIBLIOGRAPHY

1. Krishnan V, Davidovitch Z. Cellular, molecular, and tissue-level reactions to orthodontic force. *Am J Orthod Dentofacial Orthop* 2006;129:469 e461-432.
2. Melsen B. Biological reaction of alveolar bone to orthodontic tooth movement. *Angle Orthod* 1999;69:151-158.
3. Christenson RH. Biochemical markers of bone metabolism: an overview. *Clin Biochem* 1997;30:573-593.
4. Lowney JJ, Norton LA, Shafer DM, Rossomando EF. Orthodontic forces increase tumor necrosis factor alpha in the human gingival sulcus. *Am J Orthod Dentofacial Orthop* 1995;108:519-524.
5. Uematsu S, Mogi M, Deguchi T. Interleukin (IL)-1 beta, IL-6, tumor necrosis factor-alpha, epidermal growth factor, and beta 2-microglobulin levels are elevated in gingival crevicular fluid during human orthodontic tooth movement. *J Dent Res* 1996;75:562-567.
6. Dedic A, Kiliaridis S, Mombelli A, Giannopoulou C. Composition changes in gingival crevicular fluid during orthodontic tooth movement: comparisons between tension and compression sides. *Eur J Oral Sci* 2006;114:416-422.
7. Yamaguchi M, Yoshii M, Kasai K. Relationship between substance P and interleukin-1beta in gingival crevicular fluid during orthodontic tooth movement in adults. *Eur J Orthod*. 2006;28:241-246.

8. Insoft M, King GJ, Keeling SD. The measurement of acid and alkaline phosphatase in gingival crevicular fluid during orthodontic tooth movement. *Am J Orthod Dentofacial Orthop* 1996;109:287-296.
9. Perinetti G, Paolantonio M, D'Attilio M, D'Archivio D, Tripodi D, Femminella B et al. Alkaline phosphatase activity in gingival crevicular fluid during human orthodontic tooth movement. *Am J Orthod Dentofacial Orthop* 2002;122:548-556.
10. Nozoe T, Yamaguchi I M, Nogimura A, Aihara N, Kojima T, Kasai K. The level of cathepsin L in gingival crevicular fluid during orthodontic tooth movement and effect of tension-force on cathepsin L activity from periodontal ligament cells. *Orthodontic Waves* 2002;61:447-453.
11. Villela B, Cogen RB, Bartolucci AA, Birkedal-Hansen H. Crevicular fluid collagenase activity in healthy, gingivitis, chronic adult periodontitis and localized juvenile periodontitis patients. *J Periodontal Res* 1987;22:209-211.
12. Nakashima K, Roehrich N, Cimasoni G. Osteocalcin, prostaglandin E2 and alkaline phosphatase in gingival crevicular fluid: their relations to periodontal status. *J Clin Periodontol* 1994;21:327-333.
13. Champe PC, Harvey RA. Glycosaminoglycans Lippincott's Illustrated Reviews: Biochemistry Philadelphia, PA: J. B. Lippincott Company; 1994. p. 147-156.
14. Waddington RJ, Embrey G, Last KS. Glycosaminoglycans of human alveolar bone. *Arch Oral Biol* 1989;34:587-589.
15. Pender N, Samuels RH, Last KS. The monitoring of orthodontic tooth movement over a 2-year period by analysis of gingival crevicular fluid. *Eur J Orthod* 1994;16:511-520.

16. Samuels RH, Pender N, Last KS. The effects of orthodontic tooth movement on the glycosaminoglycan components of gingival crevicular fluid. *J Clin Periodontol* 1993;20:371-377.
17. Last KS, Donkin C, Embrey G. Glycosaminoglycans in human gingival crevicular fluid during orthodontic movement. *Arch Oral Biol* 1988;33:907-912.
18. Waddington RJ, Embrey G, Samuels RH. Characterization of proteoglycan metabolites in human gingival crevicular fluid during orthodontic tooth movement. *Arch Oral Biol* 1994;39:361-368.
19. Baldwin PD, Pender N, Last KS. Effects on tooth movement of force delivery from nickel-titanium archwires. *Eur J Orthod* 1999;21:481-489.
20. Kagayama M, Sasano Y, Mizoguchi I, Kamo N, Takahashi I, Mitani H. Localization of glycosaminoglycans in periodontal ligament during physiological and experimental tooth movement. *J Periodontal Res* 1996;31:229-234.
21. Peansukmanee S, Trinarong C, Kongtawelert P, Ong-chai S. Chondroitin sulfate epitopes in sera of normal and osteoarthritic horses. *Chiang Mai Veterinary Journal* 2003;1:3-10.
22. Pothacharoen P, Kalayanamitra K, Deepa SS, Fukui S, Hattori T, Fukushima N et al. Two related but distinct chondroitin sulfate mimotope octasaccharide sequences recognized by monoclonal antibody WF6. *J Biol Chem* 2007;282:35232-35246.

23. Pothacharoen P, Teekachunhatean S, Louthrenoo W, Yingsung W, Ong-Chai S, Hardingham T et al. Raised chondroitin sulfate epitopes and hyaluronan in serum from rheumatoid arthritis and osteoarthritis patients. *Osteoarthritis Cartilage* 2006;14:299-301.
24. Jaito N, Jotikasthira D, Krisanaprakornkit S, Ong-chai S, Kongtawelert P. Monitoring of chondroitin-6-sulfate levels in gingival crevicular fluid during orthodontic canine movement. In: In:Davidovitch Z, Mah J, Suthanarak S, editors. *Biological Mechanisms of Tooth Eruption, Resorption and Movement*. Boston Harvard Society for the Advancement of Orthodontics, Boston, Massachusetts; 2006. p. 289-295.
25. King GJ, Keeling SD, McCoy EA, Ward TH. Measuring dental drift and orthodontic tooth movement in response to various initial forces in adult rats. *Am J Orthod Dentofacial Orthop* 1991;99:456-465.
26. Murrell EF, Yen EH, Johnson RB. Vascular changes in the periodontal ligament after removal of orthodontic forces. *Am J Orthod Dentofacial Orthop* 1996;110:280-286.
27. von Bohl M, Maltha JC, Von Den Hoff JW, Kuijpers-Jagtman AM. Focal hyalinization during experimental tooth movement in beagle dogs. *Am J Orthod Dentofacial Orthop* 2004;125:615-623.
28. Hill PA. Bone remodeling. *British Journal of Orthodontics* 1998;25:101-107.
29. Gonzales C, Hotokezaka H, Yoshimatsu M, Yozgatian JH, Darendeliler MA, Yoshida N. Force magnitude and duration effects on amount of tooth movement and root resorption in the rat molar. *Angle Orthod* 2008;78:502-509.

30. Schwarz AM. Tissue changes incidental to orthodontic tooth movement. International Journal of Orthodontics, Oral Surgery and Radiography 1932;18:331-352.
31. Proffit WR, Fields HW, Sarver DM. Contemporary Orthodontics The Biologic Basis of Orthodontic Therapy. St. Louis, Missouri: MOSBY Elsevier; 2007. p. 340.
32. Kohno T, Matsumoto Y, Kanno Z, Warita H, Soma K. Experimental tooth movement under light orthodontic forces: rates of tooth movement and changes of the periodontium. J Orthod 2002;29:129-135.
33. Gonzales C, Hotokezaka H, Arai Y, Ninomiya T, Tominaga J, Jang I et al. An in vivo 3D micro-CT evaluation of tooth movement after the application of different force magnitudes in rat molar. Angle Orthod 2009;79:703-714.
34. Ren Y, Maltha JC, Kuijpers-Jagtman AM. Optimum force magnitude for orthodontic tooth movement: a systematic literature review. Angle Orthod 2003;73:86-92.
35. Boester CH, Johnston LE. A clinical investigation of the concepts of differential and optimal force in canine retraction. Angle Orthod 1974;44:113-119.
36. Owman-Moll P, Kurol J, Lundgren D. Effects of a doubled orthodontic force magnitude on tooth movement and root resorptions. An inter-individual study in adolescents. Eur J Orthod 1996;18:141-150.
37. Owman-Moll P, Kurol J, Lundgren D. The effects of a four-fold increased orthodontic force magnitude on tooth movement and root resorptions. An intra-individual study in adolescents. Eur J Orthod 1996;18:287-294.

38. Thiruvenkatachari B, Ammayappan P, Kandaswamy R. Comparison of rate of canine retraction with conventional molar anchorage and titanium implant anchorage. *Am J Orthod Dentofacial Orthop* 2008;134:30-35.
39. Batra P, Kharbanda O, Duggal R, Singh N, Parkash H. Alkaline phosphatase activity in gingival crevicular fluid during canine retraction. *Orthod Craniofac Res* 2006;9:44-51.
40. Yee JA, Turk T, Elekdag-Turk S, Cheng LL, Darendeliler MA. Rate of tooth movement under heavy and light continuous orthodontic forces. *Am J Orthod Dentofacial Orthop* 2009;136:150 e151-159; discussion 150-151.
41. Lundgren D, Owman-Moll P, Kurol J, Martensson B. Accuracy of orthodontic force and tooth movement measurements. *Br J Orthod* 1996;23:241-248.
42. Kojima Y, Fukui H. Numerical simulation of canine retraction by sliding mechanics. *Am J Orthod Dentofacial Orthop* 2005;127:542-551.
43. Kojima Y, Mizuno T, Umemura S, Fukui H. A numerical simulation of orthodontic tooth movement produced by a canine retraction spring. *Dent Mater J* 2007;26:561-567.
44. Hayashi K, Uechi J, Lee SP, Mizoguchi I. Three-dimensional analysis of orthodontic tooth movement based on XYZ and finite helical axis systems. *Eur J Orthod* 2007;29:589-595.
45. Alfano MC. The origin of gingival fluid. *J Theor Biol* 1974;47:127-136.
46. Lee AJ, Walsh TF, Hodges SJ, Rawlinson A. Gingival crevicular fluid osteocalcin in adult periodontitis. *J Clin Periodontol* 1999;26:252-256.

47. Tsai CC, Ho YP, Chen CC. Levels of interleukin-1 beta and interleukin-8 in gingival crevicular fluids in adult periodontitis. *J Periodontol* 1995;66:852-859.
48. Chapple IL, Garner I, Saxby MS, Moscrop H, Matthews JB. Prediction and diagnosis of attachment loss by enhanced chemiluminescent assay of crevicular fluid alkaline phosphatase levels. *J Clin Periodontol* 1999;26:190-198.
49. Armitage GC. Analysis of gingival crevice fluid and risk of progression of periodontitis. *Periodontol* 2000 2004;34:109-119.
50. Kavadia-Tsatala S, Kaklamanos EG, Tsalikis L. Effects of orthodontic treatment on gingival crevicular fluid flow rate and composition: clinical implications and applications. *Int J Adult Orthodon Orthognath Surg* 2002;17:191-205.
51. Curtis MA, Griffiths GS, Price SJ, Coulthurst SK, Johnson NW. The total protein concentration of gingival crevicular fluid. Variation with sampling time and gingival inflammation. *J Clin Periodontol* 1988;15:628-632.
52. Haffajee AD, Socransky SS. Microbial etiological agents of destructive periodontal diseases. *Periodontol* 2000 1994;5:78-111.
53. Eley BM, Cox SW. Proteolytic and hydrolytic enzymes from putative periodontal pathogens: characterization, molecular genetics, effects on host defenses and tissues and detection in gingival crevicle fluid. *Periodontol* 2000 2003;31:105-124.

54. Leethanakul C, Kittichaikarn C, Charoemratrete C, Jitpukdeebodintra S. Effects of continuous and interrupted orthodontic force on interleukin-1beta and interleukin-8 secretion in human gingival crevicular fluid. *Journal of Oral Biosciences* 2008;50:230-238.
55. Grieve WG, 3rd, Johnson GK, Moore RN, Reinhardt RA, DuBois LM. Prostaglandin E (PGE) and interleukin-1 beta (IL-1 beta) levels in gingival crevicular fluid during human orthodontic tooth movement. *Am J Orthod Dentofacial Orthop* 1994;105:369-374.
56. Clark DA, Coker R. Transforming growth factor-beta (TGF-beta). *Int J Biochem Cell Biol* 1998;30:293-298.
57. Janssens K, ten Dijke P, Janssens S, Van Hul W. Transforming growth factor-beta1 to the bone. *Endocr Rev* 2005;26:743-774.
58. Uematsu S, Mogi M, Deguchi T. Increase of transforming growth factor-beta 1 in gingival crevicular fluid during human orthodontic tooth movement. *Arch Oral Biol* 1996;41:1091-1095.
59. Lamster IB, Holmes LG, Gross KB, Oshrain RL, Cohen DW, Rose LF et al. The relationship of beta-glucuronidase activity in crevicular fluid to probing attachment loss in patients with adult periodontitis. Findings from a multicenter study. *J Clin Periodontol* 1995;22:36-44.
60. Kunimatsu K, Yamamoto K, Ichimaru E, Kato Y, Kato I. Cathepsins B, H and L activities in gingival crevicular fluid from chronic adult periodontitis patients and experimental gingivitis subjects. *J Periodontal Res* 1990;25:69-73.

61. Tsuji Y, Yamaza T, Kido MA, Goto T, Nakata S, Akamine A et al. Expression of cathepsin K mRNA and protein in odontoclasts after experimental tooth movement in the mouse maxilla by in situ hybridization and immunoelectron microscopy. *Cell Tissue Res* 2001;303:359-369.
62. Strbac GD, Monov G, Cei S, Kandler B, Watzek G, Gruber R. Cathepsin K levels in the crevicular fluid of dental implants: a pilot study. *J Clin Periodontol* 2006;33:302-308.
63. Alpagot T, Silverman S, Lundergan W, Bell C, Chambers DW. Crevicular fluid elastase levels in relation to periodontitis and metabolic control of diabetes. *J Periodontal Res* 2001;36:169-174.
64. Plagnat D, Giannopoulou C, Carrel A, Bernard JP, Mombelli A, Belser UC. Elastase, alpha2-macroglobulin and alkaline phosphatase in crevicular fluid from implants with and without periimplantitis. *Clin Oral Implants Res* 2002;13:227-233.
65. Bull H, Murray PG, Thomas D, Fraser AM, Nelson PN. Acid phosphatases. *Mol Pathol* 2002;55:65-72.
66. Ljusberg J, Wang Y, Lang P, Norgard M, Dodds R, Hultenby K et al. Proteolytic excision of a repressive loop domain in tartrate-resistant acid phosphatase by cathepsin K in osteoclasts. *J Biol Chem* 2005;280:28370-28381.
67. Halleen JM, Tiitinen SL, Ylipahkala H, Fagerlund KM, Vaananen HK. Tartrate-resistant acid phosphatase 5b (TRACP 5b) as a marker of bone resorption. *Clin Lab* 2006;52:499-509.

68. Calogero D, Scott Malone J, Wheeler TT. Current Concepts in the Biology of Orthodontic Tooth Movement. *Seminars in Orthodontics* 2002;8:6-12.
69. van de Wijngaert FP, Burger EH. Demonstration of tartrate-resistant acid phosphatase in un-decalcified, glycolmethacrylate-embedded mouse bone: a possible marker for (pre)osteoclast identification. *J Histochem Cytochem* 1986;34:1317-1323.
70. Cole AA, Walters LM. Tartrate-resistant acid phosphatase in bone and cartilage following decalcification and cold-embedding in plastic. *J Histochem Cytochem* 1987;35:203-206.
71. Rody WJ, Jr., King GJ, Gu G. Osteoclast recruitment to sites of compression in orthodontic tooth movement. *Am J Orthod Dentofacial Orthop* 2001;120:477-489.
72. Igarashi Y, Lee MY, Matsuzaki S. Acid phosphatases as markers of bone metabolism. *J Chromatogr B Analyt Technol Biomed Life Sci* 2002;781:345-358.
73. Lau KH, Onishi T, Wergedal JE, Singer FR, Baylink DJ. Characterization and assay of tartrate-resistant acid phosphatase activity in serum: potential use to assess bone resorption. *Clin Chem* 1987;33:458-462.
74. Price CP, Kirwan A, Vader C. Tartrate-resistant acid phosphatase as a marker of bone resorption. *Clin Chem* 1995;41:641-643.
75. Qin YJ, Zhang ZL, Zhang H, Hu WW, Liu YJ, Hu YQ et al. Age-related changes of serum tartrate-resistant acid phosphatase 5b and the relationship with bone mineral density in Chinese women. *Acta Pharmacol Sin* 2008;29:1493-1498.

76. Janckila AJ, Cardwell EM, Yam LT, Li CY. Hairy cell identification by immunohistochemistry of tartrate-resistant acid phosphatase. *Blood* 1995;85:2839-2844.
77. Terpos E, de la Fuente J, Szydlo R, Hatjiharissi E, Viniou N, Meletis J et al. Tartrate-resistant acid phosphatase isoform 5b: a novel serum marker for monitoring bone disease in multiple myeloma. *Int J Cancer* 2003;106:455-457.
78. Chung YC, Ku CH, Chao TY, Yu JC, Chen MM, Lee SH. Tartrate-resistant acid phosphatase 5b activity is a useful bone marker for monitoring bone metastases in breast cancer patients after treatment. *Cancer Epidemiol Biomarkers Prev* 2006;15:424-428.
79. Waddington RJ, Embery G. Proteoglycans and orthodontic tooth movement. *J Orthod* 2001;28:281-290.
80. Beck CB, Embery G, Langley MS, Waddington RJ. Levels of glycosaminoglycans in peri-implant sulcus fluid as a means of monitoring bone response to endosseous dental implants. *Clin Oral Implants Res* 1991;2:179-185.
81. Smith AJ, Addy M, Embery G. Gingival crevicular fluid glycosaminoglycan levels in patients with chronic adult periodontitis. *J Clin Periodontol* 1995;22:355-361.
82. Intachai I, Krisanaprakornkit S, Kongtawelert P, Ong-chai S, Buranastidporn B, Suzuki EY et al. Chondroitin sulphate (WF6 epitope) levels in peri-miniscrew implant crevicular fluid during orthodontic loading. *Eur J Orthod* 2009;32:60-65.

83. Peeranantrangsri A, Jotikasthira D, Krisanaprakornkit S, Ong-chai S, Kongtawelert P, Jotikasthira N. Monitoring changes in hyaluronic acid levels in gingival crevicular fluid during orthodontic canine movement. In: In: Davidovitch Z, Mah J, Suthanarak S, editors. *Biological Mechanisms of Tooth Eruption, Resorption and Movement*. Boston: Harvard Society for the Advancement of Orthodontics, Boston, Massachusetts; 2006. p. 283-288.
84. Talonpoika J, Heino J, Larjava H, Hakkinen L, Paunio K. Gingival crevicular fluid fibronectin degradation in periodontal health and disease. *Scand J Dent Res* 1989;97:415-421.
85. Kunimatsu K, Mataki S, Tanaka H, Mine N, Kiyoki M, Hosoda K et al. A cross-sectional study on osteocalcin levels in gingival crevicular fluid from periodontal patients. *J Periodontol* 1993;64:865-869.
86. Kido J, Nakamura T, Asahara Y, Sawa T, Kohri K, Nagata T. Osteopontin in gingival crevicular fluid. *J Periodontal Res* 2001;36:328-333.
87. Figueiredo CM, Gustafsson A. Increased amounts of laminin in GCF from untreated patients with periodontitis. *J Clin Periodontol* 2000;27:313-318.
88. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976;72:248-254.
89. Bartold PM, Miki Y, McAllister B, Narayanan AS, Page RC. Glycosaminoglycans of human cementum. *J Periodontal Res* 1988;23:13-17.
90. Bartold PM. Proteoglycans of the periodontium: structure, role and function. *J Periodontal Res* 1987;22:431-444.

91. Gibson GJ, Pearson CH. Sulfated galactosaminoglycans of bovine periodontal ligament. Evidence for the presence of two major types of hybrids but no chondroitin sulfate. *Connect Tissue Res* 1982;10:161-171.
92. Pearson CH, Gibson GJ. Proteoglycans of bovine periodontal ligament and skin. Occurrence of different hybrid-sulphated galactosaminoglycans in distinct proteoglycans. *Biochem J* 1982;201:27-37.
93. Ren Y, Maltha JC, Van 't Hof MA, Kuijpers-Jagtman AM. Optimum force magnitude for orthodontic tooth movement: a mathematic model. *Am J Orthod Dentofacial Orthop* 2004;125:71-77.
94. Luppanapornlarp S, Kajii TS, Surarit R, Iida J. Interleukin-1 β levels, pain intensity, and tooth movement using two different magnitudes of continuous orthodontic force. *Eur J Orthod* 2010;32:596-601.
95. Bodian CA, Freedman G, Hossain S, Eisenkraft JB, Beilin Y. The visual analog scale for pain: clinical significance in postoperative patients. *Anesthesiology* 2001;95:1356-1361.
96. DeLoach LJ, Higgins MS, Caplan AB, Stiff JL. The visual analog scale in the immediate postoperative period: intrasubject variability and correlation with a numeric scale. *Anesth Analg* 1998;86:102-106.
97. Ogura M, Kamimura H, Al-Kalaly A, Nagayama K, Taira K, Nagata J et al. Pain intensity during the first 7 days following the application of light and heavy continuous forces. *Eur J Orthod* 2009;31:314-319.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved