

## TABLE OF CONTENTS

<b>ACKNOWLEDGEMENTS</b>	iii
<b>ABSTRACT (ENGLISH)</b>	iv
<b>ABSTRACT (THAI)</b>	vi
<b>TABLE OF CONTENTS</b>	viii
<b>LIST OF TABLES</b>	xii
<b>LIST OF ILLUSTRATIONS</b>	xiv
<b>CHAPTER 1: INTRODUCTION</b>	
1.1 Principles, rationale and hypothesis	1
1.2 Research objectives	1
1.3 Research scope	2
1.4 Usefulness of the research	2
1.5 Research locations	2
<b>CHAPTER 2: LITERATURE REVIEW</b>	
2.1 Introduction	3
2.2 Fruit morphology	4
2.3 Anatomy of longan fruit pericarp	4
2.4 Thai longan cultivar	5
2.4.1 Daw	5
2.4.2 Biew Kiew	5
2.4.3 Si Chomphu	5
2.4.4 Haeo	5
2.5 Fruit growth	6
2.6 Harvest maturity	6
2.7 Harvesting method	7
2.8 Postharvest physiology	7

2.8.1	Respiration rate of longan fruit	7
2.8.2	Ethylene production of longan fruit	8
2.9	Chemical composition	9
2.10	Postharvest storage	10
2.11	Chilling injury	12
2.11.1	Chilling temperature of longan fruit	12
2.11.2	Symptoms of chilling injury	13
2.11.3	Mechanism of chilling injury	13
2.12	Phenolic compounds in fruits	14
2.13	Enzymatic browning of longan pericarp	18
2.14	Plant cell wall components	23
<b>CHAPTER 3: MATERIALS AND METHODS</b>		
3.1	Anatomical comparison between normal and chilling injured longan fruit pericarps cv. Daw and Biew Kiew	25
3.1.1	Fruit materials	25
3.1.2	Chilling injury evaluation	25
3.1.3	Anatomical of longan fruit pericarps	26
3.2	Physico-chemical changes between normal and chilling injured longan fruit pericarps	28
3.2.1	Measurement of longan fruit pericarp color	28
3.2.2	Determination of moisture content of longan fruit pericarp	28
3.2.3	Determination of electrolyte leakage	28
3.3	The relationship between polyphenol oxidase activity and electrolyte leakage of longan fruit pericarp during chilling injured of longan cv. Daw	29
3.4	Identification the main classes of phenolic compounds and other components of normal and chilling injured longan fruit pericarps	31
3.4.1	Identification of phenolic compounds in longan fruit pericarp	31


3.4.2 Quantification of phenolic compounds in longan fruit pericarp	33
<b>3.5 Quantification of other components of normal and chilling injured longan fruit pericarps cv. Daw and Biew Kiew</b>	<b>34</b>
3.5.1 Plant materials	34
3.5.2 Quantification of pectin and lignin	35
3.5.3 Quantification of total dietary fiber (TDF)	39
<b>3.6 Statistical data analysis</b>	<b>42</b>
 <b>CHAPTER 4: RESULTS AND DISCUSSION</b>	
<b>4.1 Anatomical comparison between normal and chilling injured longan fruit pericarps cv. Daw and Biew Kiew</b>	<b>43</b>
<b>4.1.1 The anatomy of normal longan fruit pericarp</b>	<b>43</b>
4.1.1.1 Visual observation of longan fruit pericarp	43
4.1.1.2 Stereomicroscope observation of longan fruit pericarp	43
4.1.1.3 Microscopic anatomy and ultrastructure of longan fruit pericarp	43
4.1.1.4 Anatomy of inner surface longan fruit pericarp	45
<b>4.1.2 The anatomy of chilling injured longan fruit pericarp</b>	<b>54</b>
4.1.2.1 Visual observation of chilling injured longan fruit pericarp	54
4.1.2.2 Stereomicroscope observation of chilling injured longan fruit pericarp	54
4.1.2.3 Microscopic anatomy and ultrastructure of chilling injured longan fruit pericarp	55
<b>4.2 Other morphology and anatomy of longan fruit</b>	<b>64</b>
4.2.1 Morphology and anatomy of longan aril	64
4.2.2 The porosity of longan fruit	64
<b>4.3 Physico-chemical changes between normal and chilling injured longan fruit pericarps</b>	<b>67</b>
4.3.1 Measurement of the pericarp color	67
4.3.2 Determination of moisture content of pericarp	67

4.3.3 Determination of electrolyte leakage	68
4.4 The relationship between polyphenol oxidase activity and electrolyte leakage of longan pericarp during chilling injury of longan fruit cv. Daw	73
4.5 To identify the main classes of phenolic compounds and other components of normal and chilling injured longan fruit pericarps	76
4.5.1 Number and identification of phenolic compounds in longan fruit pericarps cv. Daw and Biew Kiew	76
4.5.2 Analysis of unknown compounds	84
4.5.3 Quantification of phenolic compounds in longan fruit pericarp	87
4.6 Quantification of other components of normal and chilling injured longan fruit pericarps cv. Daw and Biew Kiew	94
<b>CHAPTER 5: CONCLUSIONS</b>	97
<b>BIBLIOGRAPHY</b>	99
<b>APPENDIX</b>	108
<b>PUBLICATIONS</b>	117
<b>CURRICULUM VITAE</b>	119

## LIST OF TABLES

<b>Table</b>	<b>Page</b>
2.1 Longan fruit compositions at harvest	9
2.2 The chemical compositions and nutritional values of fresh longan fruit	10
2.3 Optimal storage temperature recommendations for longan fruit cultivars	11
2.4 The major classes of phenolics in fruits	18
2.5 Cell wall components	24
4.1 L* values of outer pericarp surface cv. Daw and Biew Kiew during storage at 5 and 10°C	69
4.2 L* values of inner pericarp surface cv. Daw and Biew Kiew during storage at 5 and 10°C	70
4.3 Moisture content of longan pericarps cv. Daw and Biew Kiew during storage at 5 and 10°C	71
4.4 Electrolyte leakage of longan pericarps cv. Daw and Biew Kiew during storage at 5 and 10°C	72
4.5 The L* values, PPO activity and electrolyte leakage of longan pericarp cv. Daw during storage at 5°C for 14 days	75
4.6 Mass spectra of kaempferol glycosides in longan pericarp. Abbreviations are: K, kaempferol; glc, glucose; rha, rhamnose. A pentose is indicated by the neutral loss of 132 amu (150 amu-18 (H <sub>2</sub> O)). The +Na adduct (+Na) is observed as a mass ion 22 amu above the protonated mass ion (M+H) <sup>+</sup> +1. Neutral losses indicated by parentheses	82
4.7 The main class of phenolic compounds conjugates in freeze-dried longan pericarp was identified by HPLC-PDA-MS, including ellagic acid glycosides, quercetin glycosides, kaempferol glycosides and unknown compounds	83
4.8 Quercetin content <sup>1</sup> of freeze-dried longan pericarps cv. Daw and Biew	

	Kiew during storage at 5 and 10°C	91
4.9	Kaempferol content <sup>1</sup> of freeze-dried longan pericarps cv. Daw and Biew Kiew during storage at 5 and 10°C	92
4.10	TDF, pectin and lignin contents of normal and chilling injured longan fruit pericarps cv. Daw and Biew Kiew during storage at 5°C	95



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่  
Copyright© by Chiang Mai University  
All rights reserved

## LIST OF ILLUSTRATIONS

<b>Figure</b>	<b>Page</b>
2.1 Sigmoidal curve of longan fruit growth and development	6
2.2 Respiration rates of longan fruits stored at at 22 and 5°C	8
2.3 Sequences of metabolic events leading from a stress-induced alteration in the properties of membranes to observable macroscopic tissue damage	15
2.4 Biosynthesis pathways of hydroxycinnamic acids, hydroxybenzoic acids and flavonoids	17
2.5 Structure of selected phenolic compounds in fruits	20
2.6 Basic structures of ellagic acid, quercetin and kaempferol	22
3.1 Chilling injury index criteria of inner surface pericarp	26
3.2 Stereomicrograph of whole pericarp was separated into 2 parts, the inner and outer pericarps	30
3.3 The equipment for phenolic compound analysis	34
3.4 The equipment for galacturonic acid assay	38
4.1 The longan fruits cv. Daw and Biew Kiew after harvest	46
4.2 SEM and LM micrographs of the outer longan fruit pericarp surface	47
4.3 Transverse section micrographs of normal longan fruit pericarp cv. Daw (left) and Biew Kiew (right)	48
4.4 Transverse section micrograph of normal longan fruit pericarp cv. Biew Kiew consist of three layers; exocarp, mesocarp, and endocarp	49
4.5 Transverse section micrographs of exocarp layer	50
4.6 LM and TEM micrographs of cells in mesocarp layer	51
4.7 LM and SEM micrographs of vascular tissue	52
4.8 LM and TEM micrographs of endocarp layer	52
4.9 The inner surface of longan pericarp cv. Daw	53
4.10 Stereomicrographs of chilling injury symptoms of longan fruit pericarps cv. Biew Kiew and Daw during storage for 10 and 14 days	

- at 5°C score 4 showed water soaking and browning >2/3 of outer and inner pericarps 57
- 4.11 SEM micrographs of normal (left) and chilling injured (right) outer pericarp surface of longan fruits 58
- 4.12 Transverse sectional micrographs of normal (left) and chilling injured (right) longan fruit pericarps cv. Daw and Biew Kiew 59
- 4.13 Transverse sectional micrographs of normal (left) and chilling injured (right) of longan fruit pericarp cv. Daw 60
- 4.14 Transverse sectional micrographs of normal (left) and chilling injured (right) longan fruit pericarp when viewed with SEM 61
- 4.15 Transverse sectional micrographs of normal (left) and chilling injured (right) longan fruit pericarp when viewed with LM 62
- 4.16 TEM micrographs of normal and chilling injured longan fruit pericarp 63
- 4.17 Longan aril micrographs of longan cv. Daw and Biew Kiew 65
- 4.18 The longan fruit pericarp has numerous pores indicating by air bubbles on the surface of fruit under low pressure (hypobaric condition) 66
- 4.19 The pericarp was separated into 2 parts of whole pericarp for activity analysis showed the inner pericarp (endocarp and lower mesocarp include some part of vascular tissue) and outer pericarp (mid-upper mesocarp and endocarp) when viewed with SEM 74
- 4.20 The large number of peak overlaps in the HPLC chromatograms showed many of the compounds in the longan pericarp cv. Daw unhydrolysis (A) and acid hydrolysis (B) 79
- 4.21 HPLC chromatograms at 330 nm of three P2 column fractions. Ellagic acid conjugates in chromatograms A and B are labeled by EA, and set of related phenolic unknown compounds are labeled with (\*) in chromatogram A. The quercetin glycosides and kaempferol glycosides are labeled by Q and K in chromatogram C, respectively 80



4.22	UV and mass spectra of the major ellagic acid conjugates detected in the freeze-dried normal longan pericarp	80
4.23	UV and mass spectra of quercetin glycosides in freeze-dried longan pericarp	81
4.24	UV and mass spectra of 2 kaempferol glycosides in freeze dried longan pericarp	82
4.25	Analytical silica gel TLC of 3 related unknown compounds. The 3 main unknown compounds, A-C, were applied to 200 $\mu\text{m}$ thickness. Analtech analytical TLC plates with fluorescence indicator. Visualization was made with UV light (365 nm), and with the application of a 5% $\text{H}_2\text{SO}_4$ in ethanol spray followed by heat. No spots in lanes A-C other than the unknown compounds were detected by either UV irradiation or by charring with $\text{H}_2\text{SO}_4$ and heat	86
4.26	UV and mass spectra of 3 related unknown compounds. Mass spectra were recorded with an electrospray ionization cone voltage of +20 V	87
4.27	Fourier transform infrared spectra of unknown B (top) and citrus molasses hydroxycinnamates (bottom). Each spectrum is summations of 8 scans recorded with a Perkin Elmer (Norwalk, Conn.). Spectrum One FTIR spectrophotometer. Samples, dissolved in ethanol were dried onto Real Crystal KBr IR card (International Crystal Laboratories, Garfield, NJ, USA) prior to the FTIR measurements	88
4.28	HPLC chromatograms at 330 nm of hydrolyzed phenolic compounds extracted from longan pericarps cv. Daw (A) and Biew Kiew (B)	90
4.29	Correlation between quercetin and kaempferol contents of longan pericarp cv. Daw and Biew Kiew during storage at 5°C	93
4.30	TDF, pectin and lignin contents of longan pericarps cv. Daw and Biew Kiew between chilling injury score levels during storage at 5°C	96